

Ecological site AX001X01X203

Cryic Aquic Shrubland

Last updated: 5/15/2025
Accessed: 09/26/2026

General information

Provisional. A provisional ecological site description has undergone quality control and quality assurance review. It contains a working state and transition model and enough information to identify the ecological site.

MLRA notes

Major Land Resource Area (MLRA): 001X–Northern Pacific Coast Range, Foothills, and Valleys

This area consists of a long and narrow range of mountains with associated foothills and valleys that parallels the Pacific Ocean. This area is entirely within the Pacific Border Province of the Pacific Mountain System in Oregon and Washington. MLRA 1 is bounded on the north by the highest elevations of the Olympic Mountains and the strait of Juan de Fuca, and by the Klamath Mountains on the south. The Washington portion of this MLRA is primarily composed of young Tertiary sedimentary rocks (siltstone and sandstone) mixed with some volcanic rocks of the same age. Glacial till and outwash deposits are also found in the northern half of this area in Washington. Much of this area is accreted terrane formed by tectonic processes. The average annual precipitation ranges from 60 to 200 inches (1,525 to 5,580 millimeters), increasing with elevation. Most of the precipitation in this area occurs during low-intensity, Pacific frontal storms and is evenly distributed throughout fall, winter, and spring. The dominant soil orders in this MLRA are Andisols, Inceptisols, and Ultisols. Soil depths broadly range from shallow to very deep. Soils are primarily well drained, however poorly drained soils may be found in depressional areas and on alluvial floodplains. Surface textures are typically medial and loamy or clayey. Soils in this area dominantly have a mesic or frigid temperature regime and a udic moisture regime. Soils with aquic moisture regimes and cryic temperature regimes also occur.

Ecological site concept

Cryic Aquic Shrubland sites occur on seeps, depressions, and drains on valley walls, colluvial aprons, and structural benches above 900 meters. These sites are generally located adjacent to moving water. These sites are characterized by an association of shrubs and forbs. Soils on this site are poorly drained and are subject to occasional, brief duration ponding during the growing season. Undergreen willow (*Salix commutata*) and Sitka alder (*Alnus viridis* ssp. *sinuata*) are the most common shrubs on this site. Common forbs include American bistort (*Polygonum bistortoides*), white marsh marigold (*Caltha leptosepala*), Sitka valerian (*Valeriana sitchensis*), common cowparsnip (*Heracleum maximum*), elephanthead lousewort (*Pedicularis groenlandica*), green false hellebore (*Veratrum viride*), and black alpine sedge (*Carex nigricans*).

Associated sites

AX001X01X306	Cryic Aquic Subalpine Wet Meadow Cryic Aquic Subalpine Wet Meadow sites may be found on depressions and seeps adjacent to or surrounded by Cryic Aquic Shrublands. Cryic Aquic Subalpine Wet Meadows are more poorly drained and subject to more frequent ponding for greater duration. Cryic Aquic Subalpine Wet Meadow sites have much lower shrub cover relative to forbs and sedges.
AX001X01X305	Cryic Udic Wet Subalpine Meadow Cryic Udic Wet Subalpine Meadow sites may be found adjacent to Cryic Aquic Shrublands. Cryic Udic Wet Subalpine Meadow sites are less likely to occur adjacent to running water. Cryic Udic Wet Subalpine Meadow sites are not subject to ponding.

Similar sites

AX001X01X001	Temperate Flood Plain Shrubland Temperate Floodplain Shrubland sites are found at lower elevations and will not support subalpine fir (<i>Abies lasiocarpa</i>).
---------------------	--

Table 1. Dominant plant species

Tree	(1) <i>Abies lasiocarpa</i> (2) <i>Callitropsis nootkatensis</i>
Shrub	(1) <i>Salix commutata</i> (2) <i>Alnus viridis ssp. sinuata</i>
Herbaceous	(1) <i>Caltha leptosepala</i> (2) <i>Carex nigricans</i>

Legacy ID

F001XA203WA

Physiographic features

This site primarily occurs on glacial valley walls, colluvial aprons, and structural benches on mountains. Cryic Aquic Shrubland sites are typically located adjacent to streams. These sites receive a significant amount of additional moisture via both run-on and lateral flow.

Table 2. Representative physiographic features

Landforms	(1) Mountains (2) Glacial-valley wall (3) Colluvial apron (4) Structural bench
Flooding frequency	None
Ponding duration	Brief (2 to 7 days)
Ponding frequency	Occasional
Elevation	900 – 1,900 m
Slope	0 – 10 %
Water table depth	150 cm

Aspect	W, NW, N, NE, E, SE, S, SW
--------	----------------------------

Climatic features

This site occurs in a cryic temperature and aquic moisture regime. Precipitation arrives mostly via low-intensity, Pacific frontal storms. Precipitation is unevenly distributed, with the lowest amounts on the leeward side of the Coast Range mountains. Precipitation falls largely as snow in higher elevations. Precipitation is evenly distributed throughout the fall, winter, and spring, while summers are dry. Air temperatures vary significantly along the elevation gradient.

Table 3 Representative climatic features

Frost-free period (characteristic range)	30-90 days
Freeze-free period (characteristic range)	
Precipitation total (characteristic range)	1,500-3,990 mm

Influencing water features

Cryic Aquic Shrubland sites are subject to occasional ponding for brief duration. These sites are frequently found adjacent to streams and creeks. Dominant shrub species are tolerant of saturation and frequent surface disturbance from water erosion.

Soil features

The soils that support this ecological site occur in the cryic soil temperature regime and aquic soil moisture regime. Lakedilly soils are poorly drained, very deep, formed from colluvium from metasedimentary rock, and occur on seeps and depressions on colluvial aprons and glacial valley walls. They have a high or very high saturated hydraulic conductivity. These soils have a seasonally high water table that is within zero to 10 inches of the surface during the growing season, and occasional ponding potential. Lakedilly has greater than 35 percent rock fragments in the control section. Ponding frequency is the limiting factor for plant growth on these soils. Although representative of this site, these soils may exist across multiple ecological sites because of naturally variable slope, texture, rock fragments, and pH. An on-site soil pit and the most current ecological site key are necessary to classify a site.

Table 4. Representative soil features

Parent material	(1) Colluvium – metasedimentary rock
Surface texture	(1) Loam (2) Sandy loam
Drainage class	Poorly drained
Soil depth	150 cm
Surface fragment cover <=3"	0 – 10 %
Surface fragment cover >3"	Not specified

Available water capacity (0-101.6cm)	8.64 – 12.7 cm
Soil reaction (1:1 water) (0-25.4cm)	4.5 – 5.5
Subsurface fragment volume <=3" (0-50.8cm)	0 – 30 %
Subsurface fragment volume >3" (0-50.8cm)	0 – 20 %

Ecological dynamics

The Cryic Aquic Shrubland site is found at mid-to-high elevations in the park where snowpack persists long into the spring and summer. Long-duration snowpack cover significantly reduces growing degree days on site. These sites are typically located adjacent to running water, frequently being found between streams and upland cryic forest or moist meadow sites. Soils on this site are poorly-drained and are subject to occasional, brief-duration ponding. Alpine snowmelt generates large amounts of water and may result in high-intensity floods capable of scouring the soil surface to bare gravels, creating a nutrient-poor substrate. Undergreen willow (*Salix commutata*) and Sitka alder (*Alnus viridis* ssp. *sinuata*) are the most common shrubs on this site. Common forbs include American bistort (*Polygonum bistortoides*), white marsh marigold (*Caltha leptosepala*), Sitka valerian (*Valeriana sitchensis*), common cowparsnip (*Heracleum maximum*), elephanthead lousewort (*Pedicularis groenlandica*), green false hellebore (*Veratrum viride*), and black alpine sedge (*Carex nigricans*).

In the immediate aftermath of a surface-scouring flooding event, undergreen willow (*Salix commutata*) is generally the first shrub to begin regenerating. While undergreen willow regenerates as the top canopy stratum, fireweed (*Chamerion angustifolium*) and field horsetail (*Equisetum arvense*) establish in the understory. Within a few years, Sitka alder (*Alnus viridis* ssp. *sinuata*) recruits have begun to establish. Given enough time without significant disturbance, Alaska cedar (*Callitropsis nootkatensis*) and subalpine fir (*Abies lasiocarpa*) may begin to establish on-site. The timing of tree establishment on-site may vary depending on seed source availability. At this point, additional forb species regenerate in the understory. If the time without disturbance is very long, subalpine fir will gradually become more dominant on site due to its higher tolerance of shade, but frequent disturbance from high-intensity floods generally precludes subalpine fir dominance on site.

Windthrow events are a potential source of disturbance on this site. Blowdown events may impact individual trees or entire stands, depending on severity. Blown-down logs trap sediment, accelerating the accumulation of fine particulate matter. High soil moisture and poor fuel conditions generally inhibit fires on this site. If this site does burn, it certainly experiences a very high fire-return interval, greater than 200 years (FEIS, 2012). Wildfire occurrence may be limited to prolonged periods of drought. If they occur, fires are typically mixed to high intensity, often stand-replacing. Stand-replacing wildfires open the canopy and favor the establishment of pioneering, shade-intolerant species. Wildfire events tend not to produce enduring effects on this site due to the much higher frequency of flooding events.

State and transition model

Additional community tables

Table 5. Community 1.1 plant community composition

Group	Common Name	Symbol	Scientific Name	Annual Production ()	Foliar Cover (%)
-------	-------------	--------	-----------------	----------------------	------------------

Table 6. Community 1.2 plant community composition

Group	Common Name	Symbol	Scientific Name	Annual Production ()	Foliar Cover (%)
-------	-------------	--------	-----------------	----------------------	------------------

Table 7. Community 1.3 plant community composition

Group	Common Name	Symbol	Scientific Name	Annual Production ()	Foliar Cover (%)
-------	-------------	--------	-----------------	----------------------	------------------

Other references

- Breemen, Nico van. "How Sphagnum Bogs down Other Plants." *Trends in Ecology & Evolution* 10, no. 7 (July 1, 1995): 270–75. 90007-1.
- Dwire, K. and Kauffman, J. 2003. Fire and Riparian Ecosystems in Landscapes in the Western United States. *Forest Ecology and Management*, Vol. 178 pg. 61-74.
- Franklin, Jerry F.; Cromack, Kermit, Jr.; Denison, William; [and others]. 1981. Ecological characteristics of old-growth Douglas-fir forests. Gen. Tech. Rep. PNW-118. Portland, OR: U.S. Department of Agriculture, Forest Service, Pacific Northwest Forest and Range Experiment Station. 48 p. [7551]
- Gavin, D., Brubaker, L. (2015). The Modern Landscape of the Olympic Peninsula. In: Late Pleistocene and Holocene Environmental Change on the Olympic Peninsula, Washington. *Ecological Studies*, vol 222. Springer, Cham. https://doi.org/10.1007/978-3-319-11014-1_1
- Geertsema, Marten & Pojar, James. (2007). Influence of landslides on biophysical diversity — A perspective from British Columbia. *Geomorphology*. 89. 55-69. 10.1016/j.geomorph.2006.07.019.
- Goheen, E.M. and Willhite, E.A. 2006. Field Guide to Common Diseases and Inspect Pests of Oregon and Washington Conifers. Portland, Oregon: USDA Forest Service, Pacific Northwest Region R6-NR-FID-PR-01-06.
- Hanley, D.P and D.M. Baumgartner. 2002. Forest Ecology in Washington. Washington State University Extension Publishing. Technical Report EB 1943.
- Hanson, E.J., D.L. Azuma and B.A. Hiserote. 2002. Site Index Equations and Mean Annual Increment Equations for Pacific Northwest Research Station Forest Inventory and Analysis Inventories, 1985-2001. USDA Forest Service Pacific Northwest Research Station, Research Note PNW-RN-533.
- Hemstrom, M., Franklin, J. 1982. Fire and Other Disturbances of the Forests in Mount Rainier National Park. *Quaternary Research*, Vol 18 pp 32-61.
- James K. Agee and Mark H. Huff. 1987. Fuel succession in a western hemlock/Douglas-fir forest. *Canadian Journal of Forest Research*. 17(7): 697-704. <https://doi.org/10.1139/x87-112>
- Nielsen, E. M., R. L. Brunner, C. Copass and L. K. Wise, 2021. Olympic National Park map class descriptions. National Park Service, Fort Collins.
- Pojar J., and MacKinnon. 1994. Plants of the Pacific Northwest Coast. Lone Pine, Vancouver, British Columbia. 528 pages.
- PRISM Climate Group, Oregon State University, <http://prism.oregonstate.edu>, visited October 2023.
- Reed Wendel, Darlene Zabowski "Fire History within the Lower Elwha River Watershed, Olympic National Park, Washington," *Northwest Science*, 84(1), 88-97, (1 January 2010)
- Rocheftort, R.M. and Peterson, D.L. 1996. Temporal and Spatial Distribution of Trees in Subalpine Meadows of Mount Rainier National Park. *Arctic and Alpine Research*, Vol. 28, No. 1 pp 52-59.
- Seastedt, T.R., Adams, G.A. 2001. Effects of Mobile Tree Islands on Alpine Tundra Soils. *Ecology*, Vol 82 pp 8-17.
- Smith, K., G. Kuhn, and L. Townsend. 2008. Culmination of Mean Annual Increment for Indicator Tree Species in the State of Washington. USDA-NRCS Technical Note Forestry-9.
- United States Department of Agriculture, Forest Service, 2015. *Silvics Manual Vol 1*. http://na.fs.fed.us/spfo/pubs/silvics_manual/Volume_1/vol1_Table_of_contents.htm, visited December 2015.
- United States Department of Agriculture, Natural Resources Conservation Service, and United States Department of the Interior, National Park Service. 2014. Ecological Site Descriptions for North Cascades National Park Complex, Washington.
- United States Department of Agriculture, Natural Resources Conservation Service, and United States Department of the Interior, National Park Service. Ecological Site Descriptions for Mount Rainier National Park, Washington.
- Van Pelt, R. 2007. Identifying Mature and Old Forests in Western Washington. Washington State Department of Natural Resources, Olympia, WA. 104
- Washington Department of Natural Resources, Natural Heritage Program. 2015. Ecological Systems of Washington State. A Guide to Identification.

Wood, David, and Moral, Roger del. "Mechanisms of Early Primary Succession in Subalpine Habitats on Mount St. Helens." *Ecology* 68, no. 4 (1987).

Zhao, Yunpeng, Chengzhu Liu, Simin Wang, Yiyun Wang, Xiaoqing Liu, Wanqing Luo, and Xiaojuan Feng. "'Triple Locks' on Soil Organic Carbon Exerted by Sphagnum Acid in Wetlands." *Geochimica et Cosmochimica Acta* 315 (December 2021): 24–37. <https://doi.org/10.1016/j.gca.2021.09.028>.

Contributors

Alec Haulotte
Erin Kreutz
Abigail Field
Max Ross
Erik Dahlke

Approval

Grant Petersen, 5/15/2025

Rangeland health reference sheet

Interpreting Indicators of Rangeland Health is a qualitative assessment protocol used to determine ecosystem condition based on benchmark characteristics described in the Reference Sheet. A suite of 17 (or more) indicators are typically considered in an assessment. The ecological site(s) representative of an assessment location must be known prior to applying the protocol and must be verified based on soils and climate. Current plant community cannot be used to identify the ecological site.

Author(s)/participant(s)	
Contact for lead author	
Date	09/26/2026
Approved by	
Approval date	
Composition (Indicators 10 and 12) based on	Annual Production

Indicators

1. Number and extent of rills:

2. Presence of water flow patterns:

3. Number and height of erosional pedestals or terracettes:

4. Bare ground from Ecological Site Description or other studies (rock, litter, lichen, moss, plant canopy are not bare ground):

5. Number of gullies and erosion associated with gullies:

6. Extent of wind scoured, blowouts and/or depositional areas:

7. Amount of litter movement (describe size and distance expected to travel):

8. Soil surface (top few mm) resistance to erosion (stability values are averages - most sites will show a range of values):

9. Soil surface structure and SOM content (include type of structure and A-horizon color and thickness):

10. Effect of community phase composition (relative proportion of different functional groups) and spatial distribution on infiltration and runoff:

11. Presence and thickness of compaction layer (usually none; describe soil profile features which may be mistaken for compaction on this site):

12. Functional/Structural Groups (list in order of descending dominance by above-ground annual-production or live foliar cover using symbols: >>, >, = to indicate much greater than, greater than, and equal to):

Dominant:

Sub-dominant:

Other:

Additional:

13. Amount of plant mortality and decadence (include which functional groups are expected to show mortality or decadence):

14. Average percent litter cover (%) and depth (in):

15. Expected annual annual-production (this is TOTAL above-ground annual-production, not just forage annual-production):

16. Potential invasive (including noxious) species (native and non-native). List species which BOTH characterize degraded states and have the potential to become a dominant or co-dominant species on the ecological site if their future establishment and growth is not actively controlled by management interventions. Species that become dominant for only one to several years (e.g., short-term response to drought or wildfire) are not invasive plants. Note that unlike other indicators, we are describing what is NOT expected in the reference state for the ecological site:
-

17. Perennial plant reproductive capability:
-