

Ecological site AX001X02X402

Frigid Udic Forest

Last updated: 1/23/2025
Accessed: 09/17/2026

General information

Provisional. A provisional ecological site description has undergone quality control and quality assurance review. It contains a working state and transition model and enough information to identify the ecological site.

MLRA notes

Major Land Resource Area (MLRA): 001X–Northern Pacific Coast Range, Foothills, and Valleys

This long and narrow resource area stretches along the Pacific Border Province of the Pacific Mountain System in Oregon and Washington. The area is bounded by the Olympic Mountains on the north and the Klamath Mountains on the south. Most of the area consists of hills and low mountains with gentle to steep slopes. The parent materials are composed primarily of young Tertiary sedimentary rocks with some minor volcanic rocks. Glacial till and outwash deposits are found in the northern half of the area within Washington. In the far southern portion of the area, near the Klamath Mountains, the sedimentary rocks are older and some have been metamorphosed. The average annual precipitation ranges from 60 to 200 inches, increasing with elevation. The dominant soil orders in this MLRA are Andisols, Inceptisols, and Ultisols. Soil depth ranges from shallow to very deep. While most soils in the area are well drained and occur on foothills, mountain slopes and ridges, floodplain and depressional soils can range from well drained to very poorly drained. Soil textures are typically medial, loamy, or clayey. The dominant soils in the area have a mesic or frigid soil temperature regime and a udic soil moisture regime; however, soils with an aquic soil moisture regime or cryic soil temperature regime do occur.

LRU notes

The North Pacific Coast Range land resource unit (LRU 2) of MLRA 1 is located in the northwestern corner on the Olympic Peninsula and within the Olympic National Forest in Washington State. LRU 2 is bounded on the west by MLRA 4a Sitka Spruce Belt and MLRA 2 Willamette and Puget Sound Valleys to the east. LRU 2 encircles the Olympic National Park (LRU 1). Several major rivers have headwaters in this LRU or carved valleys through the landscape depositing more recent alluvium. These include the Duckabush, Elwha, Queets, Quinault, Skokomish, Sol Duc, and Wynoochee Rivers.

Ecological site concept

This ecological site is found on the western Coast Range on the Olympic Peninsula in Washington state. A temperate climate supports a variety of flora and fauna. Elevations are typically between 1,250 and 3,000 feet. The most common overstory species are Pacific silver fir (*Abies amabilis*), western hemlock (*Tsuga heterophylla*) and Douglas-fir (*Pseudotsuga menziesii*). Western redcedar (*Thuja plicata*) and Pacific yew (*Taxus brevifolia*) may be present as a minor component. The understory may be sparse depending on canopy cover. Common understory species may include Alaska huckleberry (*Vaccinium alaskaense*), red huckleberry (*Vaccinium parvifolium*), twinflower (*Linnaea borealis*), queencup beadlily (*Clintonia uniflora*), and western cordilleran bunchberry (*Cornus unalaschensis*). The most common disturbance on this site is from windthrow of overstory trees, which results in patchy, small pockets of open areas. These areas commonly occur in conjunction with either root-, butt- or stem-rot. Western hemlock and Pacific silver fir are highly susceptible to rot diseases from fungi such as; *Armillaria ostoyae*, *Heterobasidion annosum*, *Phellinus weirii*, and *Echinodontium tinctorium* which may exacerbate the extent and area of disturbance. Following root disease, Pacific silver fir is highly susceptible to fir engravers. The resulting openings in the canopy allow sunlight to reach the forest floor, benefiting the understory. Disturbance by fire is infrequent as a result of the high humidity, high elevation, and precipitation within the Pacific silver fir and western hemlock zones. The site has a fire interval of 500 years and may experience stand replacing catastrophic wildfires (US Department of Agriculture, 2012).

Associated sites

AX001X02X403	Frigid Udic Moist Forest Frigid Udic Moist Forest is similar but is located on depressions or lower landscape positions that collect and retain soil moisture longer than Frigid Udic Forest. The Frigid Udic Moist Forest sites often have a large presence of three-leaf foamflower (<i>Tiarella trifoliata</i>) and lady fern (<i>Athyrium filix-femina</i>).
AX001X02X410	Mesic Udic Forest Mesic Udic Forest is located in lowland forests in elevations commonly found below Frigid Udic Forest. Mesic Udic Forest typically lacks Pacific silver fir (<i>Abies amabilis</i>) in the overstory.

Table 1. Dominant plant species

Tree	(1) <i>Abies amabilis</i> (2) <i>Tsuga heterophylla</i>
Shrub	(1) <i>Vaccinium alaskaense</i> (2) <i>Vaccinium parvifolium</i>
Herbaceous	(1) <i>Clintonia uniflora</i>

Legacy ID

F001XB402WA

Physiographic features

This ecological site occurs across mountain slopes at mid elevations. This ecological site may occur on all slopes, but is most commonly found between 20 to 75 percent slopes.

Table 2. Representative physiographic features

Landforms	(1) Mountains > Mountain slope (2) Mountains > Moraine
Flooding frequency	None
Ponding frequency	None
Elevation	380 – 910 m
Slope	20 – 100 %
Aspect	W, NW, N, NE, E, SE, S, SW

Climatic features

The climate has cool, moist summers and cold, wet winters. Mean annual precipitation ranges from 70 to 220 inches. Precipitation falls as snow during the winter months and there is a moderate snowpack. Average annual temperatures range from 41 to 46 degrees F. As a result of the shorter growing season resulting from higher elevations, colder temperatures, and longer durational snowpack, these sites tend to be moderately productive.

Table 3 Representative climatic features

Frost-free period (characteristic range)	130-180 days
Freeze-free period (characteristic range)	
Precipitation total (characteristic range)	1,780-5,590 mm

Influencing water features

In general, this ecological site is not influenced by wetland or riparian water features but may be found on stream terraces or adjacent to wetland and riparian areas. This site does not experience flooding or ponding.

Soil features

Soils that support this ecological site occur in the frigid soil temperature regime and the udic soil moisture regime.

Table 4. Representative soil features

Parent material	(1) Colluvium – basalt (2) Till (3) Glaciomarine deposits – basalt (4) Colluvium – sedimentary rock
Surface texture	(1) Very gravelly, extremely gravelly sandy loam (2) Very gravelly, extremely gravelly loam
Drainage class	Moderately well drained to well drained
Soil depth	30 – 150 cm
Surface fragment cover <=3"	30 – 50 %
Surface fragment cover >3"	0 – 30 %
Clay content (2.5-17.8cm)	10 %

Subsurface fragment volume <=3" (2.5-152.4cm)	30 – 50 %
Subsurface fragment volume >3" (2.5-152.4cm)	10 – 40 %

Ecological dynamics

State and transition model

Additional community tables

Table 5. Community 1.1 plant community composition

Group	Common Name	Symbol	Scientific Name	Annual Production ()	Foliar Cover (%)
-------	-------------	--------	-----------------	----------------------	------------------

Table 6. Community 1.2 plant community composition

Group	Common Name	Symbol	Scientific Name	Annual Production ()	Foliar Cover (%)
-------	-------------	--------	-----------------	----------------------	------------------

Table 7. Community 1.3 plant community composition

Group	Common Name	Symbol	Scientific Name	Annual Production ()	Foliar Cover (%)
-------	-------------	--------	-----------------	----------------------	------------------

Table 8. Community 1.4 plant community composition

Group	Common Name	Symbol	Scientific Name	Annual Production ()	Foliar Cover (%)
-------	-------------	--------	-----------------	----------------------	------------------

Inventory data references

Other Established Classifications for Ecological Site National Vegetation Classification: G241 North Central Pacific Maritime Silver fir-Western Hemlock Rainforest USDA Forest Service Plant Association of the Olympic National Forest: Pacific silver fir/Alaska huckleberry, Pacific silver fir/Alaska huckleberry/Queen's cup, Pacific silver fir/Alaska huckleberry/beargrass, and western hemlock/Alaska huckleberry associations Washington Department of Natural Resources Ecological Systems of Washington State- North Pacific Mesic Western Hemlock-Silver fir Forest

Other references

Cope, Amy B. 1992. *Abies amabilis*. In: Fire Effects Information System, [Online]. U.S. Department of Agriculture, Forest Service, Rocky Mountain Research Station, Fire Sciences Laboratory.

Crawford, P.D. 1990. *Abies amabilis*. Silvics of North America. [Online]. U.S. Department of Agriculture, Forest Service, Northeastern Area.

Crawford, R. C., C. B. Chappell, C. C. Thompson, and F. J. Rocchio. 2009. Vegetation Classification of Mount Rainier, North Cascades, and Olympic National Parks. Natural Resource Technical Report NPS/NCCN/NRTR-2009/211. National Park Service, Fort Collins, Colorado.

Goheen, E.M. and Willhite, E.A. 2006. Field Guide to Common Diseases and Inspect Pests of Oregon and Washington Conifers. Portland, Oregon: USDA Forest Service, Pacific Northwest Region R6-NR-FID-PR-01-06.

Henderson, J., Peter, D., Leshner, R., Shaw, D. 1989. Forested Plant Associations of the Olympic National Forest. United States Department of Agriculture Forest Service, Pacific Northwest Region. Technical Paper R6-ECOL-TP 001-88.

McCain, C., Diaz, N. 2002. Field Guide to the Forested Plant Associations of the Northern Oregon Coast Range. United States Department of Agriculture Forest Service, Pacific Northwest Region. Technical Paper R6-NR-Ecol-TP-03-02

Pojar J., and MacKinnon. 1994. Plants of the Pacific Northwest Coast. Lone Pine, Vancouver, British Columbia. 528 pages.

PRISM Climate Group, Oregon State University, <http://prism.oregonstate.edu>, visited Feb., 2015.

Soil Survey Staff. 2014. Keys to Soil Taxonomy, 12th ed. USDA-Natural Resources Conservation Service, Washington, DC.

Soil Survey Staff. 1999. Soil Taxonomy: A Basic System of Soil classification for Making and Interpreting Soil Surveys. 2nd ed. USDA-Natural Resources Conservation Service. U.S. Department of Agriculture Handbook 436.

U.S. Department of Agriculture, Forest Service, Missoula Fire Sciences Laboratory. 2012. Information from LANDFIRE on fire regimes of

wet-mesic western hemlock communities. In: Fire Effects Information System, [Online]. U.S. Department of Agriculture, Forest Service, Rocky Mountain Research Station, Missoula Fire Sciences Laboratory (Producer). Available: www.fs.fed.us/database/feis/fire_regimes/Western_hemlock_wet/all.html [2019, October 17].

United States National Vegetation Classification. 2016. United States National Vegetation Classification Database, V2.0. Federal Geographic Data Committee, Vegetation Subcommittee, Washington DC. (accessed 28, November, 2016.)

Washington Department of Natural Resources, Natural Heritage Program. 2015. Ecological Systems of Washington State. A Guide to Identification.

Contributors

Erin Kreutz
Erik Dahlke

Approval

Kirt Walstad, 1/23/2025

Rangeland health reference sheet

Interpreting Indicators of Rangeland Health is a qualitative assessment protocol used to determine ecosystem condition based on benchmark characteristics described in the Reference Sheet. A suite of 17 (or more) indicators are typically considered in an assessment. The ecological site(s) representative of an assessment location must be known prior to applying the protocol and must be verified based on soils and climate. Current plant community cannot be used to identify the ecological site.

Author(s)/participant(s)	
Contact for lead author	
Date	12/03/2021
Approved by	
Approval date	
Composition (Indicators 10 and 12) based on	Annual Production

Indicators

1. Number and extent of rills:

2. Presence of water flow patterns:

3. Number and height of erosional pedestals or terracettes:

4. Bare ground from Ecological Site Description or other studies (rock, litter, lichen, moss, plant canopy are not bare ground):

5. Number of gullies and erosion associated with gullies:

6. Extent of wind scoured, blowouts and/or depositional areas:

7. Amount of litter movement (describe size and distance expected to travel):

8. Soil surface (top few mm) resistance to erosion (stability values are averages - most sites will show a range of values):

9. Soil surface structure and SOM content (include type of structure and A-horizon color and thickness):

10. Effect of community phase composition (relative proportion of different functional groups) and spatial distribution on infiltration and runoff:

11. Presence and thickness of compaction layer (usually none; describe soil profile features which may be mistaken for compaction on this site):

12. Functional/Structural Groups (list in order of descending dominance by above-ground annual-production or live foliar cover using symbols: >>, >, = to indicate much greater than, greater than, and equal to):

Dominant:

Sub-dominant:

Other:

Additional:

13. Amount of plant mortality and decadence (include which functional groups are expected to show mortality or decadence):

14. Average percent litter cover (%) and depth (in):

15. Expected annual annual-production (this is TOTAL above-ground annual-production, not just forage annual-production):

16. Potential invasive (including noxious) species (native and non-native). List species which BOTH characterize degraded states and have the potential to become a dominant or co-dominant species on the ecological site if their future establishment and growth is not actively controlled by management interventions. Species that become dominant for only one to several years (e.g., short-term response to drought or wildfire) are not invasive plants. Note that unlike other indicators, we are describing what is NOT expected in the reference state for the ecological site:
-

17. Perennial plant reproductive capability:
-