

Ecological site F002XB004OR

Fragipan Hill Group

Last updated: 12/03/2024
Accessed: 08/29/2026

General information

Provisional. A provisional ecological site description has undergone quality control and quality assurance review. It contains a working state and transition model and enough information to identify the ecological site.

MLRA notes

Major Land Resource Area (MLRA): 002X–Willamette and Puget Sound Valleys

The Willamette and Puget Sound Valleys Major Land Resource Area (MLRA 2) is located in western Washington and Oregon. It occupies a forearc basin between coast ranges and the Cascade Mountain volcanic arc. The northern part contains Pleistocene drift, outwash, lacustrine and glaciomarine deposits associated with continental glaciers. The southern part contains Late Pleistocene deposits from glacial outburst floods (Missoula Floods). Climate is mild and moist, with a long growing season. Mean annual precipitation ranges from 20 to 60 inches, falling mostly in fall, winter, and spring. Summers are dry. Soil temperature regime is mesic and soil moisture regimes are xeric and aquic. Most sites in this MLRA can support forested vegetation, but some were maintained as prairie, savanna, or woodland through cultural burning prior to Euro-American settlement. Puget Sound has a moderating effect on temperatures and humidity can be higher in the northern part of the MLRA. Douglas-fir (*Pseudotsuga menziesii*) is widespread throughout. Oregon white oak (*Quercus garryana*) is common on uplands in the south and on warm, exposed or droughty sites in the north. Pacific madrone (*Arbutus menziesii*) occurs in areas close to salt water. Western hemlock (*Tsuga heterophylla*) is codominant with Douglas-fir in the north. Floodplains usually contain black cottonwood (*Populus balsamifera* ssp. *trichocarpa*) and red alder (*Alnus rubra*). Oregon ash (*Fraxinus latifolia*) is typical of forested wetlands in the south. Forestry, urban development, and cultivated agriculture are currently the most extensive land uses (Soil Survey Staff, 2006).

LRU notes

The Portland Basin and Hills Land Resource Unit (LRU B) is located in northwest Oregon and southwest Washington. It includes the Portland Basin and surrounding hills. Isolated areas of LRU C (Willamette Valley) occur below 400 feet in the Tualitan Valley on loamy or silty Missoula Flood deposits. The Columbia River Gorge borders this LRU on the east. Brackish tidewater beginning near the town of Cathlamet marks the northwestern limit of this LRU along the Columbia River floodplain. Elevation ranges from sea level to about 1200 feet. Topography is flat to steep. Major landforms include the Columbia River floodplain, glaciofluvial terraces, hills, and foothills. The valley floor is underlain by Pleistocene fluvial deposits (Rowland Formation). Hills and foothills are underlain by Eocene to Pliocene sedimentary rocks (Yamhill, Nestucca, Scotts Mills, Molalla, and Troutdale Formations), Miocene Columbia River Basalt, or Plio-Pleistocene Boring Lavas (Orr et al., 1992). Gravelly or sandy Late Pleistocene Missoula Flood deposits can occur below 400 feet elevation. Hills are covered in loess, and fragipans (brittle subsoil layers) are common. Mean annual precipitation ranges from 35 to 60 inches. Most falls as rain between October and May. The frost-free period ranges from 160 to 210 days. Ice storms occur each winter. Locations near the Columbia River Gorge experience strong winds. Most locations experience less summer moisture stress compared with the main Willamette Valley; summertime average daily maximum temperatures at Vancouver, WA are 1 to 3 degrees F cooler compared with Corvallis, OR (Agricultural Climate Information System, 2007a, 2007b). Cultural fire use prior to Euro-American settlement was apparently less than in the main Willamette Valley, though it was used in some areas. General Land Office (GLO) land surveys conducted between 1851 and 1910 indicate that forest and woodland communities were more prevalent than prairies and savannas (Hulse et al., 2002). Forested reference community phases have been chosen for these upland ecological sites. Presence of Oregon white oak (*Quercus garryana*), and absence of western hemlock (*Tsuga heterophylla*) distinguish this area from coast range (MLRA 1) and Cascade mountain (MLRA 3) ecological types in Oregon. Relative abundance of western redcedar (*Thuja plicata*) helps distinguish this area from the Willamette Valley (LRU C).

Classification relationships

This ecological site group occurs in mosaic with well drained forested sites similar to the following LANDFIRE Biophysical Setting (BpS): • LANDFIRE Biophysical Setting: North Pacific Dry Douglas-fir Forest and Woodland (0710350)

Ecological site concept

This site occurs on loess-covered hills. A fragipan restricts roots and impedes the downward movement of water. Consequently, a perched water table is located within 20 inches of the soil surface during early spring. The rooting zone is dry 45 to 60 consecutive days during the summer. The seasonal water table restricts rooting depth for most conifers. The reference plant community is bigleaf maple - red alder / beaked hazelnut / forbs.

Table 1. Dominant plant species

Tree	(1) <i>Acer macrophyllum</i> (2) <i>Alnus rubra</i>
Shrub	(1) <i>Corylus cornuta</i>
Herbaceous	Not specified

Physiographic features

Landform: hills

Parent material: loess

Elevation: 200 to 1200 feet

Slope: 0 to 30 percent

Flooding: none

Ponding: none

This site occurs on the Dolph geomorphic surface located around the edge of the Portland Basin (Balster and Parsons, 1968; Reckendorf, 1993).

Table 2. Representative physiographic features

Landforms	(1) Hill
Flooding frequency	None
Ponding frequency	None
Elevation	60 – 370 m
Slope	0 – 30 %

Climatic features

Mean annual air temperature: 50 to 54 degrees F

Mean annual precipitation: 45 to 60 inches

Frost free period: 165 to 210 days

Influencing water features

Perched water tables rise above 20 inches of the soil surface, and the rooting zone is saturated during the early growing season.

Wetland description

None

Soil features

Drainage class: somewhat poorly or poorly drained

Parent material: loess

Soil restrictive feature(s): a fragipan (brittle subsoil layer) occurs in most soils

Soil moisture regime: aquic or xeric

Soil temperature regime: mesic

Particle-size family(s): fine-silty

Soil mineralogy: mixed

Cation exchange capacity: superactive

Soil reaction: moderately or strongly acid

Perched water tables rise above 20 inches of the soil surface, and the rooting zone is saturated during the early growing season. Fragipans are often present. Soils classify as Inceptisols. Additions of loess have rejuvenated and increased the fertility of these soils; cation-exchange activity class is higher compared with soils on the Dolph geomorphic surface in the Willamette Valley (LRU C).

Soils correlated with this site include Delena, Powell, Borges, and Wollent.

Ecological dynamics

Central Concept

This site occurs on loess-covered hills. Soils are silty. A fragipan restricts roots and impedes the downward movement of water. Consequently, a perched water table is located within 20 inches of the soil surface during early spring. The rooting zone is dry 45 to 60 consecutive days during the summer which is typical for upland sites in this LRU. The seasonal water table restricts rooting depth for most conifers. Deciduous forest tends to develop. The reference plant community is bigleaf maple - red alder / beaked hazelnut / forbs.

Disturbance

This site developed under a mixed-severity to low-severity fire regime; the estimated mean fire return interval is 36 years (Kertis et al., 2007). Cultural burning was not focused here prior to Euro-American settlement. General Land Office (GLO) surveys noted mainly conifer forest. Woodlands were rare. Fire has been suppressed since modern wildland fire suppression efforts began.

Tree-throw occurs in forested communities especially where a seasonal water table exists in the soil profile. Pocket gophers (*Thomomys* spp.) make burrows and mounds in early-seral communities (Oregon Department of Fish and Wildlife).

Plant Composition

Conifer forest was typical prior to Euro-American settlement but modern fire suppression efforts have generally led to denser forest stands through the proliferation of shade-tolerant, fire-intolerant species. However, prairie or savanna can be created and maintained on this site.

Representative native plants are listed below. Not all species are present within the same community phase. Plant lists (especially for grasses, grasslikes, and forbs) are incomplete.

TREES:

Douglas-fir (*Pseudotsuga menziesii*)

western redcedar (*Thuja plicata*)

grand fir (*Abies grandis*)

Oregon ash (*Fraxinus latifolia*)

red alder (*Alnus rubra*)

bigleaf maple (*Acer macrophyllum*)

ponderosa pine (*Pinus ponderosa*)

Oregon white oak (*Quercus garryana*)

Pacific dogwood (*Cornus nuttallii*)

cascara buckthorn (*Frangula purshiana*)

SHRUBS:

vine maple (*Acer circinatum*)

Cascade barberry (*Mahonia nervosa*)

salal (*Gaultheria shallon*)

common snowberry (*Symphoricarpos albus*)

creeping snowberry (*Symphoricarpos mollis*)

Saskatoon serviceberry (*Amelanchier alnifolia*)

beaked hazelnut (*Corylus cornuta*)
 Indian plum (*Oemleria cerasiformis*)
 rose (*Rosa* spp.)
 California blackberry (*Rubus ursinus*)
 black hawthorn (*Crataegus douglasii*)
 Oregon crab apple (*Malus fusca*)
 Pacific poison oak (*Toxicodendron diversilobum*)
 oceanspray (*Holodiscus discolor*)
 hollyleaved barberry (*Mahonia aquifolium*)

GRASSLIKES:

sedge (*Carex* spp.)
 rush (*Juncus* spp.)

State and transition model

Additional community tables

Table 3. Community 1.1 plant community composition

Group	Common Name	Symbol	Scientific Name	Annual Production ()	Foliar Cover (%)
-------	-------------	--------	-----------------	----------------------	------------------

Table 4. Community 1.2 plant community composition

Group	Common Name	Symbol	Scientific Name	Annual Production ()	Foliar Cover (%)
-------	-------------	--------	-----------------	----------------------	------------------

Table 5. Community 1.3 plant community composition

Group	Common Name	Symbol	Scientific Name	Annual Production ()	Foliar Cover (%)
-------	-------------	--------	-----------------	----------------------	------------------

Table 6. Community 2.1 plant community composition

Group	Common Name	Symbol	Scientific Name	Annual Production ()	Foliar Cover (%)
-------	-------------	--------	-----------------	----------------------	------------------

Table 7. Community 2.2 plant community composition

Group	Common Name	Symbol	Scientific Name	Annual Production ()	Foliar Cover (%)
-------	-------------	--------	-----------------	----------------------	------------------

Table 8. Community 2.3 plant community composition

Group	Common Name	Symbol	Scientific Name	Annual Production ()	Foliar Cover (%)
-------	-------------	--------	-----------------	----------------------	------------------

Other references

Agricultural Climate Information System. (2007). WETS Station Data for Corvallis State University, OR, 1971-2000. [Online]. Available at <http://agacis.rcc-acis.org/?fips=41003> (accessed on 5/7/2020).

Agricultural Climate Information System. (2007). WETS Station Data for Vancouver 4 NNE, WA, 1971-2000. [Online]. Available at <http://agacis.rcc-acis.org/?fips=53011> (accessed on 5/7/2020).

Balster, C.A., and Parsons, R.B. (1968). Geomorphology and soils Willamette Valley, Oregon. Oregon State University Experiment Station Special Report 265. <https://ir.library.oregonstate.edu/downloads/mg74qm961>

Christy, J., and Alverson, E. (2011). Historical vegetation of the Willamette Valley, Oregon, circa 1850. Northwest Science. 85(2):93-107. <https://doi.org/10.3955/046.085.0202>

Christy, J.A., Alverson, E.R., Dougherty, M.P., Kolar, S.C., Alton, C.W., Hawes, S.M., Ashkenas, L., and Minear, P. (2011). GLO historical vegetation of the Willamette Valley, Oregon, 1851-1910. ArcMap shapefile, Version 2011_04. Oregon Biodiversity Information Center, Portland State University. Available at http://www.pdx.edu/sites/www.pdx.edu.pnwlamp/files/glo_willamette_2011_04.zip (accessed on 11/14/2019).

Franklin, J., and Dyrness, C. (1973). Interior valleys of western Oregon. p. 110-129. In Natural Vegetation of Oregon and Washington. United States Department of Agriculture Forest Service, Pacific Northwest Forest and Range Experiment Station. General Technical Report PNW-8.

Hulse, D., Gregory, S., and Baker, J. (2002). Presettlement Vegetation circa 1850. p. 38-39. In Pacific Northwest Ecosystem Research

Consortium (ed.) Willamette River Basin Planning Atlas: Trajectories of Environmental and Ecological Change. [Online]. Available at http://www.fsl.orst.edu/pnwer/wrb/Atlas_web_compressed/4.Biotic_Systems/4b.presetveg_web.pdf (accessed on 9/28/2015).

Johannessen, C. L., Davenport, W.A., Millet, A., and McWilliams, S. (1971). The vegetation of the Willamette Valley. *Annals of the Association of American Geographers*. 61(2):286-302.

Kertis, J., DeMeo, T., and Kopper, K. (2007). North Pacific Dry Douglas-fir Forest and Woodland. p. 90-93. In *LANDFIRE Biophysical Setting Model Descriptions*. [Online]. Available at https://www.landfire.gov/national_veg_models_op2.php (accessed on 6/3/2020).

Oregon Department of Fish and Wildlife. Pocket Gophers. [Online]. Available at <https://myodfw.com/wildlife-viewing/species/pocket-gophers> (accessed on 5/21/2020).

Orr, E., Orr, W., and Baldwin, E. (1992). Willamette Valley. p. 203-221. In *Geology of Oregon*. 4th ed. Kendall/Hunt Publishing Company.

Reckendorf, F. (1993). Geomorphology, stratigraphy, and soil interpretations, Willamette Valley, Oregon. p. 178-199. In J.M. Kimble (ed.) *Proceedings of the Eighth International Soil Management Workshop: Utilization of Soil Survey Information for Sustainable Land Use*. Oregon, California, and Nevada. 11-24 July 1992; May 1993. United States Department of Agriculture Soil Conservation Service National Soil Survey Center.

Schoeneberger, P.J., Wysocki, D.A., Benham, E.C., and Soil Survey Staff. (2012). *Field book for describing and sampling soils*, Version 3.0. Natural Resources Conservation Service, National Soil Survey Center, Lincoln, NE.

Soil Survey Staff, Natural Resources Conservation Service, United States Department of Agriculture. (2006). *Land Resource Regions and Major Land Resource Areas of the United States, the Caribbean, and the Pacific Basin*. Agricultural Handbook 296. https://www.nrcs.usda.gov/Internet/FSE_DOCUMENTS/nrcs142p2_050898.pdf

Soil Survey Staff. (2014). *Keys to Soil Taxonomy*, 12th ed. USDA-Natural Resources Conservation Service, Washington, DC.

Soil Survey Staff, Natural Resources Conservation Service, United States Department of Agriculture. *Official Soil Series Descriptions*. Online. Available at https://www.nrcs.usda.gov/wps/portal/nrcs/detail/soils/survey/geo/?cid=nrcs142p2_053587 (accessed 2019 to 2020).

Soil Survey Staff, Natural Resources Conservation Service, United States Department of Agriculture. *Soil Survey Geographic (SSURGO) Database for Oregon (multiple counties)*. [Online]. Available at <https://websoilsurvey.sc.egov.usda.gov/App/HomePage.htm> (accessed in 2020).

Yeats, R.S., Graven, E.P., Werner, K.S., Goldfinger, C. and Popowski, T. (1996). Tectonics of the Willamette Valley, Oregon. p. 183-222. In Rogers, Albert M., Walsh, Timothy J., Kockelman, William J., and Priest, George R. (ed.) *Assessing earthquake hazards and reducing risk in the Pacific Northwest*. US Geological Survey Professional Paper 1560.

Approval

Kirt Walstad, 12/03/2024

Rangeland health reference sheet

Interpreting Indicators of Rangeland Health is a qualitative assessment protocol used to determine ecosystem condition based on benchmark characteristics described in the Reference Sheet. A suite of 17 (or more) indicators are typically considered in an assessment. The ecological site(s) representative of an assessment location must be known prior to applying the protocol and must be verified based on soils and climate. Current plant community cannot be used to identify the ecological site.

Author(s)/participant(s)	
Contact for lead author	
Date	10/03/2023
Approved by	

Approval date	
Composition (Indicators 10 and 12) based on	Annual Production

Indicators

1. Number and extent of rills:

2. Presence of water flow patterns:

3. Number and height of erosional pedestals or terracettes:

4. Bare ground from Ecological Site Description or other studies (rock, litter, lichen, moss, plant canopy are not bare ground):

5. Number of gullies and erosion associated with gullies:

6. Extent of wind scoured, blowouts and/or depositional areas:

7. Amount of litter movement (describe size and distance expected to travel):

8. Soil surface (top few mm) resistance to erosion (stability values are averages - most sites will show a range of values):

9. Soil surface structure and SOM content (include type of structure and A-horizon color and thickness):

10. Effect of community phase composition (relative proportion of different functional groups) and spatial distribution on infiltration and runoff:

11. Presence and thickness of compaction layer (usually none; describe soil profile features which may be mistaken for compaction on this site):

12. Functional/Structural Groups (list in order of descending dominance by above-ground annual-production or live foliar cover using symbols: >>, >, = to indicate much greater than, greater than, and equal to):

Dominant:

Sub-dominant:

Other:

Additional:

13. Amount of plant mortality and decadence (include which functional groups are expected to show mortality or decadence):

14. Average percent litter cover (%) and depth (in):

15. Expected annual annual-production (this is TOTAL above-ground annual-production, not just forage annual-production):

16. Potential invasive (including noxious) species (native and non-native). List species which BOTH characterize degraded states and have the potential to become a dominant or co-dominant species on the ecological site if their future establishment and growth is not actively controlled by management interventions. Species that become dominant for only one to several years (e.g., short-term response to drought or wildfire) are not invasive plants. Note that unlike other indicators, we are describing what is NOT expected in the reference state for the ecological site:

17. Perennial plant reproductive capability:
