

Ecological site F003XN947WA

Southern Washington Cascades Low Cryic Coniferous Forest

Last updated: 1/30/2025
Accessed: 09/26/2026

General information

Provisional. A provisional ecological site description has undergone quality control and quality assurance review. It contains a working state and transition model and enough information to identify the ecological site.

MLRA notes

Major Land Resource Area (MLRA): 003X—Olympic and Cascade Mountains

Steep mountains and narrow to broad, gently sloping valleys characterize this MLRA. A triple junction of two oceanic plates and one continental plate is directly offshore from Puget Sound. Subduction of the oceanic plates under the westerly and northwesterly moving continental plate contributes to volcanic activity in the Cascade Mountains. Movement among these plates has resulted in major earthquakes and the formation of large stratovolcanoes. The Cascade Mountains consist primarily of volcanic crystalline rock and some associated metasedimentary rock. The mean annual precipitation is dominantly 60 to 100 inches, but it is 30 to 60 inches on the east side of the Cascade Mountains. The soil orders in this MLRA are dominantly Andisols, Spodosols, and Inceptisols and minor areas of Entisols and Histosols. The soils are dominantly in the frigid or cryic temperature regime and the udic moisture regime. The soils generally are shallow to very deep, well drained, ashy to medial, and loamy or sandy. They are on mountain slopes and ridges.

Ecological site concept

This ecological site is in cold, moist areas. It is at middle elevations (about 2,600 to 4,900 feet) on the west side of the Cascade Crest in Mount Rainier National Park. The site commonly is on debris aprons, bedrock benches, ridges, and glacial-valley walls. The temperate climate supports a variety of flora and fauna. The soils that support this ecological site are in the low cryic soil temperature regime and the udic soil moisture regime. They are moderately well drained or well drained and are shallow to bedrock to very deep. Laughingwater soils have a seasonal high water table at a depth of 19 to 40 inches some time during the growing season. Soil moisture is not a limiting factor to forest growth because of the abundance of precipitation and the inherent water-holding properties of soils influenced by volcanic ash. Slope and aspect are not defining features of this ecological site. Pacific silver fir (*Abies amabilis*) and western hemlock (*Tsuga heterophylla*) are the most common overstory species. Some noble fir (*Abies procera*), western redcedar (*Thuja plicata*), Douglas-fir (*Pseudotsuga menziesii*), western white pine (*Pinus monticola*), and Engelmann spruce (*Picea engelmannii*) are present. Common understory shrubs include black mountain huckleberry (*Vaccinium membranaceum*), red huckleberry (*Vaccinium parvifolium*), prince's pine (*Chimaphila umbellata*), bunchberry dogwood (*Cornus canadensis*), and vine maple (*Acer circinatum*). The most common disturbance on this site is windthrow because of the seasonal high water table. The resulting openings in the canopy allow more sunlight to reach the forest floor. Other disturbances include forest pathogens, such as root rot, and high-intensity, low-frequency (300 years or more) fires that are stand replacing.

Associated sites

F003XN943WA	Southern Washington Cascades Frigid Coniferous Forest Ecological site F003XN947WA, Low Cryic Coniferous Forest is located at higher elevations from F003XN943WA. Vegetation indicators for F003XN947WA include Pacific silver fir, noble fir, and black mountain huckleberry.
F003XN951WA	Southern Washington Cascades High Cryic Coniferous Forest Ecological site F003XN951WA, High Cryic Coniferous Forest is located at higher elevations from F003XN947WA. Vegetation indicators for F003XN951WA include mountain hemlock, subalpine fir, rusty menziesia, Sitka mountain-ash, and common beargrass.

Similar sites

F003XN945WA	Southern Washington Cascades Wet Low Cryic Coniferous Forest Ecological F003XN945WA, Southern Washington Cascades Wet Low Cryic Coniferous Forest, has features similar to those of site F003XN947WA, Washington Cascades Low Cryic Coniferous Forest. The soils of site F003XN945WA are subject to frequent ponding conditions. Species adapted to the wetter conditions, such as American skunkcabbage and devilsclub, are common.
F003XN946WA	Southern Washington Cascades Moist Low Cryic Coniferous Forest Ecological F003XN946WA, Southern Washington Cascades Moist Low Cryic Coniferous Forest, has features similar to those of site F003XN947WA, Washington Cascades Low Cryic Coniferous Forest. The soils of site F003XN947WA are well drained. Species adapted to the drier conditions, such as Pacific silver fir, western hemlock, and black mountain huckleberry, are common.

Table 1. Dominant plant species

Tree	(1) <i>Abies amabilis</i> (2) <i>Tsuga heterophylla</i>
Shrub	(1) <i>Vaccinium membranaceum</i>
Herbaceous	(1) <i>Linnaea borealis</i>

Physiographic features

This ecological site is on debris aprons of mountain slopes, bedrock benches, glacial-valley walls, and ridges at the lower montane elevations (2,600 to 4,900 feet). The site is on all aspects. Slope is dominantly 5 to 45 percent. This site is the second largest in extent in Mount Rainier National Park.

Table 2. Representative physiographic features

Landforms	(1) Mountain slope (2) Bench (3) Glacial-valley wall (4) Ridge
Flooding frequency	None
Ponding frequency	None
Elevation	790 – 1,490 m
Slope	0 – 100 %
Water table depth	50 – 200 cm

Aspect	W, NW, N, NE, E, SE, S, SW
--------	----------------------------

Climatic features

Most of the annual precipitation is received in October through April. The mean annual precipitation is 72 to 87 inches, and the mean annual air temperature is 42.0 to 47.5 degrees F. Generally, the summers are cool and dry and the winters are cold and wet.

Table 3 Representative climatic features

Frost-free period (actual range)	90-130 days
Freeze-free period (actual range)	
Precipitation total (actual range)	1,830-2,210 mm

Influencing water features

Generally, this ecological site is not influenced by wetland or riparian water features. The site is not subject to flooding or ponding. The Laughingwater soils have a seasonal high water table at a depth of 19 to 40 inches some time during the growing season. The water table typically rises in spring and recedes in fall.

Soil features

Applicable soils: Longmire, Laughingwater, Arahustan, Ohanapecosh

Applicable soil map units: 8110, 8120, 8125, 8130, 8150, 9100, 9101, 9110, 9120, 9125

The Longmire and Laughingwater soils are deep or very deep, the Arahustan soils are moderately deep, and the Ohanapecosh soils are shallow. The Laughingwater soils have a seasonal high water table at a depth of 19 to 40 inches some time during the growing season. Generally, the soils have a mantle of volcanic ash over colluvium or residuum derived from andesite. The shallow Ohanapecosh series have volcanic ash directly over andesite. The mantle of volcanic ash is characterized by low bulk density, high available water-holding capacity, gravel-sized pumice parafragments, and sandy loam or loamy sand textures. The subsoil, where present, consists of colluvium derived from andesite. It is sandy loam and has andesite fragments. Podsolization is the dominant pedogenic process in the soils. All of the soils have an albic and a spodic diagnostic horizon. A thin organic horizon that consists of decomposing twigs, needles, and litter is on the soil surface. This horizon helps to protect the soil from wind and water erosion.

Table 4. Representative soil features

Parent material	(1) Acidic volcanic ash – andesite (2) Colluvium – andesite
Surface texture	(1) Ashy sandy loam (2) Paragravelly, ashy sandy loam (3) Paragravelly, ashy loamy sand
Drainage class	Well drained
Soil depth	30 – 150 cm

Surface fragment cover <=3"	0 – 10 %
Surface fragment cover >3"	0 – 10 %
Available water capacity (Depth not specified)	6.35 – 17.78 cm
Soil reaction (1:1 water) (Depth not specified)	3.5 – 10
Subsurface fragment volume <=3" (Depth not specified)	10 – 50 %
Subsurface fragment volume >3" (Depth not specified)	0 – 20 %

Ecological dynamics

This ecological site is in cold, moist areas at the middle elevations (2,600 to 4,900 feet) on the west side of the Cascade Crest. The site commonly is on debris aprons, bedrock benches, ridges, and glacial-valley walls. The mild, temperate climate supports a variety of flora and fauna.

The most common overstory species are Pacific silver fir (*Abies amabilis*) and western hemlock (*Tsuga heterophylla*). Other common species include noble fir (*Abies procera*), western redcedar (*Thuja plicata*), Douglas-fir (*Pseudotsuga menziesii*), western white pine (*Pinus monticola*), and Engelmann spruce (*Picea engelmannii*).

Common understory shrubs include black mountain huckleberry (*Vaccinium membranaceum*), red huckleberry (*Vaccinium parvifolium*), prince's pine (*Chimaphila umbellata*), bunchberry dogwood (*Cornus canadensis*), and vine maple (*Acer circinatum*). Common forbs include twinflower (*Linnaea borealis*), western swordfern (*Polystichum munitum*), deerfoot vanillaleaf (*Achlys triphylla*), myrtle pachistima (*Paxistima myrsinites*), and common beargrass (*Xerophyllum tenax*). As the stand ages, competition for moisture and sunlight increases mortality of the vegetation. This results in a wide range of snag maturation and an abundance of coarse woody debris on the forest floor.

The most common disturbance on this site is windthrow. Pacific silver fir and western hemlock are susceptible to windthrow. The resulting openings in the canopy allow more sunlight to reach the forest floor. Other disturbances include forest pathogens, such as root rot, and high-intensity, low-frequency (300 years or more) fires that are stand replacing.

State and transition model

Additional community tables

Table 5. Community 1.1 plant community composition

Group	Common Name	Symbol	Scientific Name	Annual Production ()	Foliar Cover (%)
-------	-------------	--------	-----------------	----------------------	------------------

Table 6. Community 1.1 forest overstory composition

Common Name	Symbol	Scientific Name	Nativity	Height M	Canopy Cover (%)	Diameter Cm	Basal Area (square M/hectare)
Tree							
Pacific silver fir	ABAM	<i>Abies amabilis</i>	Native	–	–	–	–
western hemlock	TSHE	<i>Tsuga heterophylla</i>	Native	–	–	–	–
Douglas-fir	PSME	<i>Pseudotsuga menziesii</i>	Native	–	–	–	–

noble fir	ABPR	<i>Abies procera</i>	Native	–	–	–	–
western redcedar	THPL	<i>Thuja plicata</i>	Native	–	–	–	–

Table 7. Community 1.1 forest understory composition

Common Name	Symbol	Scientific Name	Nativity	Height (m)	Canopy Cover (%)
Forb/Herb					
sweet after death	ACTR	<i>Achlys triphylla</i>	Native	0.3-11	1-60
common beargrass	XETE	<i>Xerophyllum tenax</i>	Native	0.3-11	1-45
twinlineflower	LIBO3	<i>Linnaea borealis</i>	Native	0.6-3.7	1-40
roughfruit berry	RULA2	<i>Rubus lasiococcus</i>	Native	0.3-3	1-25
western rattlesnake plantain	GOOB2	<i>Goodyera oblongifolia</i>	Native	0.3-3.7	1-5
Shrub/Subshrub					
thinleaf huckleberry	VAME	<i>Vaccinium membranaceum</i>	Native	0.3-29.3	1-50
Cascade barberry	MANE2	<i>Mahonia nervosa</i>	Native	0.3-11	1-40
pipsissewa	CHUM	<i>Chimaphila umbellata</i>	Native	0.3-5.5	1-40
vine maple	ACCI	<i>Acer circinatum</i>	Native	0.3-128	1-40
red huckleberry	VAPA	<i>Vaccinium parvifolium</i>	Native	0.3-18.3	1-25

Table 8. Community 1.2 plant community composition

Group	Common Name	Symbol	Scientific Name	Annual Production ()	Foliar Cover (%)
-------	-------------	--------	-----------------	----------------------	------------------

Table 9. Community 1.3 plant community composition

Group	Common Name	Symbol	Scientific Name	Annual Production ()	Foliar Cover (%)
-------	-------------	--------	-----------------	----------------------	------------------

Table 10. Community 1.4 plant community composition

Group	Common Name	Symbol	Scientific Name	Annual Production ()	Foliar Cover (%)
-------	-------------	--------	-----------------	----------------------	------------------

Table 11. Community 1.5 plant community composition

Group	Common Name	Symbol	Scientific Name	Annual Production ()	Foliar Cover (%)
-------	-------------	--------	-----------------	----------------------	------------------

Table 12. Community 1.6 plant community composition

Group	Common Name	Symbol	Scientific Name	Annual Production ()	Foliar Cover (%)
-------	-------------	--------	-----------------	----------------------	------------------

Animal community

Forests of old-growth Pacific silver fir provide important habitat for mountain goats and northern spotted owl. The old-growth forests provide nesting and forage habitat for northern spotted owl, and the less mature forests provide key dispersal habitat. The seeds of Pacific silver fir are sought after by birds, rodents, and squirrels; however, Pacific silver fir is not palatable to elk (Cope, 1992).

Recreational uses

This ecological site provides hiking and backpacking opportunities in May through October, when the site is free of snow.

Other information

Pathogen Information Annosus, armillaria, and laminated root rots can infest Pacific silver fir and western hemlock forests. White pine blister rust impacts early seral forests of western white pine, and it may increase the rate of re-establishment of firs after a stand-replacing event. Foliar diseases, such as balsam woolly adelgid and western spruce budworm, are major pests in true fir forests. Outbreaks can cause severe dieback and mortality, depending on the level and duration of the infestation. Annosus root disease (Heterobasidion annosum) affects nearly all conifer species in the Pacific Northwest. It is slow growing, but it can cause severe root and butt decay. It commonly affects stands in conjunction with armillaria and laminated root rots and precedes bark beetle infestations. Signs and symptoms of annosus commonly are difficult to differentiate from armillaria. Obvious signs include circular pockets of windthrown trees and canopy dieback. A distinctive fruiting body, called a conk, is in hollow stumps of forests affected by annosus. Other identifying

features include a reddish stain, decay, and a spongy texture in the heartwood and roots (Goheen, 2006). Regeneration of conifers may be unsuccessful for several decades, or until the soil is void of fungal inoculum. Applying borax on freshly cut tree stumps within 24 hours may reduce the rate of spread in managed stands. Stands impacted by annosus are highly susceptible to damage from the fir engraver beetle (*Scolylus ventralis*), which may cause devastating mortality of Pacific silver fir stands. Armillaria root disease (*Armillaria ostoyae*) affects conifers and shrubs in the Pacific Northwest. It has the potential to affect several acres and cause widespread mortality of trees. Western larch, cedar, ponderosa pine, western white pine, and lodgepole pine are susceptible to armillaria, but mixed conifer stands commonly are more tolerant of the disease. Young regenerating stands (less than 30 years) are most affected, which may delay the maturation and succession of the forest. Identifiers of armillaria are similar to those of other root diseases; however, armillaria produces a distinct white mycelial fan between the wood and bark. Rhizomorphs, or brown shoestrings of fungal mycelia, are common under the bark and roots (Goheen, 2006). High resin flow and excessive sapping is also common. Management may be limited to planting of less susceptible species. Hemlock and Douglas-fir forests of the Washington and Oregon Cascade Mountains are highly susceptible to laminated root rot (*Phellinus weirii*), which results in moderate disturbances and openings in the forest. The fungus can cause severe root rot and butt decay and result in stunted growth and mortality. Western hemlock, Pacific silver fir, subalpine fir, and noble fir may be affected by laminated root rot; however, these species rarely are killed by the disease. Signs and symptoms of laminated root rot include pockets of dead and fallen trees that are broken at or near ground level. Decay is identified by a brown to reddish brown speckled staining in the sapwood and wood that separates along the growth rings. Regeneration of highly susceptible species in the area affected by the fungus typically is unsuccessful (Goheen, 2006). White pine blister rust (*Cronartium ribicola*) affects early seral white pine forests and causes mortality of the young trees. The disease commonly increases the rate of succession and transition by girdling affected trees. It affects five-needle pines. Mountain pine beetle commonly attacks large stands of western white pine that are stressed or dying from the fungus. The fungus requires an alternate currant (*Ribes* spp.) host to complete its lifecycle. Identifiers of blister rust include swellings on branches, which may exude sap in spring. Cankers and pustules develop on tree branches and boles, and they have yellow-orange blisters (aeciospores) by midsummer. Management may include removing the currant (*Ribes* spp.), pruning affected branches, and planting genetically improved stock (Goheen, 2006). Balsam woolly adelgid (*Adelges piceae*) is common in the Cascade Mountains of Washington. It affects Pacific silver fir. The insects suck sap from the branches and boles of trees, resulting in an infection. Outbreaks may stress, deform, or kill trees and impair cone production. Most mortality is caused by a secondary attack by bark beetles. An identifier is swelling on the branches and tips of needles. Western spruce budworm (*Choristoneura occidentalis*) affects true firs, Douglas-fir, Engelmann spruce, and western larch. During outbreaks, it commonly feeds on pines and understory trees. The budworm larvae consume the current year's foliage during the first year of infestation. Multiple years of infestation result in dieback of branches, complete defoliation, and mortality. Identifiers include chewed and discolored needles of the current year's foliage. Needles turn reddish brown from the tips, and the discoloration extends along the branches toward the bole of the tree. Larvae are light brown and have a dark-colored head. They develop into adult moths by midsummer. Females lay larvae on the underside of the needles. The larvae overwinter in sheltered spots, such as in crevasses of the bark. Outbreaks typically last 8 years. They can disrupt the successional pattern in fir-dominant forests by opening up the understory, which allows less shade-tolerant species to become more dominant (Goheen, 2006).

Table 13. Representative site productivity

Common Name	Symbol	Site Index Low	Site Index High	CMAI Low	CMAI High	Age Of CMAI	Site Index Curve Code	Site Index Curve Basis	Citation
western hemlock	TSHE	91	137	215	307	90	–	–	
Pacific silver fir	ABAM	78	125	75	151	90	–	–	

Inventory data references

Other Established Classifications National vegetation classification: A3387—*Abies amabilis*-*Tsuga heterophylla*/*Vaccinium membranaceum* Cold Forest Alliance U.S. Department of Agriculture, Forest Service, plant association: • [ABAM/VAME/CLUN](#) • [ABAM/VAME/XETE](#) U.S. Department of the Interior, National Park Service, plant association: • [ABIAMA-\(PSEMEN\)/VACMEM/ACHTRI](#) • [ABIAMA-\(PSEMEN-ABIPRO\)/VACMEM/XERTEN](#)

Other references

Barnes, George H. 1962. Yield of even-aged stands of western hemlock. U.S. Department of Agriculture, Forest Service, Pacific Northwest Forest and Range Experiment Station Technical Bulletin 1273.

Cope, Amy B. 1992. *Abies amabilis*. In Fire Effects Information System. U.S. Department of Agriculture, Forest Service, Rocky Mountain Research Station, Fire Sciences Laboratory. <https://www.fs.fed.us/database/feis/plants/tree/abiama/all.html>

Crawford, P.D. 1990. *Abies amabilis*. In *Silvics of North America: Volume 1. Conifers*. U.S. Department of Agriculture, Forest Service, Agriculture Handbook 654. Pages 17-25. https://www.srs.fs.usda.gov/pubs/misc/ag_654_vol1.pdf

Crawford, R.C., C B. Chappell, C.C. Thompson, and F.J. Rocchio. 2009. Vegetation classification of Mount Rainier, North Cascades, and Olympic National Parks. Natural Resource Technical Report NPS/NCCN/NRTR-2009/211. National Park Service, Fort Collins, Colorado.

Hanley, D.P., and D.M. Baumgartner. 2002. Forest ecology in Washington. Washington State University Cooperative Extension Technical Report EB 1943.

Hanson, E.J., D.L. Azuma, and B.A. Hiserote. 2002. Site index equations and mean annual increment equations for Pacific Northwest Research Station forest inventory and analysis inventories, 1985-2001. U.S. Department of Agriculture, Forest Service, Pacific Northwest Research Station Research Note PNW-RN-533.

Henderson, J.A., R.D. Leshner, D.H. Peter, and D.C. Shaw. 1992. Field guide to the forested plant associations of the Mt. Baker-Snoqualmie National Forest. U.S. Department of Agriculture, Forest Service, Pacific Northwest Region Technical Paper R6-ECOL-TP-

028-91.

King, James E. 1966. Site index curves for Douglas-fir in the Pacific Northwest. Weyerhaeuser Company, Forestry Research Center Forestry Paper 8.

Kittel, G., D. Meidinger, and D. Faber-Langendoen. 2015. G240 *Pseudotsuga menziesii*-*Tsuga heterophylla*/*Gaultheria shallon* forest group. United States National Vegetation Classification. Federal Geographic Data Committee, Vegetation Subcommittee, Washington, D.C.

Pojar, J., and A. MacKinnon. 1994. Plants of the Pacific Northwest Coast. Lone Pine, Vancouver, British Columbia.

PRISM Climate Group. Oregon State University. Accessed February 2015. <http://prism.oregonstate.edu>

Smith, K., G. Kuhn, and L. Townsend. 2008. Culmination of mean annual increment for indicator tree species in the State of Washington. U.S. Department of Agriculture, Natural Resources Conservation Service, Technical Note Forestry-9.

Topik, C., N.M. Halverson, and D.G. Brockway. 1986. Plant associations and management guide for the western hemlock zone, Gifford Pinchot National Forest. U.S. Department of Agriculture, Forest Service, Pacific Northwest Region Technical Paper R6-ECOL-230A-1986.

United States Department of Agriculture, Natural Resources Conservation Service, and United States Department of the Interior, National Park Service. 2014. Ecological site descriptions for North Cascades National Park Complex, Washington.

Contributors

Philip Roberts

Erin Kreutz

Erik Dahlke

Approval

Kirt Walstad, 1/30/2025

Rangeland health reference sheet

Interpreting Indicators of Rangeland Health is a qualitative assessment protocol used to determine ecosystem condition based on benchmark characteristics described in the Reference Sheet. A suite of 17 (or more) indicators are typically considered in an assessment. The ecological site(s) representative of an assessment location must be known prior to applying the protocol and must be verified based on soils and climate. Current plant community cannot be used to identify the ecological site.

Author(s)/participant(s)	
Contact for lead author	
Date	05/10/2024
Approved by	
Approval date	
Composition (Indicators 10 and 12) based on	Annual Production

Indicators

1. Number and extent of rills:

2. Presence of water flow patterns:

3. Number and height of erosional pedestals or terracettes:

4. Bare ground from Ecological Site Description or other studies (rock, litter, lichen, moss, plant canopy are not bare ground):

5. Number of gullies and erosion associated with gullies:

6. Extent of wind scoured, blowouts and/or depositional areas:

7. Amount of litter movement (describe size and distance expected to travel):

8. Soil surface (top few mm) resistance to erosion (stability values are averages - most sites will show a range of values):

9. Soil surface structure and SOM content (include type of structure and A-horizon color and thickness):

10. Effect of community phase composition (relative proportion of different functional groups) and spatial distribution on infiltration and runoff:

11. Presence and thickness of compaction layer (usually none; describe soil profile features which may be mistaken for compaction on this site):

12. Functional/Structural Groups (list in order of descending dominance by above-ground annual-production or live foliar cover using symbols: >>, >, = to indicate much greater than, greater than, and equal to):

Dominant:

Sub-dominant:

Other:

Additional:

13. Amount of plant mortality and decadence (include which functional groups are expected to show mortality or decadence):

14. Average percent litter cover (%) and depth (in):

15. Expected annual annual-production (this is TOTAL above-ground annual-production, not just forage annual-production):

16. Potential invasive (including noxious) species (native and non-native). List species which BOTH characterize degraded states and have the potential to become a dominant or co-dominant species on the ecological site if their future establishment and growth is not actively controlled by management interventions. Species that become dominant for only one to several years (e.g., short-term response to drought or wildfire) are not invasive plants. Note that unlike other indicators, we are describing what is NOT expected in the reference state for the ecological site:

17. Perennial plant reproductive capability:
