

Ecological site AX006X03B100

Oregon East Cascades, Lava Plains

(Ponderosa pine - Western Juniper)

12-16 PZ

Last updated: 3/11/2025

Accessed: 08/23/2026

General information

Provisional. A provisional ecological site description has undergone quality control and quality assurance review. It contains a working state and transition model and enough information to identify the ecological site.

MLRA notes

Major Land Resource Area (MLRA): 006X–Cascade Mountains, Eastern Slope

Stretching from northern Washington to southern Oregon, MLRA 6 encompasses the mountain slopes, foothills, elevated plateaus and valleys on the eastern slopes of the Cascade mountains. This MLRA is a transitional area between the Cascade Mountains to the west and the lower lying Columbia Basalt Plateau to the east. Situated in the rain shadow of the Cascade Crest, this MLRA receives less precipitation than portions of the cascades further west and greater precipitation than the basalt plateaus to the east. Geologically, the majority of the MLRA is dominated by Miocene volcanic rocks, while the northern portion is dominated by Pre-Cretaceous metamorphic rocks and the southern portion is blanketed with a thick mantle of ash and pumice from Mount Mazama. The soils in the MLRA dominantly have a mesic, frigid, or cryic soil temperature regime, a xeric soil moisture regime, and mixed or glassy mineralogy. They generally are moderately deep to very deep, well drained, and loamy or ashy. Biologically, the MLRA is dominated by coniferous forest, large expanses of which are dominated by ponderosa pine, Douglas-fir or lodgepole pine. Areas experiencing cooler and moister conditions include grand fir, white fir, and western larch while the highest elevations include pacific silver fir, subalpine fir and whitebark pine. Economically, timber harvest and recreation are important land uses in these forests. Historically, many of these forests would have experienced relatively frequent, low and mixed severity fire favoring the development of mature forests dominated by ponderosa pine or Douglas-fir. In the southern pumice plateau forests, less frequent, higher severity fire was common and promoted the growth of large expanses of lodgepole pine forests.

LRU notes

This unit is characterized by ash mantled lava flows and glacial outwash plains on lower mountain slopes and foothills of the East Cascades in Oregon. Vegetation is largely dominated by forests of ponderosa pine with transitional dry mixed conifer forests where Douglas-fir and grand fir are sub dominant occurring in areas with greater effective precipitation. Historically, these forests have been influenced by a fire regime whereby frequent to moderately frequent, low and mixed severity fires would have favored the development of open stands of mature ponderosa pine. The climate of this unit is cool and dry with a predominately xeric soil moisture regime and frigid soil temperature regime. Geologically, underlying lithologies are dominated by Quaternary and late Tertiary basalt and basaltic andesite as well as mixed grain sediments deposited during Pleistocene glacial retreat. Unlike the nearby pumice plateau, this unit lacks the coarse pumice fragments that dominate the soil profile and cooler temperatures that favor lodgepole pine. This unit is south of the climate influences of the Columbia gorge and therefore does not support woodlands of Oregon white oak.

Classification relationships

Forested Plant Associations of the Oregon East Cascades (Simpson 2007) CPS111 – Ponderosa/bitterbrush sage/fescue CPS141 - Pinus ponderosa/Artemisia tridentate CPG135 - Pinus ponderosa/Festuca idahoensis Plant Associations of the Fremont National Forest (Hopkins 1979) CPS111 – Ponderosa/bitterbrush sage/fescue Plant Associations of the Central Oregon Pumice Zone (Volland 1985) CPS111 – Ponderosa Pine/Bitterbrush-Sagebrush/Fescue Landfire Biophysical Setting (Landfire 2007) 0710532 - Northern Rocky Mountain Ponderosa Pine Woodland and Savanna - Xeric

Ecological site concept

This site represents a transitional, dry ponderosa pine (*Pinus ponderosa*) – western juniper (*Juniperus occidentalis*) forest at the footslopes and lava plains of the East Cascades of Oregon. The Reference Plant Community is that of a mixed canopy of ponderosa pine and western juniper with a shrub layer of bitterbrush (*Purshia tridentata*) and big sagebrush (*Artemisia tridentata*) and an herbaceous layer dominated by cool-season perennial grasses. Occupying a transitional ecotone between ponderosa pine forests and dry juniper

woodlands, this site exhibits plant community characteristics of each. This site is found upon lava plains with shallow to moderately deep, coarsely textured, volcanic soils, and is distinguished from nearby stands with higher ponderosa productivity by having shallower soils and receiving less precipitation. As a very dry site with low productivity, this site also likely had a historical fire regime characterized by less frequent fires than other dry ponderosa pine sites, due to low fuel loads and continuity. This is a provisional ecological site that groups characteristics at a broad scale with little to no field verification and is subject to extensive review and revision before final approval. All data herein was developed using existing information and literature and should be considered provisional and contingent upon field validation prior to use in conservation planning.

Associated sites

AX006X03B101	Oregon East Cascades, Outwash Plains and Volcanic Uplands (Ponderosa pine / Antelope bitterbrush / Idaho Fescue) 14-18 PZ Deeper soils, higher precip range, higher cover ponderosa pine, scarcity of western juniper
--------------	---

Similar sites

AX006X03B101	Oregon East Cascades, Outwash Plains and Volcanic Uplands (Ponderosa pine / Antelope bitterbrush / Idaho Fescue) 14-18 PZ Deeper soils, higher precip range, higher cover ponderosa pine, scarcity of western juniper
--------------	---

Table 1. Dominant plant species

Tree	(1) <i>Juniperus occidentalis</i> (2) <i>Pinus ponderosa</i>
Shrub	(1) <i>Purshia tridentata</i> (2) <i>Artemisia tridentata ssp. vaseyana</i>
Herbaceous	(1) <i>Festuca idahoensis</i> (2) <i>Pseudoroegneria spicata</i>

Legacy ID

R006XB002OR

Physiographic features

This site typically occurs on gentle north aspects, and in flat or rolling positions on lava plains between juniper woodlands and ponderosa pine forests. Within these landscapes, this site is often found in interspaces between lava blisters. Slopes are commonly 2 to 15 percent but range from nearly level to 35 percent. Elevations are most commonly 3,750 to 4,250 feet (1,150 to 1,300 meters) but can range from 2,750 to 6,000 feet (850 to 1,850 m). This site is not subject to ponding or flooding and no water table is present within the upper 100 inches of soil.

Table 2. Representative physiographic features

Landforms	(1) Lava plain > Hillslope (2) Lava plateau > Depression
Flooding frequency	None

Ponding frequency	None
Elevation	1,070 – 1,520 m
Slope	0 – 20 %
Ponding depth	0 cm
Water table depth	250 cm
Aspect	W, NW, N, NE, E, SE, S, SW

Table 3. Representative physiographic features (actual ranges)

Flooding frequency	Not specified
Ponding frequency	Not specified
Elevation	840 – 1,830 m
Slope	0 – 40 %
Ponding depth	Not specified
Water table depth	Not specified

Climatic features

The annual precipitation ranges from 12 to 16 in (300 to 400 mm) which occurs mainly between the months of November and June, mostly in the form of rain and snow. The average annual air temperature is 45 degrees Fahrenheit (7° C), ranging from 41 to 46 degrees Fahrenheit (5 to 8° C) with extreme temperatures ranging from -26 to 104 degrees Fahrenheit (-32 to 40°C). The frost-free period is 60 to 90 days. The optimum period for plant growth is from April through June. The soil temperature regime is frigid, soil moisture regime is xeric. The graphs below are populated from the closest available weather station to representative site locations and are provided to indicate general climate patterns.

Table 4 Representative climatic features

Frost-free period (characteristic range)	60-90 days
Freeze-free period (characteristic range)	
Precipitation total (characteristic range)	310-410 mm

Frost-free period (average)	80 days
Freeze-free period (average)	
Precipitation total (average)	360 mm

- (1) SISTERS [USC00357857], Sisters, OR
- (2) BEND [USC00350694], Bend, OR

Influencing water features

This site is not influenced by water from adjacent wetlands or streams.

Wetland description

N/A

Soil features

The soils that typify this site concept are shallow to moderately deep, well to excessively drained and coarse textured. These soils are primarily formed in volcanic ash and colluvium, have loamy sand to sandy loam surface textures and occur over basalt and tuff. They often have substantial gravels, cobbles and stones on the surface and within the upper soil horizon which can reduce the water holding capacity of the soil. Permeability is moderately rapid and the available water holding capacity (AWC) is 3.0 to 6.0 inches for the profile. The potential for wind erosion is high and for water erosion is low. Soils of this site are largely classified as frigid vitrixerands (cool, dry soils formed in volcanic parent materials).

Representative soil associated with this site:

Fremkle, 0 – 15% slopes

Table 5. Representative soil features

Parent material	(1) Ash flow – volcanic rock (2) Colluvium – volcanic rock
Surface texture	(1) Sandy loam (2) Very cobbly sandy loam (3) Gravelly loamy sand
Family particle size	(1) Ashy (2) Ashy-skeletal
Drainage class	Well drained to excessively drained
Permeability class	Moderately rapid

Depth to restrictive layer	30 – 100 cm
Soil depth	30 – 100 cm
Surface fragment cover <=3"	0 – 50 %
Surface fragment cover >3"	0 – 50 %
Available water capacity (0-101.6cm)	3.81 – 6.6 cm
Soil reaction (1:1 water) (0-101.6cm)	6.6 – 10
Subsurface fragment volume <=3" (10.2-152.4cm)	10 – 20 %
Subsurface fragment volume >3" (10.2-152.4cm)	0 – 30 %

Table 6. Representative soil features (actual values)

Drainage class	Not specified
Permeability class	Not specified
Depth to restrictive layer	Not specified
Soil depth	Not specified
Surface fragment cover <=3"	Not specified
Surface fragment cover >3"	Not specified
Available water capacity (0-101.6cm)	3.56 – 7.62 cm

Soil reaction (1:1 water) (0-101.6cm)	Not specified
Subsurface fragment volume <=3" (10.2-152.4cm)	Not specified
Subsurface fragment volume >3" (10.2-152.4cm)	Not specified

Ecological dynamics

Reference plant community:

The Reference Native Plant Community under the natural disturbance regime, is characterized by an open stand of western juniper and ponderosa pine, with an understory dominated by bitterbrush, big sagebrush and cool season perennial grasses. Wax currant (*Ribes cereum*) may also be common in the shrub layer. The herbaceous layer includes an abundance of Idaho fescue (*Festuca idahoensis*) and bluebunch wheatgrass (*Pseudoroegneria spicata*), as well as lesser amounts of bottlebrush squirreltail (*Elymus elymoides*), rosy pussytoes (*Antennaria rosea*), and common yarrow (*Achillea millefolium*). Vegetative composition in this state is approximately 55 to 60 percent grasses, 3 percent forbs, and 20 percent shrubs and trees.

Potential plant communities will differ across the range of this site due to naturally occurring variability in soil, aspect and climate. An increase in stoniness results in an increase in drought tolerant species such as juniper, sagebrush, Sandberg bluegrass (*Poa secunda*), and bluebunch wheatgrass. Bluebunch wheatgrass increases on south exposures, Idaho fescue on more northerly aspects and Sandberg bluegrass on shallower soils. Bitterbrush is scarce where soils are saturated during the spring. Western juniper increases on shallow, well drained soils and at the lower end of the precipitation range. Ponderosa pine increases as precipitation approaches 14 inches or more.

Disturbance:

Dry ponderosa pine forests were often maintained by frequent, low severity surface fire, whereas juniper woodlands often had longer fire return intervals. As a transitional site between these ecological types, mean fire return interval estimates range between 10 to 50 years, somewhat longer than adjacent ponderosa pine forests with higher productivity and shorter than that of bordering western juniper forests and sagebrush steppe with lower productivity (Miller et al. 2019, Miller and Heyerdahl 2008, Landfire 2007). Historically, low severity fire would have decreased juniper, sagebrush and mature bitterbrush, while favoring high herbaceous cover and mature ponderosa pine development with open stand conditions. With longer time between fire, increased stocking of juniper and ponderosa pine and the development of a closed canopy, can promote an increased frequency of stand replacing fires and insect outbreaks. This condition characterizes much of the historically open canopy, dry ponderosa forests due to a history of selective logging and fire suppression (Ritchie et al 2005). Prolonged anthropogenic fire suppression may lead to cycles of overstocking and high severity fires, yet evidence is insufficient for the characterization of this pattern as an alternative state. Since this site occurs toward the lower precipitation range suitable for ponderosa pine reproduction and growth, ponderosa pine may struggle to reestablish post fire except for during climate cycles of increased moisture. In fact, across the Western US, many dry ponderosa pine forests existing toward the fringe of their climate suitability may be transitioning into grass shrub communities due to warming and drying growing season conditions (Davis et al. 2018). Ponderosa pine recovery following stand replacing fire will also be dependent on seed sources on site or in nearby forests.

Pine beetle outbreaks are another important disturbance agent in this system helping thin overstocked, closed canopy conditions (Ritchie et al 2005). Livestock grazing is a common land use of this site. Overgrazing may reduce Idaho fescue and Ross' sedge (*Carex rossii*), while allowing Thurber's needlegrass (*Achnatherum thurberianum*), bottlebrush squirreltail and rabbitbrush (*Ericameria* and *Chrysothamnus* sp.) cover to increase. Invasive annual grass may also occur on this site, with coarse textured soils likely favoring cheatgrass (*Bromus tectorum*) over medusahead (*Taeniatherum caput-medusae*) or ventenata (*Ventenata dubia*). Low forest productivity of these sites often makes them impractical for commercial timber harvest, however, given the favorable climate of this site, areas with suitable soils may also be converted to agricultural use.

The state and transition model below represents a generalized and simplified version of plant community change in response to major disturbance types in this ecological site. It does not attempt to model all of the complex interacting effects of grazing, fire and invasive species on ecosystem change and the potential restoration pathways emerging from these dynamics. Emerging evidence is suggesting that climate change is leading to hotter and drier conditions in western forests that will increase fire frequency and extent and lengthen fire seasons (Halofsky et al. 2020). When combined with the interacting impacts of fire suppression, drought, and insect outbreaks, it is

possible that this ecological system will experience unpredictable ecosystem shifts and additional alternative states. For warm and dry sites, these impacts may include the possibility of regeneration failure following wildfire disturbance (Halofsky et al. 2020). As this site is updated in future iterations, and further research informs our understanding of East Cascades oak woodlands, descriptions will include more thorough treatments of disturbance and ecological change. The Reference State is largely based on Landfire biophysical settings model 710532 (Landfire 2007).

State and transition model

Additional community tables

Table 7. Community 1.1 plant community composition

Group	Common Name	Symbol	Scientific Name	Annual Production ()	Foliar Cover (%)
Grass/Grasslike					
1	Grass and grasslike plants			476-644	
	Idaho fescue	FEID	Festuca idahoensis	353-392	–
	Ross' sedge	CARO5	Carex rossii	39-78	–
	bluebunch wheatgrass	PSSPS	Pseudoroegneria spicata ssp. spicata	39-78	–
	Sandberg bluegrass	POSE	Poa secunda	39-78	–
	Thurber's needlegrass	ACTH7	Achnatherum thurberianum	6-17	–
2	Big Bluegrass			0	
2	Other perennial grasses			6-17	
	needle and thread	HECOC8	Hesperostipa comata ssp. comata	0	–
Forb					
3	Other perennial forbs			6-22	
	lupine	LUPIN	Lupinus	0	–
	pussytoes	ANTEN	Antennaria	0	–
	phlox	PHLOX	Phlox	0	–
	buckwheat	ERIOG	Eriogonum	0	–
	beardtongue	PENST	Penstemon	0	–
	yarrow	ACHIL	Achillea	0	–
	flax	LINUM	Linum	0	–
Shrub/Vine					
4	Shrubs			62-163	
	antelope bitterbrush	PUTR2	Purshia tridentata	39-118	–
	mountain big sagebrush	ARTRV	Artemisia tridentata ssp. vaseyana	17-34	–
	wax currant	RICE	Ribes cereum	6-17	–
Tree					
5	Trees			45-196	
	western juniper	JUOC	Juniperus occidentalis	39-118	–
	ponderosa pine	PIPO	Pinus ponderosa	6-78	–

Table 8. Community 1.2 plant community composition

Group	Common Name	Symbol	Scientific Name	Annual Production ()	Foliar Cover (%)
-------	-------------	--------	-----------------	----------------------	------------------

Table 9. Community 1.3 plant community composition

Group	Common Name	Symbol	Scientific Name	Annual Production ()	Foliar Cover (%)
-------	-------------	--------	-----------------	----------------------	------------------

Table 10. Community 1.4 plant community composition

Group	Common Name	Symbol	Scientific Name	Annual Production ()	Foliar Cover (%)
-------	-------------	--------	-----------------	----------------------	------------------

Table 11. Community 1.5 plant community composition

Group	Common Name	Symbol	Scientific Name	Annual Production ()	Foliar Cover (%)
-------	-------------	--------	-----------------	----------------------	------------------

Animal community

Mule deer, Pinyon Jay, Coyotes

Inventory data references

The plant data contained in this document is derived from analysis of inventories, clipping studies, and ecological interpretation from field evaluations.

Type locality

Location 1: Deschutes County, OR	
General legal description	2-3 Miles SW of Plainview road

References

. (Date accessed). **Fire Effects Information System**. <http://www.fs.fed.us/database/feis/>.

. 2021 (Date accessed). **USDA PLANTS Database**. <http://plants.usda.gov>.

USGS. 2009 (Date accessed). **Landfire National Vegetation Dynamics Models**. <http://www.LANDFIRE.gov/index.php>.

Other references

Davis, Kimberley T.; Dobrowski, Solomon Z.; Higuera, Philip E.; Holden, Zachary A.; Veblen, Thomas T.; Rother, Monica T.; Parks, Sean A.; Sala, Anna; Maneta, Marco P. 2019. Wildfires and climate change push low-elevation forests across a critical climate threshold for tree regeneration. PNAS Latest Articles. doi: 10.1073/pnas.1815107116

Franklin, J., & Dyrness, C. Natural vegetation of Oregon and Washington. : Portland, Or., Pacific Northwest Forest and Range Experiment Station, Forest Service, U.S. Dept. of Agriculture.

Halofsky, J.E., Peterson, D.L. & Harvey, B.J. Changing wildfire, changing forests: the effects of climate change on fire regimes and vegetation in the Pacific Northwest, USA. fire ecol 16, 4 (2020). <https://doi.org/10.1186/s42408-019-0062-8>

Hopkins, W. 1979. Plant associations of the Fremont National Forest. Portland, Or. U.S. Dept. of Agriculture, Forest Service, Pacific Northwest Region.

LANDFIRE, 2007, Biophysical Settings Model Descriptions, LANDFIRE 1.1.0, U.S. Department of the Interior, USDA Forest service, Accessed 20 April 2020 at <https://www.landfire.gov/bps-models.php>

Miller, Richard F.; Chambers, Jeanne C.; Evers, Louisa; Williams, C. Jason; Snyder, Keirith A.; Roundy, Bruce A.; Pierson, Fred B. 2019. The ecology, history, ecohydrology, and management of pinyon and juniper woodlands in the Great Basin and Northern Colorado Plateau of the western United States. Gen. Tech. Rep. RMRS-GTR-403. Fort Collins, CO: U.S. Department of Agriculture, Forest Service, Rocky Mountain Research Station. 284 p.

Miller, R., & Heyerdahl, E. 2008. Fine-scale variation of historical fire regimes in sagebrush-steppe and juniper woodland: An example from California, USA. International Journal of Wildland Fire, 17(2), 245-254.

Ritchie, Martin W.; Maguire, Douglas A.; Youngblood, Andrew, Technical Coordinators. 2005. Proceedings of the Symposium on Ponderosa Pine: Issues, Trends, and Management. 2004 October 18-21; Klamath Falls, OR. Gen. Tech. Rep. PSW-GTR-198. Albany

CA: Pacific Southwest Research Station, Forest Service, U.S. Department of Agriculture. 281 p.

Simpson, M. 2007. Forested plant associations of the Oregon East Cascades. Portland, Or. U.S. Dept. of Agriculture, Forest Service, Pacific Northwest Region.

Volland, L. 1985. Plant associations of the central Oregon pumice zone. Portland, Or. U.S. Dept. of Agriculture, Forest Service, Pacific Northwest Region.

Contributors

Gene Hickman

Andrew Neary - 2020/2021 PES update of draft site

Approval

Kirt Walstad, 3/11/2025

Rangeland health reference sheet

Interpreting Indicators of Rangeland Health is a qualitative assessment protocol used to determine ecosystem condition based on benchmark characteristics described in the Reference Sheet. A suite of 17 (or more) indicators are typically considered in an assessment. The ecological site(s) representative of an assessment location must be known prior to applying the protocol and must be verified based on soils and climate. Current plant community cannot be used to identify the ecological site.

Author(s)/participant(s)	
Contact for lead author	
Date	03/11/2025
Approved by	
Approval date	
Composition (Indicators 10 and 12) based on	Annual Production

Indicators

1. Number and extent of rills:

2. Presence of water flow patterns:

3. Number and height of erosional pedestals or terracettes:

4. Bare ground from Ecological Site Description or other studies (rock, litter, lichen, moss, plant canopy are not bare ground):

5. Number of gullies and erosion associated with gullies:

6. Extent of wind scoured, blowouts and/or depositional areas:

7. Amount of litter movement (describe size and distance expected to travel):

8. Soil surface (top few mm) resistance to erosion (stability values are averages - most sites will show a range of values):

9. Soil surface structure and SOM content (include type of structure and A-horizon color and thickness):

10. Effect of community phase composition (relative proportion of different functional groups) and spatial distribution on infiltration and runoff:

11. Presence and thickness of compaction layer (usually none; describe soil profile features which may be mistaken for compaction on this site):

12. Functional/Structural Groups (list in order of descending dominance by above-ground annual-production or live foliar cover using symbols: >>, >, = to indicate much greater than, greater than, and equal to):

Dominant:

Sub-dominant:

Other:

Additional:

13. Amount of plant mortality and decadence (include which functional groups are expected to show mortality or decadence):

14. Average percent litter cover (%) and depth (in):

15. Expected annual annual-production (this is TOTAL above-ground annual-production, not just forage annual-production):

16. Potential invasive (including noxious) species (native and non-native). List species which BOTH characterize degraded states and have the potential to become a dominant or co-dominant species on the ecological site if their future establishment and growth is not actively controlled by management interventions. Species that become dominant for only one to several years (e.g., short-term response to drought or wildfire) are not invasive plants. Note that unlike other indicators, we are describing what is NOT expected in the reference state for the ecological site:
-

17. Perennial plant reproductive capability:
-