

Ecological site F006XC003WA

Cool Frigid Moist Xeric Mountain Slopes (Grand fir Cool Moist Shrub/Herb)

Last updated: 4/01/2025
Accessed: 08/31/2026

General information

Provisional. A provisional ecological site description has undergone quality control and quality assurance review. It contains a working state and transition model and enough information to identify the ecological site.

MLRA notes

Major Land Resource Area (MLRA): 006X–Cascade Mountains, Eastern Slope

Major Land Resource Area (MLRA): 006X–Cascade Mountains, Eastern Slope. Stretching from northern Washington to southern Oregon, MLRA 6 encompasses the mountain slopes, foothills, elevated plateaus and valleys on the eastern slopes of the Cascade mountains. This MLRA is a transitional area between the Cascade Mountains to the west and the lower lying Columbia Basalt Plateau to the east. Situated in the rain shadow of the Cascade Crest, this MLRA receives less precipitation than portions of the cascades further west and greater precipitation than the basalt plateaus to the east. Geologically, the majority of the MLRA is dominated by Miocene volcanic rocks, while the northern portion is dominated by Pre-Cretaceous metamorphic rocks and the southern portion is blanketed with a thick mantle of ash and pumice from Mount Mazama. The soils in the MLRA dominantly have a mesic, frigid, or cryic soil temperature regime, a xeric soil moisture regime, and mixed or glassy mineralogy. They generally are moderately deep to very deep, well drained, and loamy or ashy. Biologically, the MLRA is dominated by coniferous forest, large expanses of which are dominated by ponderosa pine, Douglas-fir or lodgepole pine. Areas experiencing cooler and moister conditions include grand fir, white fir, and western larch while the highest elevations include pacific silver fir, subalpine fir and whitebark pine. Economically, timber harvest and recreation are important land uses in these forests. Historically, many of these forests would have experienced relatively frequent, low and mixed severity fire favoring the development of mature forests dominated by ponderosa pine or Douglas-fir. In the southern pumice plateau forests, less frequent, higher severity fire was common and promoted the growth of large expanses of lodgepole pine forests.

LRU notes

Common Resource Area (CRA) 6.7 - Grand Fir Mixed Forest This LRU occurs predominantly on slopes of hills and mountains. The soils are dominantly in the Andisols and Inceptisols taxonomic order, with some Alfisols . Soil parent materials are dominantly colluvium and residuum from igneous, sedimentary, and metamorphic rock, glacial outwash, and glacial till, with a mantle or mixture of volcanic ash in the upper part. Taxonomic soil climate is primarily a frigid temperature regime and xeric moisture regime with average annual precipitation of about 50 inches. Other LRU'S where the site occurs: CRA 6.8 - Oak-Conifer Eastern Cascades - Columbia Foothills CRA 6.5 - Chiwaukum Hills and Lowlands CRA 6.6 - Yakima Plateau and Slopes

Classification relationships

CWS525 - grand fir/vanilla leaf (ABGR/ACTR)

Ecological site concept

This site is recognized as Grand Fir Cool Moist Shrub/Herb. It occurs on steep north slopes, benches, and bottoms. Elevations ranges from 2000 to 4800 feet. Precipitation ranges from 40 to 60 inches. Its main tree components are grand fir, Douglas-fir, western larch, and western white pine. Engelmann spruce can be present on the moist bottom sites and ponderosa pine can occur on the warmer end. Lodgepole pine will occur after severe fire. Main understory tree species are grand fir and Douglas-fir. Key understory species are vanilla leaf, Cascade Oregon grape, twinflower, western prince's pine, pachistima, sidebells pyrola, big huckleberry, baldhip rose, queencup beadlily, western rattlesnake plantain, trillium, and wester starflower. The USFS plant association for this site is Grand fir/vanilla leaf (ABGR/ACTR)

Associated sites

F006XD005WA	Frigid Xeric Mountain Slopes and Plateaus (Grand fir Warm Moderately Dry Shrub) Has western hazel in the understory.
F006XB003WA	Frigid Xeric Mountain Slopes (Grand fir Warm Moderately Dry Low Shrub/Herb) On warmer and drier sites.
F006XC003WA	Cool Frigid Moist Xeric Mountain Slopes (Grand fir Cool Moist Shrub/Herb) Moister site.

Similar sites

F006XD002WA	Cool Frigid Xeric Ashy Slopes (Grand fir Cool Dry Grass) Drier and higher elevation.
--------------------	--

Table 1. Dominant plant species

Tree	(1) <i>Abies grandis</i>
Shrub	Not specified
Herbaceous	Not specified

Physiographic features

This ecological site is on mountain slopes, glacial outwash terraces, and hillslopes. It is typically found between 1400 and 4200 feet in elevation on all aspects. Slope gradients range from 5 to 45 percent.

Table 2. Representative physiographic features

Landforms	(1) Mountains > Mountain slope (2) Hills > Outwash terrace (3) Hillslope
Flooding frequency	None
Ponding frequency	None
Elevation	430 – 1,280 m
Slope	10 – 50 %

Aspect	W, NW, N, NE, E, SE, S, SW
--------	----------------------------

Table 3. Representative physiographic features (actual ranges)

Flooding frequency	Not specified
Ponding frequency	Not specified
Elevation	150 – 1,520 m
Slope	0 – 60 %

Climatic features

Mean Annual Air Temperature

Total Range: 41 to 48 degrees Fahrenheit

Central tendency: 43 to 47 degrees Fahrenheit

Table 4 Representative climatic features

Frost-free period (characteristic range)	80-140 days
Freeze-free period (characteristic range)	
Precipitation total (characteristic range)	1,020-1,520 mm
Frost-free period (actual range)	70-160 days
Freeze-free period (actual range)	
Precipitation total (actual range)	640-1,910 mm

Climate stations used

- (1) EASTON [USC00452384], Cle Elum, WA

Influencing water features

This site is not influenced by water from a wetland or stream.

Wetland description

N/A

Soil features

This ecological site is associated with several soil mapunit components. The components are dominantly Humic Vitrixerands in the Andisols order and Andic Dystrochrepts in the Inceptisols order. Soils are dominantly moderately deep to very deep and have average available water capacity of about 4.8 inches (10.9 cm) in the 0 to 40-inches (0 to 100 cm) depth range. Soil parent material is dominantly volcanic ash over colluvium and residuum from volcanic, granitic, and metamorphic rock, and glacial outwash.

Dominant Soil Series: Bearrun, Bertolotti, Chemawa, Culving, Saydab, Sugarbowl

Parent Materials:

Kind – volcanic ash, colluvium, residuum, glacial outwash

Origin – volcanic, granitic, and metamorphic rock; mixed sources

Table 5. Representative soil features

Surface texture	(1) Ashy loam (2) Ashy sandy loam
Family particle size	(1) Ashy (2) Loamy-skeletal (3) Ashy-skeletal
Drainage class	Well drained
Soil depth	50 – 150 cm
Surface fragment cover ≤3"	0 – 20 %
Surface fragment cover >3"	0 – 10 %
Available water capacity (0-101.6cm)	6.1 – 19.56 cm
Electrical conductivity (Depth not specified)	Not specified
Sodium adsorption ratio (Depth not specified)	Not specified
Subsurface fragment volume ≤3" (Depth not specified)	0 – 60 %
Subsurface fragment volume >3" (Depth not specified)	0 – 30 %

Ecological dynamics

This site is recognized as Grand Fir Cool Moist Shrub/Herb. It occurs on steep north slopes, benches, and bottoms. Elevations ranges from 2000 to 4800 feet. Precipitation ranges from 40 to 60 inches. Its main tree components are grand fir, Douglas-fir, western larch, and western white pine. Engelmann spruce can be present on the moist bottom sites and ponderosa pine can occur on the warmer end. Lodgepole pine will occur after severe fire. These moist cool areas have a long fire intervals allowing grand fir to become a main overstory species. Older Douglas-fir and western larch will be present. Western white pine would have been more prominent if not for the white pine blister rust reducing its numbers.

Mixed severity fires will create a mosaic of stands with older remnant patches along with early seral species like larch and Douglas-fir. Main understory tree species are grand fir and Douglas-fir.

Key understory species are vanilla leaf, Cascade Oregon grape, twinflower, western prince's pine, pachistima, sidebells pyrola, big huckleberry, baldhip rose, queencup beadlelily, western rattlesnake plantain, trillium, and wester starflower.

The USFS plant association for this site is Grand fir/vanilla leaf (ABGR/ACTR)

Main insect and disease problems are armillaria root rot, Douglas-fir beetle, fir engraver, spruce budworm, Douglas-fir tussock moth, western pine beetle and mountain pine beetle. Dwarf mistletoe may occur in western larch.

State and transition model

Additional community tables

Table 6. Community 1.1 plant community composition

Group	Common Name	Symbol	Scientific Name	Annual Production ()	Foliar Cover (%)
-------	-------------	--------	-----------------	----------------------	------------------

Table 7. Community 1.2 plant community composition

Group	Common Name	Symbol	Scientific Name	Annual Production ()	Foliar Cover (%)
-------	-------------	--------	-----------------	----------------------	------------------

Table 8. Community 1.3 plant community composition

Group	Common Name	Symbol	Scientific Name	Annual Production ()	Foliar Cover (%)
-------	-------------	--------	-----------------	----------------------	------------------

Table 9. Representative site productivity

Common Name	Symbol	Site Index Low	Site Index High	CMAI Low	CMAI High	Age Of CMAI	Site Index Curve Code	Site Index Curve Basis	Citation
Douglas-fir	PSME	60	115	—	—	—	—	—	
ponderosa pine	PIPO	100	135	—	—	—	—	—	
western larch	LAOC	50	65	—	—	—	—	—	
lodgepole pine	PICO	80	90	—	—	—	—	—	
grand fir	ABGR	75	105	—	—	—	—	—	

Inventory data references

Forest Service Plant Association: CWS525 - grand fir/vanilla leaf (ABGR/ACTR) Information presented here has been derived from NRCS data. Field observations from range trained personnel were also used. Other sources used as references include USDA NRCS Water and Climate Center, USDA NRCS National Range and Pasture Handbook, and USDA NRCS Soil Surveys from various counties.

Other references

Field Guide for Forest Plant Associations of the Wenatchee National Forest. Lillybridge et al. PNW-GTR-359. October 1995
Washington Natural Heritage Program. Ecosystems of Washington State, A Guide to Identification, Rocchio and Crawford, 2015 – East Cascades Mesic Montane Mixed-Conifer Forest and Woodland
USDA, NRCS Forest-Soil Eco classifications.
Field reconnaissance by Gary Kuhn, ACES Forester and Steve Campbell NRCS Soil Scientist. July 2019

Contributors

Approval

Kirt Walstad, 4/01/2025

Rangeland health reference sheet

Interpreting Indicators of Rangeland Health is a qualitative assessment protocol used to determine ecosystem condition based on benchmark characteristics described in the Reference Sheet. A suite of 17 (or more) indicators are typically considered in an assessment. The ecological site(s) representative of an assessment location must be known prior to applying the protocol and must be verified based on soils and climate. Current plant community cannot be used to identify the ecological site.

Author(s)/participant(s)	
Contact for lead author	
Date	04/01/2025
Approved by	
Approval date	
Composition (Indicators 10 and 12) based on	Annual Production

Indicators

1. Number and extent of rills:

2. Presence of water flow patterns:

3. Number and height of erosional pedestals or terracettes:

4. Bare ground from Ecological Site Description or other studies (rock, litter, lichen, moss, plant canopy are not bare ground):

5. Number of gullies and erosion associated with gullies:

6. Extent of wind scoured, blowouts and/or depositional areas:

7. Amount of litter movement (describe size and distance expected to travel):

8. Soil surface (top few mm) resistance to erosion (stability values are averages - most sites will show a range of values):

9. Soil surface structure and SOM content (include type of structure and A-horizon color and thickness):

10. Effect of community phase composition (relative proportion of different functional groups) and spatial distribution on infiltration and runoff:

11. Presence and thickness of compaction layer (usually none; describe soil profile features which may be mistaken for compaction on this site):

12. Functional/Structural Groups (list in order of descending dominance by above-ground annual-production or live foliar cover using symbols: >>, >, = to indicate much greater than, greater than, and equal to):

Dominant:

Sub-dominant:

Other:

Additional:

13. Amount of plant mortality and decadence (include which functional groups are expected to show mortality or decadence):

14. Average percent litter cover (%) and depth (in):

15. Expected annual annual-production (this is TOTAL above-ground annual-production, not just forage annual-production):

16. Potential invasive (including noxious) species (native and non-native). List species which BOTH characterize degraded states and have the potential to become a dominant or co-dominant species on the ecological site if their future establishment and growth is not actively controlled by management interventions. Species that become dominant for only one to several years (e.g., short-term response to drought or wildfire) are not invasive plants. Note that unlike other indicators, we are describing what is NOT expected in the reference state for the ecological site:

17. Perennial plant reproductive capability:
