

Ecological site F006XY702OR East Crater Lake Pumice Drainages

Accessed: 08/16/2026

General information

Provisional. A provisional ecological site description has undergone quality control and quality assurance review. It contains a working state and transition model and enough information to identify the ecological site.

Associated sites

F006XY707OR	East Crater Lake Stratovolcano Slopes This site doesnot have a brush component in it.
-------------	---

Similar sites

F003XY707OR	HC High Pumice Basin This site is in a different MLRA and higher precip zone.
-------------	---

Table 1. Dominant plant species

Tree	(1) <i>Pinus contorta</i>
Shrub	(1) <i>Ribes cereum</i> (2) <i>Purshia tridentata</i>
Herbaceous	(1) <i>Achnatherum occidentale ssp. occidentale</i>

Physiographic features

This site is on flat to gently sloping excessively drained, deep soils.

Table 2. Representative physiographic features

Landforms	(1) Ash flow
Flooding frequency	None
Ponding frequency	None
Elevation	1,370 – 1,830 m

Slope	0 – 10 %
Water table depth	150 cm
Aspect	Aspect is not a significant factor

Climatic features

Winter are long, cold and snowy. Snow makes up a large amount of the effective precipitation. Summers days are warm, and nights are cool. Summer precipitation comes as infreqesnt rain storms. Summer thunderstorms can drop moderately heavy amounts of rain, but only for a short period of time.

Table 3 Representative climatic features

Frost-free period (average)	50 days
Freeze-free period (average)	110 days
Precipitation total (average)	760 mm

Influencing water features

Soil features

This site is found on volcanic pumice and ash flow deposits.

Table 4. Representative soil features

Surface texture	(1) Paragravelly loamy sand (2) Ashy loamy sand
Family particle size	(1) Sandy
Drainage class	Excessively drained
Permeability class	Rapid
Soil depth	150 cm
Surface fragment cover <=3"	0 – 20 %
Surface fragment cover >3"	0 – 30 %

Available water capacity (0-101.6cm)	9.14 – 15.24 cm
Calcium carbonate equivalent (0-101.6cm)	Not specified
Electrical conductivity (0-101.6cm)	Not specified
Sodium adsorption ratio (0-101.6cm)	Not specified
Soil reaction (1:1 water) (0-101.6cm)	5.6 – 6.5
Subsurface fragment volume <=3" (Depth not specified)	30 – 50 %
Subsurface fragment volume >3" (Depth not specified)	20 – 40 %

Ecological dynamics

Lodgepole pine is the only tree specie in the overstory of the historic climax plant community. The position of this site, flat basins, is influenced by cold air drainage which affects tree seedling survival. Lodgepole seedlings are the only ones that have adapted to these conditions.

Heavy to very heavily stocked stands are susceptible to mountain pine beetle infestations. If an outbreak occurs all lodgepole can be killed. If seedlings are present then a new stand will start. Large areas killed may take many years to return to a lodgepole forest. Planting seedlings will quicken the successional process.

State and transition model

Additional community tables

Table 5. Community 1.1 plant community composition

Group	Common Name	Symbol	Scientific Name	Annual Production ()	Foliar Cover (%)
-------	-------------	--------	-----------------	----------------------	------------------

Type locality

Location 1: Klamath County, OR

Latitude	42° 51'2"
Longitude	121° 58'34"

Contributors

Craig Ziegler

Rangeland health reference sheet

Interpreting Indicators of Rangeland Health is a qualitative assessment protocol used to determine ecosystem condition based on benchmark characteristics described in the Reference Sheet. A suite of 17 (or more) indicators are typically considered in an assessment. The ecological site(s) representative of an assessment location must be known prior to applying the protocol and must be verified based on soils and climate. Current plant community cannot be used to identify the ecological site.

Author(s)/participant(s)	
Contact for lead author	
Date	
Approved by	
Approval date	
Composition (Indicators 10 and 12) based on	Annual Production

Indicators

1. Number and extent of rills:

2. Presence of water flow patterns:

3. Number and height of erosional pedestals or terracettes:

4. Bare ground from Ecological Site Description or other studies (rock, litter, lichen, moss, plant canopy are not bare ground):

5. Number of gullies and erosion associated with gullies:

6. Extent of wind scoured, blowouts and/or depositional areas:

7. Amount of litter movement (describe size and distance expected to travel):

8. Soil surface (top few mm) resistance to erosion (stability values are averages - most sites will show a range of values):

9. Soil surface structure and SOM content (include type of structure and A-horizon color and thickness):

10. Effect of community phase composition (relative proportion of different functional groups) and spatial distribution on infiltration and runoff:

11. Presence and thickness of compaction layer (usually none; describe soil profile features which may be mistaken for compaction on this site):

12. Functional/Structural Groups (list in order of descending dominance by above-ground annual-production or live foliar cover using symbols: >>, >, = to indicate much greater than, greater than, and equal to):

Dominant:

Sub-dominant:

Other:

Additional:

13. Amount of plant mortality and decadence (include which functional groups are expected to show mortality or decadence):

14. Average percent litter cover (%) and depth (in):

15. Expected annual annual-production (this is TOTAL above-ground annual-production, not just forage annual-production):

16. Potential invasive (including noxious) species (native and non-native). List species which BOTH characterize degraded states and have the potential to become a dominant or co-dominant species on the ecological site if their future establishment and growth is not actively controlled by management interventions. Species that become dominant for only one to several years (e.g., short-term response to drought or wildfire) are not invasive plants. Note that unlike other indicators, we are describing what is NOT expected in the reference state for the ecological site:

17. Perennial plant reproductive capability:
