

Ecological site F006XY709OR

Mesic Xeric Foothills 20-25 PZ

Last updated: 4/01/2025
Accessed: 08/15/2026

General information

Provisional. A provisional ecological site description has undergone quality control and quality assurance review. It contains a working state and transition model and enough information to identify the ecological site.

MLRA notes

Major Land Resource Area (MLRA): 006X–Cascade Mountains, Eastern Slope

Stretching from northern Washington to southern Oregon, MLRA6 encompasses the mountain slopes, foothills, elevated plateaus and valleys on the eastern slopes of the Cascade mountains. This MLRA is a transitional area between the Cascade Mountains to the west and the lower lying Columbia Basalt Plateau to the east. Situated in the rain shadow of the Cascade Crest, this MLRA receives less precipitation than portions of the cascades further west and greater precipitation than the basalt plateaus to the east. Geologically, the majority of the MLRA is dominated by Miocene volcanic rocks, while the northern portion is dominated by Pre-Cretaceous metamorphic rocks and the southern portion is blanketed with a thick mantle of ash and pumice from Mount Mazama. The soils in the MLRA dominantly have a mesic, frigid, or cryic soil temperature regime, a xeric soil moisture regime, and mixed or glassy mineralogy. They generally are moderately deep to very deep, well drained, and loamy or ashy. Biologically, the MLRA is dominated by coniferous forest, large expanses of which are dominated by ponderosa pine, Douglas-fir or lodgepole pine. Areas experiencing cooler and moister conditions include grand fir, white fir, and western larch while the highest elevations include pacific silver fir, subalpine fir and whitebark pine. Economically, timber harvest and recreation are important land uses in these forests. Historically, many of these forests would have experienced relatively frequent, low and mixed severity fire favoring the development of mature forests dominated by ponderosa pine or Douglas-fir. In the southern pumice plateau forests, less frequent, higher severity fire was common and promoted the growth of large expanses of lodgepole pine forests.

LRU notes

This unit is characterized by ash mantled lava flows and glacial outwash plains on lower mountain slopes and foothills of the East Cascades in Oregon. Vegetation is largely dominated by forests of ponderosa pine with transitional dry mixed conifer forests where Douglas-fir and grand fir are sub dominant occurring in areas with greater effective precipitation. Historically, these forests have been influenced by a fire regime whereby frequent to moderately frequent, low and mixed severity fires would have favored the development of open stands of mature ponderosa pine. The climate of this unit is cool and dry with a predominately xeric soil moisture regime and frigid soil temperature regime. Geologically, underlying lithologies are dominated by Quaternary and late Tertiary basalt and basaltic andesite as well as mixed grain sediments deposited during Pleistocene glacial retreat. Unlike the nearby pumice plateau, this unit lacks the coarse pumice fragments that dominate the soil profile and cooler temperatures that favor lodgepole pine. This unit is south of the climate influences of the Columbia gorge and therefore does not support woodlands of Oregon white oak.

Classification relationships

Forested Plant Associations of the Oregon East Cascades (Simpson 2007): CPC733 - *Pinus ponderosa*-*Calocedrus decurrens*/*Arctostaphylos patula* Plant Associations of the Commercial Forest of the Warm Springs Indian Reservation (Marsh 1987): PIPO/PUTR-ARPA - *Pinus ponderosa* /*Purshia tridentata* - *Arctostaphylos patula* Landfire Biophysical Setting (Landfire 2007): 0710531 - Northern Rocky Mountain Ponderosa Pine Woodland and Savanna

Ecological site concept

This site represents a warm ponderosa pine site in the foothills of the Eastern Cascades of Oregon. An overstory dominated by ponderosa pine (*Pinus ponderosa*) with a lesser components of incense cedar (*Calocedrus decurrens*), occasional sub-canopy Douglas-fir (*Pseudotsuga menziesii*) and a shrub understory of greenleaf manzanita (*Arctostaphylos patula*) largely characterize the visual aspect of this reference plant community. Occupying a dry, warm ponderosa pine elevation band, this site transitions into ponderosa-bitterbrush (*Purshia tridentata*) and Oregon white oak (*Quercus garryana*) communities at its lower elevations and Douglas-fir, grand fir (*Abies grandis*) – white fir (*Abies concolor*) communities at its upper elevations. In comparison to adjacent warm, dry ponderosa pine forest

types with a bitterbrush dominated understory, this site receives greater precipitation (20 to 25 in) creating conditions favorable to the less drought tolerant Douglas-fir. In comparison to other sites with ponderosa pine overstories and manzanita dominated shrub layers, this site has a mesic, rather than frigid or cryic, soil temperature regime allowing for the persistence of incense cedar. While this site may occasionally host minor tree species in the understory such as grand fir and western larch (*Larix occidentalis*), unlike sites where these species may become dominant overtime, this site receives lower precipitation and has a more frequent fire rotation. Mixed and low severity fire were historically a critical element of the disturbance regime of this site, acting to thin crowded understories and allow mature fire-resistant ponderosa pine and Douglas-fir stands to attain an open, savanna-like appearance. This is a provisional ecological site that groups characteristics at a broad scale with little to no field verification and is subject to extensive review and revision before final approval. All data herein was developed using existing information and literature and should be considered provisional and contingent upon field validation prior to use in conservation planning.

Associated sites

AX006X03C105	Oregon East Cascades, Volcanic Uplands and Mountainbases (Ponderosa pine - Douglas-fir - (Grand fir) / Snowbrush) 20-30 PZ Adjacent upslope positions, frigid soil temperature regime, CEVE common
F006XY710OR	Mesic Xeric Foothills 14-20 PZ Adjacent downslope positions, ARPA less common
R006XB208OR	Shallow Slopes 14-20 PZ Shallow soils, south aspects, non-forested plant community

Similar sites

F006XB802OR	Mesic Xeric North Slopes 15-25 PZ Occupying steep north slopes, lower precipitation, PSME common
F006XY710OR	Mesic Xeric Foothills 14-20 PZ Lower precipitation, lower elevations
AX006X03C105	Oregon East Cascades, Volcanic Uplands and Mountainbases (Ponderosa pine - Douglas-fir - (Grand fir) / Snowbrush) 20-30 PZ Higher precipitation, frigid soil temperature regime, CADE less common

Table 1. Dominant plant species

Tree	(1) <i>Pinus ponderosa</i>
Shrub	(1) <i>Purshia tridentata</i> (2) <i>Arctostaphylos patula</i>
Herbaceous	(1) <i>Festuca idahoensis</i>

Physiographic features

This site is largely found occupying benches and side slopes of the foothills of the Eastern Cascades in Oregon. Its range is largely restricted to areas north of Black Butte, mainly on the Confederated Tribes of the Warm Springs reservation. Elevations commonly range from 2,500 to 3,500 feet (750 to 1,050 m). Slopes are most often nearly level to 12 percent but can be as steep as 40 percent. This site is found on all aspects. This site is not subject to ponding or flooding and no water table is present within 100 inches of the soil surface.

Table 2. Representative physiographic features

Landforms	(1) Foothills > Bench (2) Foothills > Hillside or mountainside
Flooding frequency	None
Ponding frequency	None
Elevation	760 – 1,070 m
Slope	0 – 10 %
Ponding depth	0 cm
Water table depth	250 cm
Aspect	W, NW, N, NE, E, SE, S, SW

Table 3. Representative physiographic features (actual ranges)

Flooding frequency	Not specified
Ponding frequency	Not specified
Elevation	760 – 1,070 m
Slope	0 – 40 %
Ponding depth	Not specified
Water table depth	Not specified

Climatic features

The average annual precipitation ranges from 20 to 25 inches (500 to 625 mm) which occurs mainly between the months of November and June, mostly in the form of rain and snow. The average annual air temperature ranges from 45 to 46 degrees Fahrenheit (7 to 8 °C) and the frost-free period ranges from 80 to 110 days. Soil temperature regime is mesic, soil moisture regime is xeric. The graphs below

are populated from the closest available weather station to representative site locations and are provided to indicate general climate patterns.

Table 4 Representative climatic features

Frost-free period (characteristic range)	80-110 days
Freeze-free period (characteristic range)	
Precipitation total (characteristic range)	510-640 mm
Frost-free period (average)	100 days
Freeze-free period (average)	
Precipitation total (average)	560 mm

- (1) SISTERS [USC00357857], Sisters, OR
- (2) DUFUR [USC00352440], Dufur, OR

Influencing water features

This site is not influenced by water from a wetland or stream.

Wetland description

N/A

Soil features

The soils that typify this site concept are moderately deep to very deep over paralithic bedrock. These are primarily well drained, volcanic soils with surface layers with components of volcanic ash over colluvium and residuum derived from basalt, andesite and sedimentary rock. Surface soil textures are commonly loams. These soils sometimes contain substantial cobbles and stones in the upper surface layer. Volcanic ash will increase water holding capacity and productivity of these soils while rock fragments will decrease water holding capacity. Taxonomic classification of these soils is predominately Vitrandic Argixerolls (Grassland soils with influences of volcanic materials). See Hehe and Teewee (high precipitation phases) for representative soils associated with this site concept.

Table 5. Representative soil features

Parent material	(1) Volcanic ash (2) Colluvium – volcanic rock (3) Residuum – volcanic rock (4) Colluvium – sedimentary rock (5) Residuum – sedimentary rock
-----------------	--

Surface texture	(1) Loam (2) Very cobbly loam (3) Very stony loam
Family particle size	(1) Loamy-skeletal (2) Fine-loamy
Drainage class	Well drained
Permeability class	Moderately slow to moderate
Depth to restrictive layer	50 – 200 cm
Soil depth	50 – 200 cm
Surface fragment cover <=3"	0 – 20 %
Surface fragment cover >3"	0 – 20 %
Available water capacity (0-101.6cm)	8.64 – 13.21 cm
Soil reaction (1:1 water) (0-101.6cm)	5.6 – 6.5
Subsurface fragment volume <=3" (10.2-152.4cm)	0 – 10 %
Subsurface fragment volume >3" (10.2-152.4cm)	30 – 50 %

Table 6. Representative soil features (actual values)

Drainage class	Not specified
Permeability class	Not specified

Depth to restrictive layer	Not specified
Soil depth	Not specified
Surface fragment cover <=3"	Not specified
Surface fragment cover >3"	Not specified
Available water capacity (0-101.6cm)	4.83 – 15.75 cm
Soil reaction (1:1 water) (0-101.6cm)	5.6 – 10
Subsurface fragment volume <=3" (10.2-152.4cm)	0 – 20 %
Subsurface fragment volume >3" (10.2-152.4cm)	10 – 50 %

Ecological dynamics

Reference Plant community:

The reference native plant community under the natural disturbance regime, is characterized by an overstory composed of ponderosa pine, with a lesser component of incense cedar, and a sub-canopy dominated by greenleaf manzanita. The tendency for incense cedar to attain codominance in the overstory sets this forested plant community apart from other dry ponderosa sites, and may be a unique feature of this site within the Oregon Eastern Cascades (Simpson 2007). In addition to greenleaf manzanita, bitterbrush and snowbrush (*Ceanothus velutinus*) are common shrubs and Douglas-fir may occur in the sub-canopy. Herbaceous species usually do not make up a dominant proportion of vegetative cover but may include Idaho fescue (*Festuca idahoensis*), Ross' sedge (*Carex rossii*), bluebunch wheatgrass (*Pseudoroegneria spicata*), prairie junegrass (*Koeleria macrantha*), Sandberg bluegrass (*Poa secunda*), arrowleaf balsamroot (*Balsamorhiza sagittata*) and tailcup lupine (*Lupinus caudatus*).

Disturbance:

A lack of specific fire rotation information is available for this forest type, yet warm ponderosa pine forests were historically subject to frequent surface fires primarily ignited by lightning strikes. Little evidence exists for local Native American cultural burning in this dry forest type, a practice more commonly used in wetter mixed conifer forests of the area (Steen-Adams et al. 2019). These low intensity fires would have decreased the density of young regenerating understory trees, which may otherwise act as ladder fuels to ignite crown fires and lead to stand replacing events. Overtime, frequent low intensity fires, as well as occasional mixed severity fires, would have favored the development of mature, even-aged ponderosa pine stands with open canopies (Landfire 2007). Fire resistant ponderosa is well adapted to these conditions, developing increasing fire resistance with age by growing thick bark and self-thinning lower limbs (Fryer 2008). Incense cedar is much less fire tolerant than ponderosa, yet more shade tolerant. Given long periods without fire, it may become more prominent in the overstory (Tollefson 2008). Douglas-fir is moisture limited on this site and therefore often occupies northern aspects and declines without overstory ponderosa (Marsh et al. 1987). Manzanita and snowbrush will increase following fire and may form shrub fields on sites with higher moisture following stand replacing fires. With longer time between fire, increased development of understory fuels, especially shade tolerant trees such as incense cedar and Douglas-fir, along with the development of a closed canopy, can promote an increased frequency of stand replacing fires and insect outbreaks. This condition characterizes much of the historically

open canopy, dry ponderosa forests due to a history of selective logging and fire suppression (Ritchie et al 2005). Prolonged anthropogenic fire suppression may lead to cycles of overstocking and high severity fires, yet evidence is insufficient for the characterization of this pattern as an alternative state.

Historically, low elevation ponderosa forests were harvested extensively for timber products (Ritchie et al. 2005). Sites with higher productivity are especially attractive for commercial timber harvesting which will have varying effects on stand structure and composition depending on harvest practices. Selective logging of large shade tolerant ponderosa trees may favor the development of stands dominated by more shade tolerant tree species such as incense cedar and Douglas-fir overtime (Hessberg et al. 2005). This site has been historically utilized for livestock grazing and hosts elk and deer wintering range despite low herbaceous cover (Marsh et al. 1987). Grazing pressure will alter herbaceous cover and composition. Idaho fescue will likely decrease while bottlebrush squirreltail (*Elymus elymoides*), cheatgrass (*Bromus tectorum*), Sandberg bluegrass and Ross' sedge will likely increase (Marsh et al. 1987). Advanced invasion of exotic annual grasses in dry pine sites may increase fine fuel loads and fuel continuity, thereby increasing fire rotation and altering seasonality.

The state and transition model below represents a generalized and simplified version of forest change in response to fire in this ecological site. It does not attempt to model the complex effects of forestry practices, insect outbreaks or climate change on ecosystem function or process. Emerging evidence is suggesting that climate change is leading to hotter and drier conditions in western forests that will increase fire frequency and extent and lengthen fire seasons (Halofsky et al. 2020). When combined with the interacting impacts of fire suppression, drought, and insect outbreaks, it is possible that this ecological system will experience unpredictable ecosystem shifts and additional alternative states. For warm and dry sites, this may include the possibility of regeneration failure following wildfire disturbance (Halofsky et al. 2020). The reference state of the current model is largely based on Landfire biophysical settings model Rocky Mountain Ponderosa Pine Woodland and Savanna, 0710531 (Landfire 2007).

State and transition model

Additional community tables

Table 7. Community 1.1 plant community composition

Group	Common Name	Symbol	Scientific Name	Annual Production ()	Foliar Cover (%)
-------	-------------	--------	-----------------	----------------------	------------------

Table 8. Community 1.2 plant community composition

Group	Common Name	Symbol	Scientific Name	Annual Production ()	Foliar Cover (%)
-------	-------------	--------	-----------------	----------------------	------------------

Table 9. Community 1.3 plant community composition

Group	Common Name	Symbol	Scientific Name	Annual Production ()	Foliar Cover (%)
-------	-------------	--------	-----------------	----------------------	------------------

Table 10. Community 1.4 plant community composition

Group	Common Name	Symbol	Scientific Name	Annual Production ()	Foliar Cover (%)
-------	-------------	--------	-----------------	----------------------	------------------

Table 11. Community 1.5 plant community composition

Group	Common Name	Symbol	Scientific Name	Annual Production ()	Foliar Cover (%)
-------	-------------	--------	-----------------	----------------------	------------------

References

. (Date accessed). **Fire Effects Information System**. <http://www.fs.fed.us/database/feis/>.

. 2021 (Date accessed). **USDA PLANTS Database**. <http://plants.usda.gov>.

. 2021 (Date accessed). **USNVC [United States National Vegetation Classification]. 2019. United States National Vegetation Classification Database, V2.03. Federal Geographic Data Committee, Vegetation Subcommittee, Washington DC.. USNVC: <http://usnvc.org/>.**

Other references

Franklin, J., & Dyrness, C. Natural vegetation of Oregon and Washington. : Portland, Or., Pacific Northwest Forest and Range Experiment Station, Forest Service, U.S. Dept. of Agriculture.

Fryer, Janet L. 2018. *Pinus ponderosa* var. *benthamiana*, P. p. var. *ponderosa*: Ponderosa pine. In: Fire Effects Information System, [Online]. U.S. Department of Agriculture, Forest Service, Rocky Mountain Research Station, Missoula Fire Sciences Laboratory (Producer). Available: www.fs.fed.us/database/feis/plants/tree/pinponp/all.html

Halofsky, J.E., Peterson, D.L. & Harvey, B.J. Changing wildfire, changing forests: the effects of climate change on fire regimes and vegetation in the Pacific Northwest, USA. *fire ecol* 16, 4 (2020). <https://doi.org/10.1186/s42408-019-0062-8>

Hessburg, P.F., Agee, J.K., & Franklin, J.F. (2005). Dry forests and wildland fires of the inland Northwest USA: Contrasting the landscape ecology of the pre-settlement and modern eras.

LANDFIRE, 2007, Biophysical Settings Model Descriptions, LANDFIRE 1.1.0, U.S. Department of the Interior, USDA Forest service, Accessed 20 April 2020 at <https://www.landfire.gov/bps-models.php>

Marsh, Frank; Helliwell, Richard; Rodgers, Jean. 1987. Plant Association Guide for the Commercial Forest of the Warm Springs Indian Reservation. Portland, Or. U.S. Dept. of Agriculture, Forest Service, Pacific Northwest Region.

Ritchie, Martin W.; Maguire, Douglas A.; Youngblood, Andrew, Technical Coordinators. 2005. Proceedings of the Symposium on Ponderosa Pine: Issues, Trends, and Management. 2004 October 18-21; Klamath Falls, OR. Gen. Tech. Rep. PSW-GTR-198. Albany CA: Pacific Southwest Research Station, Forest Service, U.S. Department of Agriculture. 281 p.

Simpson, M. 2007. Forested plant associations of the Oregon East Cascades. Portland, Or. U.S. Dept. of Agriculture, Forest Service, Pacific Northwest Region.

Steen-Adams, Michelle M.; Charnley, Susan; McLain, Rebecca J.; Adams, Mark D.O.; Wendel, Kendra L. 2019. Traditional knowledge of fire use by the Confederated Tribes of Warm Springs in the eastside Cascades of Oregon. *Forest Ecology and Management*. 450: 117405-. <https://doi.org/10.1016/j.foreco.2019.06.002>.

Tollefson, Jennifer E. 2008. *Calocedrus decurrens*. In: Fire Effects Information System, [Online]. U.S. Department of Agriculture, Forest Service, Rocky Mountain Research Station, Fire Sciences Laboratory (Producer). Available: <https://www.fs.fed.us/database/feis/plants/tree/caldec/all.html> [2020, May 28].

Volland, L. 1985 Plant associations of the central Oregon pumice zone. Portland, Or. U.S. Dept. of Agriculture, Forest Service, Pacific Northwest Region.

Contributors

Andrew Neary - Original PES site concept

Approval

Kirt Walstad, 4/01/2025

Rangeland health reference sheet

Interpreting Indicators of Rangeland Health is a qualitative assessment protocol used to determine ecosystem condition based on benchmark characteristics described in the Reference Sheet. A suite of 17 (or more) indicators are typically considered in an assessment. The ecological site(s) representative of an assessment location must be known prior to applying the protocol and must be verified based on soils and climate. Current plant community cannot be used to identify the ecological site.

Author(s)/participant(s)	
Contact for lead author	
Date	04/01/2025
Approved by	

Approval date	
Composition (Indicators 10 and 12) based on	Annual Production

Indicators

1. Number and extent of rills:

2. Presence of water flow patterns:

3. Number and height of erosional pedestals or terracettes:

4. Bare ground from Ecological Site Description or other studies (rock, litter, lichen, moss, plant canopy are not bare ground):

5. Number of gullies and erosion associated with gullies:

6. Extent of wind scoured, blowouts and/or depositional areas:

7. Amount of litter movement (describe size and distance expected to travel):

8. Soil surface (top few mm) resistance to erosion (stability values are averages - most sites will show a range of values):

9. Soil surface structure and SOM content (include type of structure and A-horizon color and thickness):

10. Effect of community phase composition (relative proportion of different functional groups) and spatial distribution on infiltration and runoff:

11. Presence and thickness of compaction layer (usually none; describe soil profile features which may be mistaken for compaction on this site):

12. Functional/Structural Groups (list in order of descending dominance by above-ground annual-production or live foliar cover using symbols: >>, >, = to indicate much greater than, greater than, and equal to):

Dominant:

Sub-dominant:

Other:

Additional:

13. Amount of plant mortality and decadence (include which functional groups are expected to show mortality or decadence):

14. Average percent litter cover (%) and depth (in):

15. Expected annual annual-production (this is TOTAL above-ground annual-production, not just forage annual-production):

16. Potential invasive (including noxious) species (native and non-native). List species which BOTH characterize degraded states and have the potential to become a dominant or co-dominant species on the ecological site if their future establishment and growth is not actively controlled by management interventions. Species that become dominant for only one to several years (e.g., short-term response to drought or wildfire) are not invasive plants. Note that unlike other indicators, we are describing what is NOT expected in the reference state for the ecological site:

17. Perennial plant reproductive capability:
