

Ecological site R008XY720WA

Riparian Complex

Last updated: 5/23/2025
Accessed: 09/01/2026

General information

Provisional. A provisional ecological site description has undergone quality control and quality assurance review. It contains a working state and transition model and enough information to identify the ecological site.

MLRA notes

Major Land Resource Area (MLRA): 008X–Columbia Plateau

MLRA 8 encompasses about 50,100 square kilometers mainly in Washington and Oregon, with a small area in Idaho. This MLRA is characterized by loess hills, surrounding scablands, and alluvial deposits. This MLRA consists mostly of Miocene Columbia River Basalt covered with up to 200 feet of loess and volcanic ash. The dominant soil order in this MLRA is Mollisols. Soils in this MLRA dominantly have a mesic temperature regime, a xeric moisture regime, and mixed minerology.

Classification relationships

Major Land Resource Area (MLRA): 8 – Columbia Plateau LRU – Common Resource Areas (CRA): 8.1 - Channeled Scablands 8.2 - Loess Islands 8.3 - Okanogan Drift Hills 8.4 - Moist Pleistocene Lake Basins 8.5 - Moist Yakima Folds 8.6 - Lower Snake and Clearwater Canyons 8.7 - Okanogan Valley

Ecological site concept

NRCS Washington has never had an ecological site for riparian areas in MLRA 7, 8 and 9, so this is a starting point. Riparian complex is extremely general and will continue to be a work in progress for some time. There are precipitation ranges for these sites, but precipitation is not the main driver of the site development. Diagnostics: A bottomland tree-shrub site, Riparian Complex sits in the floodplain position of the landscape as a narrow, linear corridor along perennial streams or spring fed reaches on intermittent streams. It also occurs as a small patch near ponds, lakes and springs, or on bottoms, depressions and basins. This site receives seasonal flooding and/or discharging groundwater from sites uphill. Riparian complex is part of the lotic or flowing water ecosystem. Soils are often cobbly and well drained, so they remain saturated for only a short period. The soils are not hydric, nor is this ecological site a wetland. In the sagebrush steppe and grassland steppe regions, bunchgrasses, shrubs and forbs are common, but trees are rare. Riparian Complex stands out because of trees and tall shrubs. Principle Vegetative Drivers: The vegetative expression of Riparian complex is driven by three situations. First, this site receives off-site water - overbank flooding and surface runoff for some sites or, discharging groundwater for sites influenced by springs or seeps. Second, the soils are deep and have unrestricted rooting. Third, the soils are well drained and supports trees and shrubs instead of aquatic plants

Associated sites

R008XY435WA	Loamy 14-20 PZ Goldendale Prairie
R008XY402WA	Loamy Bottom 10-16 PZ

Similar sites

R008XY402WA	Loamy Bottom 10-16 PZ
-------------	-----------------------

R008XY980WA	Wet Meadow
--------------------	-------------------

Table 1. Dominant plant species

Tree	(1) <i>Populus balsamifera ssp. trichocarpa</i> (2) <i>Populus tremuloides</i>
Shrub	(1) <i>Symphoricarpos albus</i> (2) <i>Cornus sericea ssp. sericea</i>
Herbaceous	Not specified

Physiographic features

The landscape is part of the Columbia basalt plateau. Riparian complex sites occur as a narrow, linear corridor on floodplains and terraces along perennial streams or spring fed reaches of intermittent streams. There are also Riparian Complex patches on draws, basins, depressions, and near ponds, lakes or springs. In the upland setting ecological sites are often expansive, and thus, can be delineated and separated on aerial photos. But in the landscape position of bottoms, basins and depressions this is rarely the case as small changes in soil chemistry, the water table and elevation or aspect results in significant changes in plant community composition. In short distances there are often big swings of available water holding capacity, and soils can go from hydric to non-hydric, or from saline-sodic to not. So, in bottoms, riparian areas and depressions, ecological sites and community phases occur as small spots, strips and patches, or as narrow rings around vernal ponds. Generally, in a matter of steps one can walk across several ecological sites. On any given site location, two or more of these sites occur as a patchwork – Loamy Bottom, Alkali Terrace, Sodic Flat, Wet Meadow, Herbaceous Wetland and Riparian Complex. These ecological sites may need to be mapped as a complex when doing resource inventory.

Physiographic Division: Intermontane Plateau

Physiographic Province: Columbia Plateau

Physiographic Sections: Walla Walla Plateau Section

Landscapes: Valleys, hills and plateaus

Landform: floodplains, drainageways on concave positions

Table 2. Representative physiographic features

Slope shape across	(1) Concave
Slope shape up-down	(1) Concave
Landforms	(1) Valley (2) Hills (3) Plateau (4) Flood plain (5) Drainageway
Flooding frequency	Rare to frequent

Ponding frequency	None to frequent
Elevation	310 – 1,100 m
Slope	0 %
Water table depth	30 – 150 cm
Aspect	W, NW, N, NE, E, SE, S, SW

Table 3. Representative physiographic features (actual ranges)

Flooding frequency	Not specified
Ponding frequency	Not specified
Elevation	Not specified
Slope	0 – 10 %
Water table depth	Not specified

Climatic features

The climate is characterized by moderately cold, wet winters, and hot, dry summers, with limited precipitation due to the rain shadow effect of the Cascades. Taxonomic soil climate is either xeric (12 to 16 inches PPT) or aridic moisture regimes (10 to 12 inches PPT) with a mesic temperature regime.

Table 4 Representative climatic features

Frost-free period (characteristic range)	110-160 days
Freeze-free period (characteristic range)	
Precipitation total (characteristic range)	250-410 mm
Frost-free period (actual range)	90-200 days
Freeze-free period (actual range)	
Precipitation total (actual range)	

Influencing water features

A plant's ability to grow on a site and overall plant production is determined by soil-water-plant relationships:

1. Whether rain and melting snow run off-site or infiltrate into the soil
2. Whether soil condition remain aerobic or become saturated and anaerobic
3. How quickly the soil reaches the wilting point

Most sites experience overbank flooding and surface runoff. But for sites influenced by springs or seeps, discharging groundwater is the important driver. The soils are deep, well drained, and often cobbly, and thus, remain saturated for only a short period in late winter to early spring. With adequate cover of live plants and litter, there are no water infiltrating restrictions on Riparian Complex.

Soil features

This ecological site components are dominantly Cumulic, Fluventic and Torrifluventic taxonomic subgroups of Haploxerolls, Endoaquolls great groups of the Mollisols taxonomic order but also includes some Entisols. Soils are dominantly very deep. Average available water capacity of about five inches (12.7 cm) in the zero to 40 inches (zero to 100 cm) depth range.

Soil parent material is dominantly mixed alluvium.

The associated soils are Bridgewater, Colville, Weirman, Xerofluvents and similar soils.

Dominant soil surface is silt loam to cobbly loamy coarse sand, with ashy modifier sometimes occurring as well.

Dominant particle-size class is fine-silty to coarse-loamy but includes limited sandy-skeletal.

Table 5. Representative soil features

Parent material	(1) Alluvium
Surface texture	(1) Silt loam (2) Cobbly loamy coarse sand
Family particle size	(1) Fine-silty (2) Coarse-loamy
Depth to restrictive layer	50 – 150 cm
Soil depth	150 cm
Surface fragment cover <=3"	10 %
Surface fragment cover >3"	10 %
Available water capacity (0-101.6cm)	12.7 cm

Calcium carbonate equivalent (Depth not specified)	0 – 10 %
Electrical conductivity (Depth not specified)	Not specified
Sodium adsorption ratio (Depth not specified)	0 – 10
Soil reaction (1:1 water) (0-25.4cm)	6.1 – 8.4
Subsurface fragment volume <=3" (Depth not specified)	20 %
Subsurface fragment volume >3" (Depth not specified)	10 %

Table 6. Representative soil features (actual values)

Depth to restrictive layer	Not specified
Soil depth	Not specified
Surface fragment cover <=3"	0 – 30 %
Surface fragment cover >3"	0 – 40 %
Available water capacity (0-101.6cm)	2.54 – 23.11 cm
Calcium carbonate equivalent (Depth not specified)	Not specified
Electrical conductivity (Depth not specified)	Not specified

Sodium adsorption ratio (Depth not specified)	Not specified
Soil reaction (1:1 water) (0-25.4cm)	Not specified
Subsurface fragment volume <=3" (Depth not specified)	0 – 50 %
Subsurface fragment volume >3" (Depth not specified)	0 – 30 %

Ecological dynamics

Riparian complex produces about 2,000 pounds/acre of biomass annually up to 4.5 feet
Understory production is dependent on canopy cover.
Low canopy = high production. High canopy = low production

Ecological Dynamics Overview:

Riparian areas are heavily influenced by fluvial processes. This provisional ecological site concept is composed of a variety of different riverine systems and will require more detailed field investigations to refine the site concepts and likely develop several new sites that are correlated to similar geologic structure and processes, hydrologic regimes, and vegetation characteristics. This ecological site concept captures variety of typical riparian vegetation expressions. The band of riparian vegetation may be broader if part of a larger river system, or narrower if part of a small stream system.

Basic Understanding of Riparian Systems – Abiotic Factors/Primary Disturbance
(Kendra Moseley, NRCS):

Riparian forests are a complex interaction of many various physical and biologic factors, including function of valley morphology, physical processes, vegetative legacies, and life history strategies. The watershed geomorphology and physical processes form the basis for understanding the spatial extent of the riparian forests, which includes the valley shape, hillslope processes, fluvial processes, soil processes, and hydrologic processes. Soil development within alluvial environments is highly variable. Frequent erosional and depositional disturbances from flooding create a complex mosaic of soil conditions in the active floodplain that fundamentally influences vegetation colonization and establishment. Well-drained soil or recently deposited mineral alluvium may be found adjacent to very poorly drained organic soils in abandoned high-flow channels. This variability in soil conditions is a major factor in maintaining the high plant diversity typical of riparian ecological sites.

The disturbances that drive this ecological site concept are dependent on the type, frequency, predictability, extent, magnitude, and timing of the disturbance. The fluvial processes that are dominant in this riparian ecological site concept include stream power, basal shear stress, channel migration, and sediment deposition. The characteristic vegetation pattern of these low-gradient valleys is maintained by fluvial disturbances and geomorphology. The amount of force exerted on the channel bed and vegetation growing in the active channel and floodplain during a flood is a product of fluid density, gravitational acceleration, flow depth, and water surface slope.

Basic understanding of Channel Evolution
(From Stream Visual Assessment, Version 2, December 2009:

Some understanding of stream geomorphology helps our understanding of the ecological dynamics of these fluvial systems.

Channel slope is directly related to topography, geology, sinuosity, bed material and watershed size. Straight stream channels are indicative of strong geologic structure (bedrock) or human control. Braided streams have multiple interwoven channels. Meandering channels are highly variable and sinuous.

The shape of a stream channel changes constantly, imperceptibly, or dramatically, depending on the condition of the stream corridor

(channel, riparian area, and flood plain) and how it transports water and materials. Channel condition is a description of the geomorphic stage of the channel as it adjusts its shape relative to its flood plain. Channel adjustments resulting in a dramatic drop in streambed elevation (incision or degradation) or excessive deposition of bedload that raises the bed elevation (aggradation) affect the degree of bank shear and often decrease stream channel stability. Such channel adjustments can have substantial effects on the condition of streams, adjacent riparian areas, associated habitats, and their biota. For example, the greater the incision in a channel, the more it is separated from its flood plain, both physically and ecologically. Conversely, the greater the aggradation, the wider and shallower a stream becomes, which can affect riparian vegetation, surface water temperatures, and stream and riparian habitat features.

Conceptual models of how a channel evolves or adjusts over time illustrate the sequence of geomorphic changes in a stream that result from disturbances in the watershed. Such sequences are useful for evaluating trends in channel condition. The stages of the Schumm Channel Evolution Model (CEM), as shown in figure 3, provide a visual orientation of the pattern of streambed adjustment in an incising stream, its gradual detachment from the existing flood plain, and eventual formation of a new flood plain at a lower elevation. A similar model by Simon (1989) is also described in the Stream Corridor Restoration Handbook (FISRWG 1998) available in most NRCS field offices.

Evolution of a Stream:

Stage I channels are generally stable and have frequent interaction with their flood plains. The relative stability of the streambed and banks is because the stream and its flood plain are connected, and flooding occurs at regular intervals (Q2). Consequently, the stream's banks and flood plain are well vegetated. The Stage I channel undergoes initial incision. In Stage II the bed degrades, and banks are stable. During Stage III the bed aggrades, banks are unstable, and the channel goes through the widening phase. Stage IV is the stabilizing phase where the bed continues aggrading, but the banks are stable. Then in Stage V the slow aggrading continues but now banks are stable, and the new floodplain is forming. The new floodplain is lower and narrower than the original floodplain.

Especially on larger streams, variations on riparian areas (clumps and strips) are common. Variations are controlled by valley width, sediment type and overbank flooding. Clumps and strips of dominant trees species found on riparian areas can include – black cottonwood, aspen, water birch, white alder and several different willow species. Smaller streams are less diverse than larger streams. Overbank flooding and gravel bars are required for tree regeneration for many riparian trees, especially for black cottonwood.

Ecological Dynamics (from Frank G.):

The riparian woodland sites in MLRAs 7, 8, and 9 will vary based on their locations in various watersheds of the region and available moisture and soil texture/depth of the sites. Sites that have sufficient soil texture and depth will have a tall tree component like black cottonwood and/or quaking aspen. Sites that are drier will have a short tree or shrub component like hawthorn, chokecherry or mockorange. Also, the size of the watershed that drains an area and adjacent land cover will influence the amount of water entering the riparian area.

Riparian woodlands are essential to keep streambanks stable, stream temperatures cool, and buffer potential impacts from adjacent land uses. Riparian woodlands provide valuable wildlife habitat for a variety of species. Native riparian woodlands can recover from low intensity fires through tree/shrub root sprouting or seed regeneration. Serious impacts can occur when these areas are continually overgrazed or removed/reduced for another land use.

Fire Ecology:

Forested riparian areas (Riparian Woodlands) are unique ecosystems which provide critical habitat for many species of terrestrial and aquatic species. Forested riparian zones occur adjacent to two broadly associated upland ecosystems. In dryer areas, the adjacent upland type was historically steppe or shrub/steppe, and in areas of increasing rainfall, the adjacent upland was dominated by coniferous forests. In large scale watersheds with appreciable climatic gradients, the riparian zone passes through and connects these two upland types.

Riparian areas are subject to a number of ecologic disturbance influences, including wildfire. Wildfire frequency (the historic intervals between various types of fires), severity (the impacts of any given fire episode), and the type (surface, crown, mixed) of fire occurring within these riparian zones was highly variable.

Riparian ecosystems that were adjacent to prairie vegetation experienced a relatively frequent surface fire return interval. Fire typically entered into the riparian zone from fires originating in the upland grass dominated vegetation. Black cottonwood hardwoods were likely the only true tree species, growing along with adapted understory shrubs and grass species.

Riparian areas adjacent to upland conifer forests have a wider expression of tree and understory species. These areas would typically burn in the same upland fire event, and experience the impacts similar to the surrounding upland forests, especially when upland the fire episode was more severe.

Fire impacts, and natural adaptation to common riparian tree, shrub and grass species:

Black cottonwood is easily killed by fire, but coppice sprouting is common. Fire can improve seedling establishment by increasing light penetration and exposing mineral soil to allow seedling establishment if moisture is available. It can endure on site as well as invade following fire.

Ponderosa pine is resistant to fire by the development of thick, platy bark which begins to form at a young age. It naturally prunes the lower bole, decreasing the likelihood of crown ignition, and the open nature of the needles resist damage when upper crow fires do occur. It is also long lived, and will regenerate on open, sunny areas which are often produced by surface fire.

Quaking aspen regenerates by clonal regeneration following light to moderately severe fire.

Various species of willows, sedges and grasses regrow following all but severe fires, from root crowns or underground rhizomes.

Lupine, manzanita and Ceanothus produce hard shelled seeds that require fire scarification to germinate and become established.

Many other plant species produce lightweight seeds which easily travel on wind currents to disturbed areas.

In the past century, in what is referred to as the period of post-European settlement, riparian forests have undergone many human caused changes. Logging up to the edge of the stream, conversion to agriculture, fire exclusion, forest fragmentation and especially unregulated livestock grazing have all contributed to the degradation and proper function of riparian woodlands. These factors have all contributed to changes from the historic fire regimes and impacts of native riparian woodland forests.

State and transition model

Additional community tables

Table 7. Community 1.1 plant community composition

Group	Common Name	Symbol	Scientific Name	Annual Production ()	Foliar Cover (%)
-------	-------------	--------	-----------------	----------------------	------------------

Table 8. Community 1.2 plant community composition

Group	Common Name	Symbol	Scientific Name	Annual Production ()	Foliar Cover (%)
-------	-------------	--------	-----------------	----------------------	------------------

Table 9. Community 2.1 plant community composition

Group	Common Name	Symbol	Scientific Name	Annual Production ()	Foliar Cover (%)
-------	-------------	--------	-----------------	----------------------	------------------

Table 10. Community 3.1 plant community composition

Group	Common Name	Symbol	Scientific Name	Annual Production ()	Foliar Cover (%)
-------	-------------	--------	-----------------	----------------------	------------------

Other references

Crawford, Rex, Riparian Vegetation Classification of the Columbia Basin, Washington, Washington State Department of Natural Resources, Natural Heritage Report 2003-03, March 2003

Dwire, Kathleen A. and Kauffman, J. Boone. Fire and riparian ecosystems in landscapes of the western USA. In Forest Ecology and Management 178 (2003) p. 61-74. 14.

Environmental Protection Agency, map of Level III and IV Ecoregions of Washington, June 2010 Natural Resources Conservation Service, map of Common Resource Areas of Washington, 2003 NRCS Stream Visual Assessment Protocol, Version 2, December 2009

Rocchio, Joseph & Crawford, Rex C., Ecological Systems of Washington State. A Guide to Identification. Washington State Department of Natural Resources, October 2015. Pages 156-161

Inter-Mountain Basin Big Sagebrush. Soil Conservation Service, Range Sites for MLRA 9 from 1980s

Trees and Shrubs for Riparian Plantings. USDA, NRCS Washington. Plant Materials Technical Note #24.

Contributors

Gary Kuhn
Frank Gariglio
Kevin Guinn
R. Fleenor
K. Paup-Lefferts
C. Smith

Approval

Kirt Walstad, 5/23/2025

Acknowledgments

Provisional Site Author: Kevin Guinn Technical Team: K. Moseley, G. Fults, R. Fleenor, W. Keller, C. Smith, K. Bomberger, C. Gaines, K. Paup-Lefferts

Rangeland health reference sheet

Interpreting Indicators of Rangeland Health is a qualitative assessment protocol used to determine ecosystem condition based on benchmark characteristics described in the Reference Sheet. A suite of 17 (or more) indicators are typically considered in an assessment. The ecological site(s) representative of an assessment location must be known prior to applying the protocol and must be verified based on soils and climate. Current plant community cannot be used to identify the ecological site.

Author(s)/participant(s)	
Contact for lead author	
Date	05/08/2025
Approved by	
Approval date	
Composition (Indicators 10 and 12) based on	Annual Production

Indicators

1. Number and extent of rills:

2. Presence of water flow patterns:

3. Number and height of erosional pedestals or terracettes:

4. Bare ground from Ecological Site Description or other studies (rock, litter, lichen, moss, plant canopy are not bare ground):

5. Number of gullies and erosion associated with gullies:

6. Extent of wind scoured, blowouts and/or depositional areas:

7. Amount of litter movement (describe size and distance expected to travel):

8. Soil surface (top few mm) resistance to erosion (stability values are averages - most sites will show a range of values):

9. Soil surface structure and SOM content (include type of structure and A-horizon color and thickness):

10. Effect of community phase composition (relative proportion of different functional groups) and spatial distribution on infiltration and runoff:

11. Presence and thickness of compaction layer (usually none; describe soil profile features which may be mistaken for compaction on this site):

12. Functional/Structural Groups (list in order of descending dominance by above-ground annual-production or live foliar cover using symbols: >>, >, = to indicate much greater than, greater than, and equal to):

Dominant:

Sub-dominant:

Other:

Additional:

13. Amount of plant mortality and decadence (include which functional groups are expected to show mortality or decadence):

14. Average percent litter cover (%) and depth (in):

15. Expected annual annual-production (this is TOTAL above-ground annual-production, not just forage annual-production):

16. Potential invasive (including noxious) species (native and non-native). List species which BOTH characterize degraded states and have the potential to become a dominant or co-dominant species on the ecological site if their future

establishment and growth is not actively controlled by management interventions. Species that become dominant for only one to several years (e.g., short-term response to drought or wildfire) are not invasive plants. Note that unlike other indicators, we are describing what is NOT expected in the reference state for the ecological site:

17. Perennial plant reproductive capability:
