

Ecological site R010XC036OR

SR Shallow Cool

9-12 PZ

Accessed: 08/27/2026

General information

Provisional. A provisional ecological site description has undergone quality control and quality assurance review. It contains a working state and transition model and enough information to identify the ecological site.

Associated sites

R010XC030OR	SR Cool 9-12 PZ SR Cool 9-12 PZ
R010XC038OR	SR Very Shallow 9-12 PZ SR Very Shallow 9-12" PZ
R010XC043OR	SR South 9-12 PZ SD South 9-12 PZ
BX010X00C050	Shallow South 9-12 PZ Snake River Warm Plains SR Shallow South 9-12" PZ
R010XC065OR	SR Cool North 9-12 PZ SR Cool North 9-12 PZ

Similar sites

R010XC030OR	SR Cool 9-12 PZ SR Cool 9-12 PZ (deeper soil, higher production)
--------------------	--

Table 1. Dominant plant species

Tree	Not specified
Shrub	(1) <i>Artemisia tridentata ssp. wyomingensis</i>
Herbaceous	(1) <i>Festuca idahoensis</i>

Physiographic features

This site occurs on terraces, tablelands and rolling uplands. Slopes range from 2 to 12%. Elevations typically range from 3,500 to 4,200 feet.

Table 2. Representative physiographic features

Landforms	(1) Terrace (2) Hill
Elevation	1,070 – 1,280 m
Slope	0 – 10 %
Aspect	Aspect is not a significant factor

Climatic features

The annual precipitation ranges from 9 to 12 inches, most of which occurs in the form of snow during the months of December through March. Localized convection storms occasionally occur during the summer. The soil temperature regime is cool mesic with a mean air temperature of 45 degrees F. Temperature extremes range from 100 to -20 degrees F. The frost free period ranges from 60 to 100 days. The optimum growth period for plant growth is April through June.

Table 3 Representative climatic features

Frost-free period (average)	100 days
Freeze-free period (average)	0 days
Precipitation total (average)	310 mm

Influencing water features

Soil features

The soils of this site are typically shallow and well drained. Typically the surface layer is a silt loam to stony clay loam about 6 inches thick. The subsoil is a silty clay loam to gravelly clay about 5 inches thick. Depth to bedrock or an indurated pan is 10 to 20 inches. Permeability is slow. The available water holding capacity (AWC) is about 2 to 4 inches for the profile. The erosion potential is moderate to severe.

Table 4. Representative soil features

Surface texture	(1) Gravelly silt loam (2) Stony clay loam
Family particle size	(1) Clayey

Drainage class	Well drained
Permeability class	Slow
Soil depth	30 – 50 cm
Available water capacity (0-101.6cm)	5.08 – 10.16 cm

Ecological dynamics

The potential native plant community is dominated by Wyoming big sagebrush and Idaho fescue. Bluebunch wheatgrass, Thurber needlegrass and Sandberg bluegrass are common. A variety of forbs and other shrubs are present. Vegetative composition of the community is approximately 70 percent grasses, 10 percent forbs and 20 percent shrubs. Approximate ground cover is 40-60 percent (basal and crown).

Range in Characteristics:

Idaho fescue dominance is marginal as it is near the edge of its ecological amplitude and has low site resiliency. Bluebunch wheatgrass increases on slight south and west exposures. Thurber needlegrass increases on shallower soils and soils with gravelly surfaces. Shrubs will increase over fractured substratum. Production will increase on deeper soils and at the upper end of the precipitation zone.

Response to Disturbance:

When the condition of the site deteriorates as a result of over grazing Idaho fescue rapidly decreases. Wyoming big sagebrush, Sandberg bluegrass and squirreltail increase. With continued overgrazing Wyoming big sagebrush and Sandberg bluegrass become dominant. When present, bulbous bluegrass invades. Annual invasion is limited unless major ground disturbance occurs. With further deterioration, bare ground increases and excessive erosion reduces the site productivity and contributes to downstream sedimentation.

States: ARTRW/POSE-Bare Ground

State and transition model

Additional community tables

Table 5. Community 1.1 plant community composition

Group	Common Name	Symbol	Scientific Name	Annual Production ()	Foliar Cover (%)
Grass/Grasslike					
1	Dominant, perennial deep-rooted bunchgrass			392-549	
	Idaho fescue	FEID	<i>Festuca idahoensis</i>	392-549	–
2	Sub-dominant, deep rooted perennial grasses			55-141	
	bluebunch wheatgrass	PSSPS	<i>Pseudoroegneria spicata ssp. spicata</i>	39-78	–
	Thurber's needlegrass	ACTH7	<i>Achnatherum thurberianum</i>	16-63	–
4	Dominant, perennial shallow-rooted grass			24-63	
	Sandberg bluegrass	POSE	<i>Poa secunda</i>	24-63	–
5	Other perennial grasses			17-71	
	threadleaf sedge	CAFI	<i>Carex filifolia</i>	0-24	–
	squirreltail	ELEL5	<i>Elymus elymoides</i>	11-24	–
	prairie Junegrass	KOMA	<i>Koeleria macrantha</i>	6-24	–
Forb					

7	Dominant, perennial forbs			30-86	
	arrowleaf balsamroot	BASA3	<i>Balsamorhiza sagittata</i>	8-24	–
	fleabane	ERIGE2	<i>Erigeron</i>	6-16	–
	buckwheat	ERIOG	<i>Eriogonum</i>	6-16	–
	desertparsley	LOMAT	<i>Lomatium</i>	6-16	–
	phlox	PHLOX	<i>Phlox</i>	6-16	–
9	Other perennial forbs			30-93	
	common yarrow	ACMI2	<i>Achillea millefolium</i>	3-8	–
	milkvetch	ASTRA	<i>Astragalus</i>	3-8	–
	stoneseed	LITHO3	<i>Lithospermum</i>	3-8	–
	lupine	LUPIN	<i>Lupinus</i>	3-8	–
	brittle pricklypear	OPFR	<i>Opuntia fragilis</i>	0-8	–
	tapertip hawksbeard	CRAC2	<i>Crepis acuminata</i>	3-8	–
	larkspur	DELPH	<i>Delphinium</i>	0-6	–
	bitter root	LERE7	<i>Lewisia rediviva</i>	2-6	–
	sagebrush buttercup	RAGL	<i>Ranunculus glaberrimus</i>	2-6	–
	brodiaea	BRODI	<i>Brodiaea</i>	2-6	–
	agoseris	AGOSE	<i>Agoseris</i>	2-6	–
	onion	ALLIU	<i>Allium</i>	2-6	–
	pussytoes	ANTEN	<i>Antennaria</i>	2-6	–
	bushy bird's beak	CORA5	<i>Cordylanthus ramosus</i>	0-3	–
	rough eyelashweed	BLSC	<i>Blepharipappus scaber</i>	0-3	–
Shrub/Vine					
11	Dominant evergreen shrub			78-157	
	Wyoming big sagebrush	ARTRW8	<i>Artemisia tridentata ssp. wyomingensis</i>	78-157	–
15	Other shrubs			24-126	
	basin big sagebrush	ARTRT	<i>Artemisia tridentata ssp. tridentata</i>	16-39	–
	wild crab apple	PERA4	<i>Peraphyllum ramosissimum</i>	8-24	–
	antelope bitterbrush	PUTR2	<i>Purshia tridentata</i>	0-16	–
	horsebrush	TETRA3	<i>Tetradymia</i>	0-8	–
	yellow rabbitbrush	CHVI8	<i>Chrysothamnus viscidiflorus</i>	0-8	–
	spiny hopsage	GRSP	<i>Grayia spinosa</i>	0-8	–
	granite prickly phlox	LIPU11	<i>Linanthus pungens</i>	0-8	–
	scabland sagebrush	ARRI2	<i>Artemisia rigida</i>	0-8	–
	threetip sagebrush	ARTR4	<i>Artemisia tripartita</i>	0-8	–

Animal community

Livestock Grazing: This site is suitable for livestock grazing use in the spring, early summer and fall under a planned grazing system. Use should be postponed until the soils are firm enough to prevent trampling damage and soil compaction. Grazing management should be keyed for Idaho fescue. Deferred grazing or rest is recommended at least once every three years. Native Wildlife Associated with the Potential Climax Community: This site is commonly used by pronghorn antelope, mule deer, rabbits, rodents, upland birds and various predators. It is a preferred site for sage grouse leks, nesting, rearing and wintering. Antelope and mule deer make excellent use of the site for winter and spring forage.

Hydrological functions

The soils of this site are typically in an upland topographic position. They have moderate runoff potential and medium infiltration rates when the hydrologic cover is high. Hydrologic cover is high when the Idaho fescue deep rooted bunchgrass component is >70 percent of potential. Under lower seral conditions runoff potential is high. This occurs when deep rooted perennial bunchgrass cover is low and bare

ground increases.

Contributors

BLM/SCS Team - Burns, A. Bahn, G. Kuehl, H. Barrett
Bob Gillaspy
M. Parks (OSU)
T.Bloomer, B.Gillaspy, E. Petersen, A.Bahn.

Rangeland health reference sheet

Interpreting Indicators of Rangeland Health is a qualitative assessment protocol used to determine ecosystem condition based on benchmark characteristics described in the Reference Sheet. A suite of 17 (or more) indicators are typically considered in an assessment. The ecological site(s) representative of an assessment location must be known prior to applying the protocol and must be verified based on soils and climate. Current plant community cannot be used to identify the ecological site.

Author(s)/participant(s)	Bruce Frannsen and Jeff Repp
Contact for lead author	State Rangeland Management Specialist for NRCS in Oregon.
Date	08/07/2012
Approved by	
Approval date	
Composition (Indicators 10 and 12) based on	Annual Production

Indicators

1. Number and extent of rills: none, moderate to severe sheet & rill erosion hazard

2. Presence of water flow patterns: None

3. Number and height of erosional pedestals or terracettes: None to very few (some frost heaving)

4. Bare ground from Ecological Site Description or other studies (rock, litter, lichen, moss, plant canopy are not bare ground):
10-25%

5. Number of gullies and erosion associated with gullies: None

6. Extent of wind scoured, blowouts and/or depositional areas: None, slight wind erosion hazard

7. Amount of litter movement (describe size and distance expected to travel): Fine - limited movement

8. Soil surface (top few mm) resistance to erosion (stability values are averages - most sites will show a range of values):
Moderate to significant resistance to erosion: aggregate stability = 4-6

9. Soil surface structure and SOM content (include type of structure and A-horizon color and thickness): Shallow well drained
gravelly clay loam (6 inches thick): Low to moderate OM (0-2%)

10. Effect of community phase composition (relative proportion of different functional groups) and spatial distribution on
infiltration and runoff: Moderate ground cover (30-50%) and gentle slopes (0-15%) moderately limit rainfall impact and overland flow

11. Presence and thickness of compaction layer (usually none; describe soil profile features which may be mistaken for
compaction on this site): None

12. Functional/Structural Groups (list in order of descending dominance by above-ground annual-production or live foliar
cover using symbols: >>, >, = to indicate much greater than, greater than, and equal to):

Dominant: Idaho fescue > Wyoming big sagebrush > Bluebunch wheatgrass > other grasses > forbs > other shrubs

Sub-dominant:

Other:

Additional:

13. Amount of plant mortality and decadence (include which functional groups are expected to show mortality or decadence):
Normal decadence and mortality expected

14. Average percent litter cover (%) and depth (in):

15. Expected annual annual-production (this is TOTAL above-ground annual-production, not just forage annual-production):
Faorable: 1100, Normal: 600, Unfavorable: 300 lbs/acre/year at high RSI (HCPC)

16. Potential invasive (including noxious) species (native and non-native). List species which BOTH characterize degraded states and have the potential to become a dominant or co-dominant species on the ecological site if their future establishment and growth is not actively controlled by management interventions. Species that become dominant for only one to several years (e.g., short-term response to drought or wildfire) are not invasive plants. Note that unlike other indicators, we are describing what is NOT expected in the reference state for the ecological site: Western Juniper readily invades the site. Cheatgrass and Medusahead invade sites that have lost deep rooted perennial grass functional groups.
-

17. Perennial plant reproductive capability: All species should be capable of reproducing annually
-