

## Ecological site R011XA003ID

### Shallow Loam

### 8-12 PZ ARTRT/PSSPS

Last updated: 4/06/2020

Accessed: 09/22/2026

#### General information

**Provisional.** A provisional ecological site description has undergone quality control and quality assurance review. It contains a working state and transition model and enough information to identify the ecological site.

#### MLRA notes

Major Land Resource Area (MLRA): 011X–Snake River Plains

Major Land Resource Area (MLRA): 011X – Snake River Plains Precipitation or Climate Zone: 8-12" P.Z.

#### Classification relationships

*Artemisia tridentata*/*Agropyron spicatum* HT in "Hironaka, M., M.A. Fosberg, A. H. Winward. 1983. Sagebrush- Grass Habitat Types of Southern Idaho. University of Idaho. Moscow, Idaho. Bulletin Number 35." Land Resource Region: B (Northwest Wheat and Range)  
MLRA: 11 (Snake River Plains) EPA Eco Region: Level III (Snake River Plain)

#### Ecological site concept

Site does not receive additional moisture Soils are: Not saline or saline sodic Shallow, with >35% coarse fragments (by volume), skeletal not strongly or violently effervescent in the surface mineral 10" Surface textures range from sandy loam to silt loam the surface mineral 4" Slope is 30% Clay content is =35% in surface mineral 4" Site does not have an argillic horizon with >35% clay

#### Associated sites

|             |                                    |
|-------------|------------------------------------|
| R011XA009ID | Loamy 8-12 PZ ARTRT/PSSPS          |
| R011XA014ID | Sandy 8-14 PZ ARTRT/HECOC8-ACHY    |
| R011XB003ID | Stony Loam 8-12 PZ ARTRW8/PSSPS    |
| R011XB006ID | Loamy 8-12 PZ ARTRT/LECI4          |
| R011XB009ID | Shallow Stony 8-12 PZ ARTRW8/PSSPS |
| R011XB016ID | Sand 8-12 PZ ARTRT-PUTR2/HECOC8    |

|                    |                                         |
|--------------------|-----------------------------------------|
| <b>R011XY001ID</b> | <b>Loamy 8-12 PZ</b>                    |
| <b>R011XY007ID</b> | <b>Gravelly 10-12 PZ</b>                |
| <b>R011XY008ID</b> | <b>South Slope 10-12 PZ</b>             |
| <b>R011XY015ID</b> | <b>Loamy Bottom 8-14 PZ ARTRT/LECI4</b> |

### Similar sites

|                    |                                  |
|--------------------|----------------------------------|
| <b>R011XA009ID</b> | <b>Loamy 8-12 PZ ARTRT/PSSPS</b> |
|--------------------|----------------------------------|

**Table 1. Dominant plant species**

|            |                                                        |
|------------|--------------------------------------------------------|
| Tree       | Not specified                                          |
| Shrub      | (1) <i>Artemisia tridentata</i> ssp. <i>tridentata</i> |
| Herbaceous | (1) <i>Pseudoroegneria spicata</i> ssp. <i>spicata</i> |

### Physiographic features

This site occurs on nearly level to hilly uplands with slopes ranging from 2 to 15 percent with occasional slopes to 25 percent on all aspects. Elevations range from 3000 to 5500 feet (915-1675 meters). Small lava outcrops may occur within and adjacent to the site.

**Table 2. Representative physiographic features**

|                    |                            |
|--------------------|----------------------------|
| Landforms          | (1) Lava plain<br>(2) Hill |
| Flooding frequency | None                       |
| Ponding frequency  | None                       |
| Elevation          | 910 – 1,680 m              |
| Slope              | 0 – 20 %                   |

|        |                                    |
|--------|------------------------------------|
| Aspect | Aspect is not a significant factor |
|--------|------------------------------------|

## Climatic features

The Central Snake River Plain, MLRA 11A, has a mean elevation of 3929 feet above sea level, and varies from a minimum of 2575 feet to a maximum of 8586 feet. The average annual precipitation is 10.03 inches, with a range of 8.30 to 11.46 inches, based on 10 long term climate stations located throughout the MLRA. In general, annual precipitation is below the national average, especially during the summer months. Temperatures show considerable variation during the year. A maximum temperature of 112° Fahrenheit was recorded at the Hagerman climate station (# 103932; elevation 2880 feet), and a minimum of -38° was recorded at the Richfield station. Richfield has also recorded up to 186 days below freezing during the year.

The frost-free period ranges from 116 to 140 days. The freeze-free period can be as short as 144 days to as long as 169 days. Each period is greatest on the west side of the MLRA. In general, morning and afternoon relative humidity is at or far below the national average, especially during the months of May through September.

Table 3 Representative climatic features

|                               |          |
|-------------------------------|----------|
| Frost-free period (average)   | 140 days |
| Freeze-free period (average)  | 170 days |
| Precipitation total (average) | 280 mm   |

## Influencing water features

This site is not influenced by adjacent wetlands, streams or run on.

## Soil features

The soils supporting this site are shallow, well drained, with moderate to moderately rapid permeability. Runoff is very low to high. The erosion hazard is slight to very severe by water and moderate by wind. The available water holding capacity (AWC) is very low. The soils are 10 to 20 inches deep over bedrock. The surface is generally light colored loams. The surface in some areas can be very stony. There may be minor amounts of gravels or stones in the profile but not sufficient amounts to limit production. Some areas may have soils with a clay subsoil at 20 inches limiting effective rooting depth. There may be cracks or fractures in the lava bedrock that extend 60 inches or more into the lava but the soil is limited to the width of the cracks in the lava. These soils are characterized by a xeric or aridic bordering on xeric soil moisture regime. Soil temperature regime is mesic.

Table 4. Representative soil features

|                    |                                                        |
|--------------------|--------------------------------------------------------|
| Surface texture    | (1) Very stony silt loam<br>(2) Sandy loam<br>(3) Loam |
| Drainage class     | Well drained                                           |
| Permeability class | Moderate to moderately rapid                           |
| Soil depth         | 30 – 50 cm                                             |

|                                                          |                |
|----------------------------------------------------------|----------------|
| Surface fragment cover <=3"                              | 0 – 10 %       |
| Surface fragment cover >3"                               | 0 – 30 %       |
| Available water capacity<br>(0-101.6cm)                  | 1.78 – 7.62 cm |
| Sodium adsorption ratio<br>(0-101.6cm)                   | 0 – 10         |
| Soil reaction (1:1 water)<br>(0-101.6cm)                 | 6.1 – 7.8      |
| Subsurface fragment volume <=3"<br>(Depth not specified) | 0 – 10 %       |
| Subsurface fragment volume >3"<br>(Depth not specified)  | 0 – 40 %       |

## Ecological dynamics

### Ecological Dynamics of the Site:

The dominant visual aspect of this site is basin big sagebrush with an understory of bluebunch wheatgrass. Threetip sagebrush may be present. Composition by weight is approximately 40 to 60 percent grasses, 10 to 20 percent forbs and 25 to 40 percent shrubs.

During the last few thousand years, this site has evolved in an arid climate characterized by dry summers and cold, wet winters. Herbivory has historically occurred on this site at low levels of utilization. Herbivores include mule deer, lagomorphs and small rodents. Pronghorn antelope occasionally use the site in the winter.

Fire has historically occurred on the site at intervals of 20-50 years.

The Historic Climax Plant Community (HCPC) moves through many phases depending on the natural and man-made forces that impact the community over time. State 1, described later, indicates some of these phases. The HCPC is Phase A. This plant community is dominated by bluebunch wheatgrass in the understory and basin big sagebrush in the overstory. Subdominant species includes Sandberg bluegrass, Thurber's needlegrass, tapertip hawksbeard, and arrowleaf balsamroot. There is a large variety of other grasses, forbs and shrubs that can occur in minor amounts. Threetip sagebrush can be a subdominant in some plant communities. The plant species composition of Phase A is listed later under "HCPC Plant Species Composition".

Total annual production is 650 pounds per acre (728 kilograms per hectare) in a normal year. Production in a favorable year is 975 pounds per acre (1092 kilograms per hectare). Production in an unfavorable year is 400 pounds per acre (448 kilograms per hectare). Structurally, cool season deep rooted perennial bunchgrasses are very dominant, followed by tall shrubs being more dominant than perennial forbs while shallow rooted perennial bunchgrasses are subdominant.

### FUNCTION:

This site is suited for livestock grazing in the spring, early summer, and fall. The distance to water may be a problem in some areas.

The site provides winter and spring range for mule deer. Pronghorn antelope use the site occasionally in the winter. It is also used by small mammals, raptors, and songbirds. It has some value as brood rearing habitat for sage grouse.

The site has limited value for recreation but does provide some hunting, hiking, off-road vehicles, and photography opportunities.

Due to gentle slopes and relatively low production, this site can be easily degraded from improper livestock management. A mixed stand of shrubs and perennial grasses is necessary to reach the potential of the site.

#### Impacts on the Plant Community.

##### Influence of fire:

In the absence of normal fire frequency, basin big sagebrush can gradually increase on the site. Grasses and forbs decrease as shrubs increase. With the continued absence of fire, basin big sagebrush can displace most of the primary understory species.

When fires become more frequent than historic levels (20-50 years), basin big sagebrush is reduced significantly. Rabbitbrush can increase slightly. With continued short fire frequency, basin big sagebrush can be completely eliminated along with many of the desirable understory species such as bluebunch wheatgrass and Thurber's needlegrass. These species may be replaced by Sandberg bluegrass along with a variety of annual and perennial forbs including noxious and invasive plants. Cheatgrass will invade the site. Bulbous bluegrass may invade at lower elevations. These fine fuels will increase the fire frequency.

Threetip sagebrush may be a component of the plant community in some areas. Threetip sagebrush has been found to be a weak sprouter in some locations and a strong sprouter in others. This suggests the species has ecotypic variation from one geographic location to another. Fire in one location may result in killing most of the plants while in another location, the plant community may become dominated by threetip sagebrush following a fire event. Threetip sagebrush in this site description is considered to be a weak sprouter, but more data is needed.

##### Influence of improper grazing management:

Season-long grazing and/or excessive utilization can be very detrimental to this site. This type of management leads to reduced vigor of the bunchgrasses. With reduced vigor, recruitment of these species declines. As these species decline, the plant community becomes susceptible to increase in basin big sagebrush and noxious and invasive plants. Threetip sagebrush may increase if present in the community.

Continued improper grazing management influences fire frequency by increasing fine fuels. As cheatgrass increases along with other annuals, fires become more frequent.

Proper grazing management that addresses frequency, duration, and intensity of grazing can also keep fine fuels from developing, thereby reducing fire frequency. This can lead to gradual increases in basin big sagebrush. A planned grazing system can be developed to intentionally accumulate fine fuels in preparation for a prescribed burn. Any brush management should be carefully planned, as a reduction in shrubs without a suitable understory of perennial grasses can increase cheatgrass which leads to more frequent fire intervals.

##### Weather influences:

Above normal precipitation in March, April, and May can dramatically increase total annual production of the plant community. These weather patterns can also increase viable seed production of desirable species to provide for recruitment. Likewise, below normal precipitation during these spring months can significantly reduce total annual production and be detrimental to viable seed production. Overall plant composition is normally not affected when perennials have good vigor.

Below normal temperatures in the spring can have an adverse impact on total production regardless of precipitation. An early, hard freeze can occasionally kill some plants.

Prolonged drought adversely affects this plant community in several ways. Vigor, recruitment, and production are usually reduced. Mortality can occur. Prolonged drought can lead to a reduction in fire frequency.

##### Influence of Insects and disease:

Outbreaks can affect vegetation health. Mormon cricket and grasshopper outbreaks occur periodically. Outbreaks seldom cause plant mortality since defoliation of the plant occurs only once during the year of the outbreak.

##### Influence of noxious and invasive plants:

Many of these species add to the fine-fuel component and lead to increased fire frequency.

Perennial and annual invasive species compete with desirable plants for moisture and nutrients. The result is reduced production and change in composition of the understory.

Influence of wildlife:

Big game animals use this site primarily in the spring and winters. Their numbers are seldom high enough to adversely affect the plant community. Burrowing rodents can impact the community providing microsites for invasion of noxious plants.

Watershed:

Decreased infiltration and increased runoff occur with an increase in basin big sagebrush. Desired understory species can be reduced. This composition change can affect nutrient and water cycles. Increased runoff also causes sheet and rill erosion. Abnormally short fire frequency also gives the same results, but to a lesser degree. The long-term effect is a transition to a different state. See the “Hydrology” section for more detail.

Plant Community and Sequence:

Transition pathways between common vegetation states and phases:

State 1.

Phase A to B. Develops with improper grazing management.

Phase A to C. Develops with fire.

Phase B to A. Develops with prescribed grazing.

Phase C to A. Develops with prescribed grazing and no fire.

State 1 Phase C to State 2. Develops through frequent fire or continued improper grazing management. The site has crossed a threshold. It is generally not economically feasible to move this state back to State 1 with accelerated practices.

State 2 to 3. Develops with range seeding.

State 2 to unknown site. Excessive soil loss and changes in the hydrologic cycle caused by continued improper grazing management and/or frequent fire cause this state to cross a threshold and retrogress to a new site with reduced potential. It is generally not economically feasible to move this state back to State 1 with accelerated practices.

Practice Limitations:

Few limitations exist for implementing vegetative management practices on this site although distance to water may be a limiting factor. Slight to moderate limitations exist for implementing facilitating practices on this site. Shallow soil depth may present concerns for livestock watering pipelines, other water developments, and fencing. Moderate limitations exist for implementing accelerating practices. Low average annual precipitation, low available water holding capacity, and the wind erosion hazard on some soils require careful planning and implementation of rangeland seeding, brush management, and prescribed burning. Adequate vegetative cover must be maintained to prevent wind erosion on some soils. Cheatgrass can invade and increase on the site without careful planning and implementation of improvement practices.

State and transition model

Additional community tables

Table 5. Community 1.1 plant community composition

| Group | Common Name | Symbol | Scientific Name | Annual Production () | Foliar Cover (%) |
|-------|-------------|--------|-----------------|----------------------|------------------|
|-------|-------------|--------|-----------------|----------------------|------------------|

Table 6. Community 2.1 plant community composition

| Group | Common Name | Symbol | Scientific Name | Annual Production () | Foliar Cover (%) |
|-------|-------------|--------|-----------------|----------------------|------------------|
|-------|-------------|--------|-----------------|----------------------|------------------|

Table 7. Community 3.1 plant community composition

| Group | Common Name | Symbol | Scientific Name | Annual Production () | Foliar Cover (%) |
|-------|-------------|--------|-----------------|----------------------|------------------|
|-------|-------------|--------|-----------------|----------------------|------------------|

Table 8. Community 4.1 plant community composition

| Group | Common Name | Symbol | Scientific Name | Annual Production () | Foliar Cover (%) |
|-------|-------------|--------|-----------------|----------------------|------------------|
|-------|-------------|--------|-----------------|----------------------|------------------|

Table 9. Community 5.1 plant community composition

| Group | Common Name | Symbol | Scientific Name | Annual Production () | Foliar Cover (%) |
|-------|-------------|--------|-----------------|----------------------|------------------|
|-------|-------------|--------|-----------------|----------------------|------------------|

Table 10. Community 6.1 plant community composition

| Group | Common Name | Symbol | Scientific Name | Annual Production () | Foliar Cover (%) |
|-------|-------------|--------|-----------------|----------------------|------------------|
|-------|-------------|--------|-----------------|----------------------|------------------|

## Animal community

Wildlife Interpretations. Animal Community – Wildlife Interpretations This rangeland ecological site provides diverse habitat for many native wildlife species. Large herbivore use of this ecological site is dominated by mule deer, pronghorn antelope, and elk. Important seasonal habitat is provided for resident and migratory animals including western toad, sagebrush lizard, western rattlesnake, shrews, bats, jackrabbits, ground squirrels, mice, coyote, red fox, badger, sage-grouse, Ferruginous hawk, prairie falcon, horned lark and western meadowlark. Area sensitive species include pygmy rabbit, burrowing owl, great basin ground squirrel and Townsend pocket gopher. Changes in the plant community composition can reduce the number and diversity of wildlife species in the area. With reduced shrub cover shrub obligate avian and mammal species become rare including sage-grouse, brewer's sparrow, sage thrasher, and pygmy rabbits. Encroachment of noxious and invasive plant species (cheatgrass and bulbous bluegrass) can replace native plant species which provide critical feed, brood-rearing and nesting cover for a variety of native wildlife. Water features are sparse provided by seasonal streams, artificial water catchments and springs. This rangeland ecological site is commonly associated with pre-historic lava flows which may provide unique cave habitats for several sensitive animal species, including the Blind Cave Leioidid Beetle, Cave Obligate Mite, Bats and Cave Obligate Harvestman. State 1 Phase 1.1 - Basin Big Sagebrush/ Bluebunch Wheatgrass/ Thurber's Needlegrass/ Sandberg Bluegrass Reference Plant Community (RPC): The RPC provides a diversity of grasses, forbs and shrubs, used by native insect communities who assist in the pollination process for the plant community. The reptile and amphibian community is represented by leopard lizard, short horned lizard, sagebrush lizard, western skink, western rattlesnake, western toad and northern leopard frog. Amphibians are associated with springs and isolated water bodies adjacent to this plant community. Spring developments that capture all available water would preclude the use of these sites by amphibians. Shrub-steppe obligate avian species include the Brewer's sparrow, sage sparrow, sage thrasher and sage-grouse. Brood-rearing habitat for sage-grouse is provided by this plant community. The plant community provides seasonal (winter and spring) food and cover for mule deer, antelope, and elk. A diverse small mammal population including golden-mantled ground squirrels and chipmunks would utilize the site. State 1 Phase 1.2- Basin Big Sagebrush/ Sandberg Bluegrass Plant Community: This plant community is the result of improper grazing management. An increase in canopy cover of sagebrush contributes to a sparse herbaceous understory and can lead to an increase in threetip sagebrush. The reduced herbaceous understory results in lower diversity and numbers of insects. The reptile community is represented by leopard lizard, short horned lizard, sagebrush lizard, western skink and western rattlesnake. You can expect a decrease in populations and diversity of reptiles due to the reduced diversity and canopy cover of herbaceous vegetation and associated invertebrate production. Reduced herbaceous understory is a key factor in limiting the use of this plant community by avian species. Key shrub-steppe obligate avian species include Brewer's sparrow, sage sparrow, sage thrasher and sage-grouse. Brood-rearing habitat for sage-grouse is limited due to the reduced diversity and canopy cover of herbaceous vegetation. This plant community provides a shorter spring forage season for mule deer, antelope, and elk. With an increase in threetip sagebrush the quality of forage for large mammals would be reduced due to the poor palatability and nutritional value of threetip. Winter thermal cover and young of year cover is provided by this phase. A small mammal population including golden-mantled ground squirrels and chipmunks would utilize this plant community. State 1 Phase 1.3 - Bluebunch Wheatgrass/ Sandberg Bluegrass /Bottlebrush Squirreltail Plant Community: This plant community is the result of frequent fire. The plant community, dominated by herbaceous vegetation with little or no sagebrush provides less vertical structure, limiting use by shrub obligate animals. Insect diversity would be reduced but a diverse native forb plant community would still support select pollinators. An increase in rabbitbrush will add fall pollinator habitat. Reptile use would be limited or excluded due to the absence of sagebrush. The dominance of herbaceous vegetation with little sagebrush canopy cover would severely reduce or eliminate use of these areas for nesting by Brewer's sparrow, sage sparrow and sage thrasher. This plant community provides limited brood-rearing habitat for sage grouse when sagebrush cover is nearby. The site would not be suitable for sage-grouse winter habitat.. The dominant herbaceous vegetation improves habitat for grassland avian species (horned lark and western meadowlark). Grassland animal species including savannah sparrow, deer mouse and kangaroo rat would utilize this site for nesting and/or foraging. Mule deer, antelope, and elk forage use would be seasonal (spring through fall) and the site would offer little thermal cover and young of year cover. State 2 - Sandberg Bluegrass / Cheatgrass and Annual Plant Community: This state has developed due to frequent fires or improper grazing management. The reduced forb component in the plant community would support a very limited population of pollinators. Most reptilian species would not be supported with food, water or cover. The plant community would not provide suitable habitat for sage-grouse, sage thrasher, Brewer's sparrow or sage sparrow. Birds of prey including hawks and falcons may range throughout these areas looking for prey species. Large mammals may utilize the herbaceous vegetation in the early part of the year when the invasive annuals (cheatgrass) would be more palatable. At other times of the year large mammals would not regularly utilize these areas due to poor food and cover conditions. The populations of small mammals would be dominated by open grassland species including deer mouse, savannah sparrow, kangaroo rat and Columbian ground squirrel. State 3 - Range Seeding Plant Community: The proposed seeding mixture (native or non-native) would determine the animal species that would utilize the area. A diverse seed mixture of grasses and forbs would provide similar habitat conditions as in the herbaceous plant community described in State 1 phase 1.3. A diverse seed mixture of grasses, forbs and shrubs would provide similar habitat conditions as described in State 1 phase 1.1 or 1.2. A monoculture of non-native grass species would not support diverse populations of insects, reptiles, avians or mammals. Sagebrush obligate animal species would not be supported with a monoculture of grass species. Grassland animal species including western meadowlark, horned lark, savannah sparrow, deer mouse, kangaroo rat and elk would utilize this site for nesting and/or foraging. Birds of prey including hawks and falcons may range throughout these areas looking for prey species.

**Grazing Interpretations:** This site is suited for livestock grazing in the spring, early summer, and fall. The distance to water may be a problem in some areas. Estimated initial stocking rate will be determined with the landowner or decision-maker. They will be based on the inventory which includes species, composition, similarity index, production, past use history, season of use and seasonal preference. Calculations used to determine estimated initial stocking rate will be based on forage preference ratings

## Hydrological functions

The soils on this site are in hydrologic groups B and C. When hydrologic conditions of the vegetative cover is good, the natural erosion hazard is slight to moderate.

## Recreational uses

The site has limited value for recreation but does provide some hunting, hiking, off-road vehicles and photography opportunities.

## Wood products

None

## Other products

None

## Inventory data references

Information presented here has been derived from NRCS clipping and other inventory data. Also, field knowledge of range-trained personnel was used. Those involved in developing this site description include: Dave Franzen, co-owner, Intermountain Rangeland Consultants, LLC Jacy Gibbs, co-owner, Intermountain Rangeland Consultants, LLC Jim Cornwell, Range Management Specialist, IASCD Joe May, State Rangeland Management Specialist, NRCS, Idaho Leah Juarros, Resource Soil Scientist, NRCS, Idaho Lee Brooks, Range Management Specialist, IASCD

## Other references

Hironaka, M., M.A. Fosberg, A. H. Winward. 1983. Sagebrush- Grass Habitat Types of Southern Idaho. University of Idaho. Moscow, Idaho. Bulletin Number 35

USDA Forest Service, Rocky Mountain Research Station. 2004. Restoring Western Ranges and Wildlands. General Technical Report RMRS-GTR-136-vols. 1-3.

USDA, NRCS.2001. The PLANTS Database, Version 3.1 (<http://plants.usda.gov>). National Plant Data Center, Baton Rouge, LA 70874-4490 USA

USDA, Forest Service, Fire Effects Information Database. 2004. [www.fs.fed.us/database/feis](http://www.fs.fed.us/database/feis)

USDI Bureau of Land Management, US Geological Survey; USDA Natural Resources Conservation Service, Agricultural Research Service; Interpreting Indicators of Rangeland Health. Technical Reference 1734-6; version 4-2005.

## Contributors

DLF

## Approval

Kendra Moseley, 4/06/2020

## Rangeland health reference sheet

**Interpreting Indicators of Rangeland Health** is a qualitative assessment protocol used to determine ecosystem condition based on benchmark characteristics described in the Reference Sheet. A suite of 17 (or more) indicators are typically considered in an assessment. The ecological site(s) representative of an assessment location must be known prior to applying the protocol and must be

verified based on soils and climate. Current plant community cannot be used to identify the ecological site.

|                                                |                                                                                                  |
|------------------------------------------------|--------------------------------------------------------------------------------------------------|
| Author(s)/participant(s)                       | Dave Franzen and Jacy Gibbs.                                                                     |
| Contact for lead author                        | Brendan Brazee, State Range Conservationist USDA-NRCS<br>9173 W. Barnes, Suite C Boise, ID 83709 |
| Date                                           | 03/28/2007                                                                                       |
| Approved by                                    |                                                                                                  |
| Approval date                                  |                                                                                                  |
| Composition (Indicators 10 and 12)<br>based on | Annual Production                                                                                |

## Indicators

1. **Number and extent of rills:** Rills: rarely occur on this site.

---

2. **Presence of water flow patterns:** Water-Flow Patterns: rarely occur on this site. When they do occur, they are short, disrupted by cool season perennial grasses and tall shrubs and are not extensive.

---

3. **Number and height of erosional pedestals or terracettes:** Pedestals and/or Terracettes: are rare on this site. In areas where flow patterns and/or rills are present, a few pedestals and terracettes may be expected.

---

4. **Bare ground from Ecological Site Description or other studies (rock, litter, lichen, moss, plant canopy are not bare ground):**  
Bare Ground: no data on bare ground but is expected to range from 30-40 percent.

---

5. **Number of gullies and erosion associated with gullies:** Gullies: do not occur on this site.

---

6. **Extent of wind scoured, blowouts and/or depositional areas:** Wind-Scoured, Blowouts, and/or Deposition Areas: usually not present. The light textured soils in some areas can blow following a fire and evidence may be found in mounded soil deposition in the crowns of bunchgrass.

---

7. Amount of litter movement (describe size and distance expected to travel): Litter Movement: fine litter in the interspaces may move up to 2 feet or further following a significant run-off event. Coarse litter generally does not move.

---

8. Soil surface (top few mm) resistance to erosion (stability values are averages - most sites will show a range of values): Soil Surface Resistance to Erosion: values should range from 4 to 6 but needs to be tested.

---

9. Soil surface structure and SOM content (include type of structure and A-horizon color and thickness): Soil Surface Loss or Degradation: The A or A1 horizon is typically 1 to 4 inches thick. Structure ranges from weak very fine or fine granular. Soil organic matter (SOM) is 0.5 to 3 percent.

---

10. Effect of community phase composition (relative proportion of different functional groups) and spatial distribution on infiltration and runoff: Plant Community Composition and Distribution Relative to Infiltration: bunchgrasses, especially deep-rooted perennials, slow run-off and increase infiltration. Shrubs accumulate snow in the interspaces. Terracettes provide a favorable micro-site for vegetation establishment, which further increases infiltration.

---

11. Presence and thickness of compaction layer (usually none; describe soil profile features which may be mistaken for compaction on this site): Compaction Layer: not present.

---

12. Functional/Structural Groups (list in order of descending dominance by above-ground annual-production or live foliar cover using symbols: >>, >, = to indicate much greater than, greater than, and equal to):

Dominant: Functional/Structural Groups: cool season deep-rooted perennial bunchgrasses &>>tall shrubs&> perennial forbs&> shallow rooted grasses.

Sub-dominant:

Other:

Additional:

---

13. Amount of plant mortality and decadence (include which functional groups are expected to show mortality or decadence): Plant Mortality/Dcadence: basin big sagebrush will become decadent in the absence of fire and ungulate grazing. Grass and forb mortality will occur as tall shrubs increase.

---

14. Average percent litter cover (%) and depth ( in): Litter Amount: annual litter cover in the interspaces will be 5-10 percent to a depth of

---

15. Expected annual annual-production (this is TOTAL above-ground annual-production, not just forage annual-production): Annual Production: is 650 lbs. per acre in a year with normal precipitation and temperatures. Perennial grasses produce 40-60 percent of the total, forbs 10-20 percent, and shrubs 25-40 percent.

---

16. Potential invasive (including noxious) species (native and non-native). List species which BOTH characterize degraded states and have the potential to become a dominant or co-dominant species on the ecological site if their future establishment and growth is not actively controlled by management interventions. Species that become dominant for only one to several years (e.g., short-term response to drought or wildfire) are not invasive plants. Note that unlike other indicators, we are describing what is NOT expected in the reference state for the ecological site: Invasive Plants: cheatgrass, clasping pepperweed, beggars ticks, tansymustard, Jim Hill tumblemustard, yellow salsify, burr buttercup, medusahead, Russian thistle, annual kochia, and halogeton. Bulbous bluegrass can invade at lower elevations.
- 

17. Perennial plant reproductive capability: Reproductive Capability of Perennial Plants: all functional groups have the potential to reproduce in normal years.
-