

Ecological site R013XY054ID

Marsh

TYLA-SCAC3

Last updated: 9/23/2020
Accessed: 09/08/2026

General information

Provisional. A provisional ecological site description has undergone quality control and quality assurance review. It contains a working state and transition model and enough information to identify the ecological site.

MLRA notes

Major Land Resource Area (MLRA): 013X–Eastern Idaho Plateaus

Land Resource Region: B (Northwestern Wheat and Range) MLRA: 13 (Eastern Idaho Plateaus) EPA EcoRegion: Level III (Middle Rockies)

LRU notes

013X-Eastern Idaho Plateaus Additional moisture site. <https://soils.usda.gov/survey/geography/mlra/index.html>

Classification relationships

No data.

Ecological site concept

Site receives additional water. Soils are: Not saline or saline-sodic. Deep to very deep, not skeletal within 20" of soil surface. Season water table 40" with high amounts of organic matter Poorly drained, standing water is common until late summer Not strongly or violently effervescent in surface mineral 10". Slope is 30%. Clay content is = 35% in surface mineral 4". Site does not have an argillic horizon with > 35% clay.

Associated sites

R013XY038ID	Meadow DECA18-CANE2
R013XY039ID	Dry Meadow PONE-PHAL2
R013XY045ID	Loamy Bottom 12-16 PZ ARTRT/LECI4-ELLAL
R013XY050ID	Riparian Wet Meadow SALIX/CAREX
R013XY053ID	Wet Meadow CAREX-JUNCUS

Table 1. Dominant plant species

Tree	Not specified
Shrub	Not specified
Herbaceous	Not specified

Physiographic features

This site generally occurs on nearly level margins of streams, ponds, lakes, springs and reservoirs, as stringers paralleling stream and river channels. This site can also occur as parts of old stream courses, oxbows and potholes. Slopes are less than 2 percent. Elevation ranges between 4,000 to 6,800 feet (1200 to 2100 meters).

Table 2. Representative physiographic features

Landforms	(1) Lake terrace (2) Lakebed (3) Flood plain
Flooding duration	Long (7 to 30 days) to very long (more than 30 days)
Flooding frequency	Occasional to frequent
Ponding duration	Brief (2 to 7 days) to long (7 to 30 days)
Ponding frequency	None to frequent
Elevation	1,220 – 2,070 m
Slope	0 %
Ponding depth	10 – 30 cm
Water table depth	0 cm
Aspect	Aspect is not a significant factor

Climatic features

Influencing water features

The Marsh site is influenced by additional water from either adjacent streams through seasonal flooding, water table, seeps or springs or from run-on from adjacent sites. The site may include the following wetlands and stream types.

Soil features

Typically these soils are erratically stratified with a surface mat of roots 4 or more inches deep over 3 to 6 inches of organic material over clays, clay loams, or silty clay loams that are over 60 inches deep. These soils are poorly drained and alluvial in origin. The soils are usually slightly acid to slightly alkaline in pH. Soils are saturated throughout the growing season and usually have standing water at or above the surface into the late summer.

Flooding can be frequent during snowmelt and just after snowmelt. Ponding is normal. The plant community is dependent on saturated soils and standing water during the growing season.

Soil Series Correlated to this Ecological Site -

Bear Lake
Bloomington
Dingle
Dinswamp
Marsh

Table 3. Representative soil features

Surface texture	(1) Silty clay loam (2) Silt loam
Drainage class	Somewhat poorly drained to poorly drained
Permeability class	Moderately slow to moderate
Soil depth	150 cm
Surface fragment cover <=3"	Not specified
Surface fragment cover >3"	Not specified
Available water capacity (0-101.6cm)	18.03 – 28.7 cm
Calcium carbonate equivalent (0-101.6cm)	Not specified
Electrical conductivity (0-101.6cm)	Not specified

Sodium adsorption ratio (0-101.6cm)	Not specified
Soil reaction (1:1 water) (0-101.6cm)	6.6 – 10
Subsurface fragment volume <=3" (Depth not specified)	Not specified
Subsurface fragment volume >3" (Depth not specified)	Not specified

Ecological dynamics

The dominant visual aspect of this site is broadleaf cattail and/or hardstem bulrush. Hardstem bulrush is more dominant where water is 6 to 8 inches deep or deeper throughout the growing season. Broadleaf cattail is often more dominant where the water is at the soil surface through most of the growing season and may dry out at the surface later in the summer. When the two species occur together, broadleaf cattail is in shallower water than the hardstem bulrush. This is common on margins of ponds, lakes, and reservoirs.

The site often occurs within a complex of wetland sites when adjacent to streams or rivers. In these situations, the soil surface of the area where the site is found is typically slightly undulating causing small depressions and high spots with variable soil moisture regimes. The plant communities found on these areas are sites within the complex. Characteristics of these sites are as follows:

1. Marsh site. Deeper depressions with the water table at or near the surface or slightly above the surface for the entire growing season. This site is dominated by broadleaf cattail, hardstem bulrush, and common threesquare.
2. Wet Meadow site. Shallow to depression areas with the water table at or near the surface for the entire growing season. This site is dominated by *Carex* spp. and *Junus* spp.
3. Meadow site. Slightly higher areas that are drier during the growing season. The water table is down to a depth of 20 to 40 inches by the end of the growing season. This site is dominated by tufted hairgrass and Nebraska sedge.
4. Dry Meadow site. Even higher areas that are even drier during the growing season and the water table is > 40 inches deep by the end of the growing season. The plant community is dominated by Nevada bluegrass and alpine timothy.

Either hardstem bulrush or broadleaf cattail can form impenetrable monotypic stands. Composition by weight is 95 percent grass or grass-like and 5 percent forbs.

During the last few thousand years, this site has evolved in a semi-arid and montane climate characterized by dry summers and cold, moist or wet winters. The site has evolved on deep alluvial soils that are saturated at the soil surface or standing water throughout most of the growing season. Herbivory has historically occurred on this site at very low levels of utilization. Herbivory by some small mammals such as muskrats does occur.

Fire has had little influence on the development of the site. Rare wildfires can occur following consecutive drought years, particularly broadleaf cattail stands that go dry late in the summer.

The conditions for the plant community of this site are variable due to differences in the duration and depth of standing water. Dominant species are mostly rhizomatous and can form monotypic stands.

An infinite number of combinations of factors that influence the ecology of potential plant communities exist. For practical purposes, four plant communities where the depth to the water table drives the vegetative composition have been described. They are:

- Dry meadow Water table at >40" at end of growing season
- Meadow Water table at 20-40" at end of growing season
- Wet meadow Water table at 10-20" at end of growing season

- Marsh Water at surface to

State and transition model

Additional community tables

Table 4. Community 1.1 plant community composition

Group	Common Name	Symbol	Scientific Name	Annual Production ()	Foliar Cover (%)
-------	-------------	--------	-----------------	----------------------	------------------

Table 5. Community 2.1 plant community composition

Group	Common Name	Symbol	Scientific Name	Annual Production ()	Foliar Cover (%)
-------	-------------	--------	-----------------	----------------------	------------------

Animal community

Wildlife Interpretations. Animal Community – Wildlife Interpretations This ecological site provides habitat for unique wetland wildlife species. The duration of hydrology results in abundant herbaceous cover and invertebrate production. The insects provide an extremely important prey base for shorebirds and waterfowl. Wetland dependant animal species utilizing the site include marsh wren, great blue heron, ibis, teal, mallard, yellow-head blackbird, red-winged blackbird, northern leopard frog, western toad, muskrat, and beaver. Large herbivores infrequently use the site due to the extreme hydrologic conditions and poor nutritional value of the forage. Loss of site hydrology significantly reduces habitat value of the adjacent ecological sites. Open water is being provided by runoff, ponding, flooding, seasonal high water table, and natural springs. State 1 Phase 1.1 – Broadleaf Cattail/ Hardstem Bulrush Reference Plant Community (RPC): The RPC and associated hydrology with deep ponded water, provide a diverse population of invertebrates (benthic and terrestrial) unique to wetlands and beneficial to water loving birds. Arthropods such as dragonflies, damselflies, midges, backswimmers, scuds, and waterfleas make up a large portion of potential invertebrates in these wetlands. They are a basic food source for many higher trophic level organisms. The reptile and amphibian community is represented by western skink, rubber boa, western rattlesnake, western toad, boreal chorus frog, and northern leopard frog. A diverse amphibian population is a key indicator of good ecological health on this site. Loss of hydrology will limit or exclude amphibians from this ecological site. The RPC may provide nesting, rearing, roosting, and forage habitat for bird species such as Canada goose, American wigeon, mallard, pied billed grebe, great blue heron, sora, whimbrel, Wilson's snipe, marsh wren, and long-billed curlew. Small mammal populations are extremely limited due to the hydrologic conditions. Deer mouse, montane vole, and western jumping mouse would utilize the site for water throughout the year or during the winter when water is frozen. State 2 Carex sp./ Juncus Sp./ Forbs/ Grasses Plant Community: The animal community may be similar to Wet Meadow, Meadow, Dry Meadow, or upland sites in low to mid-seral status depending on the severity of hydrology manipulation from State 1 Phase 1.1. The management of the resulting plant communities will be a major factor in determining the diversity and populations of the animal community. Refer to the above mentioned ecological sites for MLRA 13 for descriptions of the potential wildlife community. Grazing Interpretations. Due to standing water and low value forage, grazing during the growing season is rare. Grazing may occur during the dormant period. If dry soil conditions occur into the spring and livestock remove the tops of either broadleaf cattail or hardstem bulrush, the plants can die if water then covers them. Care must be exercised in grazing adjacent sites to avoid downcutting of stream channels that can influence standing water in this site. The potential of the site can be lost. Estimated initial stocking rate will be determined with the landowner or decision-maker. They will be based on the inventory, past use history and type, condition of vegetation, production, season of use, and seasonal preference. Calculations used to determine estimated initial stocking rate will be based on forage preference ratings.

Hydrological functions

No data.

Recreational uses

This site presents an aesthetically pleasing view of lush vegetation consisting primarily of grass-like plants and standing water. Hikers and fisherman often traverse the edges of this site. Fishing opportunities often exist on open water adjacent to this site. Waterfowl hunting can occur on or adjacent to the site.

Wood products

None.

Other products

None.

Other information

Field Offices American Falls, ID Blackfoot, ID Burley, ID Driggs, ID Ft. Hall, ID Idaho Falls, ID Malad, ID Pocatello, ID Rexburg, ID Soda Springs, ID St. Anthony, ID

Inventory data references

Information presented here has been derived from NRCS clipping and other inventory data. Also, field knowledge of range-trained personnel was used. Those involved in developing this site description include: Dave Franzen, co-owner, Intermountain Rangeland Consultants, LLC Jacy Gibbs, co-owner, Intermountain Rangeland Consultants, LLC Jim Cornwell, Range Management Specialist, IASCD Dan Ogle, Plant Materials Specialist, NRCS, Idaho Chris Hoag, Wetland Plant Ecologist, NRCS, Idaho Brendan Brazee, State Rangeland Management Specialist, NRCS, Idaho Kristen May, Resource Soil Scientist, NRCS, Idaho Lee Brooks, Range Management Specialist, IASCD

Other references

USDA, NRCS.2001. The PLANTS Database, Version 3.1 (<http://plants.usda.gov>.) National Plant Data Center, Baton Rouge, LA 70874-4490 USA.

USDA NRCS. 1992. Major Land Resource Area, Owyhee High Plateau, Range Site Descriptions. Reno, Nevada.

USDA NRCS. Major Land Resource Area, Owyhee High Plateau, Range Site Descriptions. Portland, Oregon.

USDA, Forest Service. 2004. (www.fs.fed.us/database/feis/plants).

USDI, BLM and Idaho Conservation Data Center. 2001. Riparian and Wetland Plant Associations of Southwestern Idaho.

USDI Bureau of Land Management, US Geological Survey; USDA Natural Resources Conservation Service, Agricultural Research Service; Interpreting Indicators of Rangeland Health. Technical Reference 1734-6; Version 4-2005.

Approval

Kendra Moseley, 9/23/2020

Rangeland health reference sheet

Interpreting Indicators of Rangeland Health is a qualitative assessment protocol used to determine ecosystem condition based on benchmark characteristics described in the Reference Sheet. A suite of 17 (or more) indicators are typically considered in an assessment. The ecological site(s) representative of an assessment location must be known prior to applying the protocol and must be verified based on soils and climate. Current plant community cannot be used to identify the ecological site.

Author(s)/participant(s)	Dave Franzen and Jacy Gibbs Intermountain Range Consultants 17700 Fargo Rd. Wilder, ID 83676
Contact for lead author	Brendan Brazee, State Rangeland Management Specialist USDA-NRCS 9173 W. Barnes Drive, Suite C, Boise, ID 83709
Date	06/24/2009
Approved by	
Approval date	

Composition (Indicators 10 and 12) based on	Annual Production
---	-------------------

Indicators

1. Number and extent of rills: rills do not occur on this site.
2. Presence of water flow patterns: water-flow patterns do not occur.
3. Number and height of erosional pedestals or terracettes: neither occurs on this site.
4. Bare ground from Ecological Site Description or other studies (rock, litter, lichen, moss, plant canopy are not bare ground): none.
5. Number of gullies and erosion associated with gullies: none.
6. Extent of wind scoured, blowouts and/or depositional areas: blowouts and depositional areas do not occur.
7. Amount of litter movement (describe size and distance expected to travel): litter may move in the standing water.
8. Soil surface (top few mm) resistance to erosion (stability values are averages - most sites will show a range of values): values should range from 4 to 6 but needs to be tested.
9. Soil surface structure and SOM content (include type of structure and A-horizon color and thickness): Structure ranges weak and moderate fine and medium granular to weak fine subangular blocky. Soil organic matter (SOM) ranges from 10 to 80 percent. Surface color is usually black to very dark brown to very dark gray. The A or A1 horizon is typically 2 to 11 inches thick.
10. Effect of community phase composition (relative proportion of different functional groups) and spatial distribution on infiltration and runoff: site has standing water.
11. Presence and thickness of compaction layer (usually none; describe soil profile features which may be mistaken for compaction on this site): is not present.

12. **Functional/Structural Groups** (list in order of descending dominance by above-ground annual-production or live foliar cover using symbols: >>, >, = to indicate much greater than, greater than, and equal to):

Dominant: rhizomatous species

Sub-dominant: perennial forbs

Other:

Additional:

13. **Amount of plant mortality and decadence** (include which functional groups are expected to show mortality or decadence): little mortality occurs on the site.
-

14. **Average percent litter cover (%) and depth (in):** not applicable.
-

15. **Expected annual annual-production** (this is TOTAL above-ground annual-production, not just forage annual-production): is 4500 pounds per acre (5040 Kg/ha) in a year with normal amounts of standing water and temperatures. Rhizomatous species produce 90-95 percent of the total production and forbs less than 10 percent.
-

16. **Potential invasive (including noxious) species** (native and non-native). List species which BOTH characterize degraded states and have the potential to become a dominant or co-dominant species on the ecological site if their future establishment and growth is not actively controlled by management interventions. Species that become dominant for only one to several years (e.g., short-term response to drought or wildfire) are not invasive plants. Note that unlike other indicators, we are describing what is NOT expected in the reference state for the ecological site: includes purple loosestrife, common reed.
-

17. **Perennial plant reproductive capability:** all functional groups have the potential to reproduce in most years. Most of the plants can reproduce vegetatively.
-