

Ecological site R019XI102CA  
Shallow uplands  
13-24" p.z.

Accessed: 09/24/2026

General information

**Provisional.** A provisional ecological site description has undergone quality control and quality assurance review. It contains a working state and transition model and enough information to identify the ecological site.

Table 1. Dominant plant species

|            |                                                                                    |
|------------|------------------------------------------------------------------------------------|
| Tree       | Not specified                                                                      |
| Shrub      | (1) <i>Eriogonum arborescens</i><br>(2) <i>Eriogonum grande</i> var. <i>grande</i> |
| Herbaceous | (1) <i>Nassella</i>                                                                |

Physiographic features

This ecological site is found on the backslopes of hills and mountains on slopes ranging from 15 to 75 percent. The elevations range from 49 to 2440 feet, and it is found on all aspects.

Table 2. Representative physiographic features

|                    |                                    |
|--------------------|------------------------------------|
| Landforms          | (1) Hill<br>(2) Mountain           |
| Flooding frequency | None                               |
| Ponding frequency  | None                               |
| Elevation          | 20 – 740 m                         |
| Slope              | 20 – 80 %                          |
| Aspect             | Aspect is not a significant factor |

Climatic features

This ecological site is found only on Santa Cruz Island, and due to its size, the temperature and precipitation ranges have been grouped together to capture the entire island's variance.

The average annual precipitation is 19 inches with a range between 13 and 24 inches, mostly in the form of rain in the winter months (November through April). The average annual air temperature is approximately 56 to 73 degrees Fahrenheit, and the frost-free (>32F) season is 320 to 365 days.

**Table 3 Representative climatic features**

|                               |          |
|-------------------------------|----------|
| Frost-free period (average)   | 370 days |
| Freeze-free period (average)  | 370 days |
| Precipitation total (average) | 610 mm   |

### Influencing water features

This site is not influenced by wetland or riparian water features.

### Soil features

These soils formed in residuum from schist (Delphine) or in residuum from volcanic breccia, andesite, or basalt (Spinnaker). They are shallow to bedrock, well-drained or somewhat excessively drained, with loamy surface and subsurface textures. Mean annual soil temperatures (MAST) range from 59 and 71 degrees F, and are classified as thermic.

This ecological site is found in the following map units on these soil components:

#### SSA Map symbol Component

CA688 101 Spinnaker  
CA688 240 Delphine  
CA688 241 Delphine  
CA688 250 Spinnaker  
CA688 251 Spinnaker  
CA688 260 Spinnaker  
CA688 270 Spinnaker  
CA688 271 Spinnaker  
CA688 291 Spinnaker

**Table 4. Representative soil features**

|                      |                                                                     |
|----------------------|---------------------------------------------------------------------|
| Surface texture      | (1) Gravelly<br><br>(2) Extremely gravelly<br><br>(3) Very gravelly |
| Family particle size | (1) Loamy                                                           |
| Drainage class       | Well drained to somewhat excessively drained                        |
| Permeability class   | Moderately slow to moderately rapid                                 |
| Soil depth           | 20 – 50 cm                                                          |

|                                                          |                |
|----------------------------------------------------------|----------------|
| Surface fragment cover <=3"                              | 60 %           |
| Available water capacity<br>(0-101.6cm)                  | 1.52 – 4.32 cm |
| Electrical conductivity<br>(0-101.6cm)                   | Not specified  |
| Sodium adsorption ratio<br>(0-101.6cm)                   | Not specified  |
| Soil reaction (1:1 water)<br>(0-101.6cm)                 | 5.6 – 7.3      |
| Subsurface fragment volume <=3"<br>(Depth not specified) | 40 – 70 %      |

### Ecological dynamics

This ecological site is found primarily on shallow soils on the interior ridge tops on very steep slopes and rock outcrops. The shallow rocky soils of this site have low water holding capacities that will generally inhibit germination and plant growth, which determines the species composition of this plant community and maintains the site as a sparsely vegetated, rocky habitat.

The historical state is sparsely vegetated with common plants including Santa Cruz Island buckwheat (*Eriogonum arborescens*), redflower buckwheat (*Eriogonum grande* var. *grande*), dove weed (*Croton setigerus*), island broom (*Lotus dendroideus*), threeawns (*Aristida* spp.) and needlegrasses (*Nassella* spp.). The reference state is similar to the historical state, except the herbaceous cover is dominated by non-native annual species such as oats (*Avena* spp.) and bromes (*Bromus* spp.). In the reference state, the native grasses have a scattered cover due to the invasiveness of the non-native species.

The natural successional cycle for this ecological site is driven primarily by patchy disturbances that may affect small portions of the site but rarely the entire site. Sites that are on these exposed, rocky ridges may experience rock or land slides that could potentially disturb the vegetation, opening the site for new buckwheats and grasses to establish. Several years of drought could also affect the site, by causing many of the plants to die over large areas. Once the drought ends, the species would begin to reestablish from the local seedbank.

The invasion of non-native annuals into this site can also shift the successional patterns by bringing fire disturbance into the system. Fire was not likely an issue in the natural system, given the sparse nature of the plants on rocky, exposed ground. Larger patches may have burned, but likely would not have spread beyond that patch. The non-native annuals that have invaded the system provide a continuous fuel source for fires to spread throughout the site, burning out large expanses of the native vegetation. Frequent fires can completely remove many of the native species from the site by removing seed sources. After a fire, the non-native annuals will be extremely competitive during germination and establishment, and will outcompete many of the natives that attempt to reestablish.

Overgrazing by domestic livestock and non-native wildlife can also disrupt the natural disturbance regime of this site. Overgrazing causes chronic, severe defoliation of the buckwheats and native grasses and could facilitate the invasion by non-native annual grasses and forbs into the site.

### State and transition model

Figure 3. State Transition Model

Additional community tables

Table 5. Community 1.1 plant community composition

| Group           | Common Name                 | Symbol | Scientific Name                     | Annual Production () | Foliar Cover (%) |
|-----------------|-----------------------------|--------|-------------------------------------|----------------------|------------------|
| Shrub/Vine      |                             |        |                                     |                      |                  |
| 1               | shrubs                      |        |                                     | 39-1233              |                  |
|                 | redflower buckwheat         | ERGRG5 | <i>Eriogonum grande var. grande</i> | 1-673                | –                |
|                 | Santa Cruz Island buckwheat | ERAR6  | <i>Eriogonum arborescens</i>        | 34-560               | –                |
|                 | dove weed                   | CRSE11 | <i>Croton setigerus</i>             | 0-22                 | –                |
|                 | island broom                | LODE2  | <i>Lotus dendroideus</i>            | 1-11                 | –                |
| Grass/Grasslike |                             |        |                                     |                      |                  |
| 2               | grasses                     |        |                                     | 34-224               |                  |
|                 | threeawn                    | ARIST  | <i>Aristida</i>                     | 0-112                | –                |
|                 | oat                         | AVENA  | <i>Avena</i>                        | 11-112               | –                |
|                 | needlegrass                 | NASSE  | <i>Nassella</i>                     | 11-78                | –                |
|                 | red brome                   | BRRU2  | <i>Bromus rubens</i>                | 11-39                | –                |
|                 | annual fescue               | VUMY   | <i>Vulpia myuros</i>                | 0-11                 | –                |
|                 | soft brome                  | BRHO2  | <i>Bromus hordeaceus</i>            | 0-11                 | –                |
| Forb            |                             |        |                                     |                      |                  |
| 3               |                             |        |                                     | 1-84                 |                  |
|                 | island bristleweed          | HADE4  | <i>Hazardia detonsa</i>             | 0-56                 | –                |
|                 | pricklypear                 | OPUNT  | <i>Opuntia</i>                      | 0-17                 | –                |
|                 | spikemoss                   | SELAG  | <i>Selaginella</i>                  | 0-6                  | –                |
|                 | cudweed                     | GNAPH  | <i>Gnaphalium</i>                   | 0-6                  | –                |
|                 | longbeak stork's bill       | ERBO   | <i>Erodium botrys</i>               | 0-2                  | –                |
|                 | smooth cat's ear            | HYGL2  | <i>Hypochaeris glabra</i>           | 0-1                  | –                |

Table 6. Community 2.1 plant community composition

| Group | Common Name | Symbol | Scientific Name | Annual Production () | Foliar Cover (%) |
|-------|-------------|--------|-----------------|----------------------|------------------|
|-------|-------------|--------|-----------------|----------------------|------------------|

Table 7. Community 3.1 plant community composition

| Group | Common Name | Symbol | Scientific Name | Annual Production () | Foliar Cover (%) |
|-------|-------------|--------|-----------------|----------------------|------------------|
|-------|-------------|--------|-----------------|----------------------|------------------|

Animal community

This ecological site has steeper slopes and bare ground and so is not utilized by many animal communities; however small rodents and birds may use it for cover and food.

Hydrological functions

N/A

Recreational uses

N/A

Wood products

N/A

Other products

N/A

## Other information

N/A

## Inventory data references

The following NRCS plots were used to describe this ecological plot. SC-355A lbs SCV-102 lbs SCV-103 % SC-379 % SCV-6 lbs- Site location

## Type locality

|                                      |                                                                                                                                |
|--------------------------------------|--------------------------------------------------------------------------------------------------------------------------------|
| Location 1: Santa Barbara County, CA |                                                                                                                                |
| UTM zone                             | N                                                                                                                              |
| UTM northing                         | 3765595                                                                                                                        |
| UTM easting                          | 0244469                                                                                                                        |
| General legal description            | The site location is on Santa Cruz Island, about 1000' from the South Ridge Road, on the east side of the Willows Canyon Road. |

## Other references

Junak, Steve; Ayers, Tina; Scott, Randy; Wilken, Dieter; and Young, David (1995). A Flora of Santa Cruz Island. Santa Barbara Botanic Garden, Santa Barbara, CA.

## Contributors

Munnecke

## Rangeland health reference sheet

**Interpreting Indicators of Rangeland Health** is a qualitative assessment protocol used to determine ecosystem condition based on benchmark characteristics described in the Reference Sheet. A suite of 17 (or more) indicators are typically considered in an assessment. The ecological site(s) representative of an assessment location must be known prior to applying the protocol and must be verified based on soils and climate. Current plant community cannot be used to identify the ecological site.

|                          |  |
|--------------------------|--|
| Author(s)/participant(s) |  |
| Contact for lead author  |  |
| Date                     |  |

|                                             |                   |
|---------------------------------------------|-------------------|
| Approved by                                 |                   |
| Approval date                               |                   |
| Composition (Indicators 10 and 12) based on | Annual Production |

## Indicators

### 1. Number and extent of rills:

---

### 2. Presence of water flow patterns:

---

### 3. Number and height of erosional pedestals or terracettes:

---

### 4. Bare ground from Ecological Site Description or other studies (rock, litter, lichen, moss, plant canopy are not bare ground):

---

### 5. Number of gullies and erosion associated with gullies:

---

### 6. Extent of wind scoured, blowouts and/or depositional areas:

---

### 7. Amount of litter movement (describe size and distance expected to travel):

---

### 8. Soil surface (top few mm) resistance to erosion (stability values are averages - most sites will show a range of values):

---

### 9. Soil surface structure and SOM content (include type of structure and A-horizon color and thickness):

---

### 10. Effect of community phase composition (relative proportion of different functional groups) and spatial distribution on infiltration and runoff:

---

### 11. Presence and thickness of compaction layer (usually none; describe soil profile features which may be mistaken for compaction on this site):

---

12. Functional/Structural Groups (list in order of descending dominance by above-ground annual-production or live foliar cover using symbols: >>, >, = to indicate much greater than, greater than, and equal to):

Dominant:

Sub-dominant:

Other:

Additional:

---

13. Amount of plant mortality and decadence (include which functional groups are expected to show mortality or decadence):

---

14. Average percent litter cover (%) and depth ( in):

---

15. Expected annual annual-production (this is TOTAL above-ground annual-production, not just forage annual-production):

---

16. Potential invasive (including noxious) species (native and non-native). List species which BOTH characterize degraded states and have the potential to become a dominant or co-dominant species on the ecological site if their future establishment and growth is not actively controlled by management interventions. Species that become dominant for only one to several years (e.g., short-term response to drought or wildfire) are not invasive plants. Note that unlike other indicators, we are describing what is NOT expected in the reference state for the ecological site:

---

17. Perennial plant reproductive capability:

---