

Ecological site F022BI125CA

Cold Frigid Tephra Over Outwash Plains Or Lake Terraces

Accessed: 10/03/2026

General information

Provisional. A provisional ecological site description has undergone quality control and quality assurance review. It contains a working state and transition model and enough information to identify the ecological site.

MLRA notes

Major Land Resource Area (MLRA): 022B–Southern Cascade Mountains

Site concept: Landform: (1) Outwash plain, (2) Lake terrace Elevation (feet): 5850-6360 Slope (percent): 0-15 Water Table Depth (inches): 19 to greater than 60 Flooding-Frequency: None Ponding-Frequency: None Aspect: Non-influencing Mean annual precipitation (inches): 23.0-49.0 Primary precipitation: Winter months in the form of snow Mean annual temperature: 42 and 44 degrees F (5.5 and 6.6 degrees C) Restrictive Layer: Silica-cemented duripan occurs at depths between 20 to 60 inches Temperature Regime: Frigid Moisture Regime: Xeric Parent Materials: Mixed tephra and outwash over outwash from volcanic rocks Surface Texture: (1) Very bouldery medial loamy coarse sand, (2) Ashy coarse sand Surface Fragments =3" (% Cover): 5-70 Surface Fragments > 3" (% Cover): 0-20 Soil Depth (inches): 20-60 Vegetation: Although the heavy dominance of lodgepole pine here is partly in response to high fire frequency, the root restrictive layer and cold air drainage may exclude other conifers from establishing on this site. Notes: This ecological site occurs on glacial outwash plains and lake terraces.

Classification relationships

Forest Alliance = *Pinus contorta* ssp. *murrayana* – Lodgepole pine forest; Association = (no matching species). (Sawyer, John O., Keeler-Wolf, Todd, and Evens, Julie M. 2009. A Manual of California Vegetation. 2nd ed. California Native Plant Society Press. Sacramento, California.)

Associated sites

F022BI108CA	Frigid Moist Sandy Lake Or Stream Terraces This is a wet lodgepole site found on lake and stream terraces.
F022BI126CA	Cold Frigid Tephra Over Moraine Slopes This site is found on the moraines interspersed around the outwash and has more Jeffrey pine.

Similar sites

F022BI105CA	Frigid Sandy Loam Debris Flow On Stream Terraces This is a Sierra lodgepole pine-quaking aspen site.
F022BI125CA	Cold Frigid Tephra Over Outwash Plains Or Lake Terraces This is a white fir-Sierra lodgepole pine site found in wetter conditions.

F022BI123CA	Frigid Flat Outwash Terraces This is a white fir-Sierra lodgepole pine site with some aspen.
F022BI117CA	Frigid Coarse Glaciolacustrine Gentle Slopes This is a California red fir-Sierra lodgepole pine site found at higher elevations.

Table 1. Dominant plant species

Tree	(1) <i>Pinus contorta</i> var. <i>murrayana</i>
Shrub	Not specified
Herbaceous	(1) <i>Elymus elymoides</i>

Physiographic features

This ecological site occurs on glacial outwash plains and lake terraces at 5,850 to 6,360 feet in elevation. Slopes range from 0 to 15 percent.

This site has a seasonal water table associated with the Humic Haploxerands, moist lake terrace component. The water table fluctuates from 19 to 30 inches during the wetter months then drops to below 60 inches from July to November.

Table 2. Representative physiographic features

Landforms	(1) Outwash plain (2) Lake terrace
Flooding frequency	None
Ponding frequency	None
Elevation	1,780 – 1,940 m
Slope	0 – 20 %
Water table depth	50 cm

Climatic features

This ecological site receives most of its annual precipitation in the winter months in the form of snow. The mean annual precipitation ranges from 23 and 49 inches (594 mm to 1,245 mm) and the mean annual temperature is between 42 and 44 degrees F (5.5 and 6.6 degrees C). The frost free (>32F) season is 60 to 85 days. The freeze free (>28F) season is 75 to 190 days.

There are no representative climate stations for this site. The nearest one is Manzanita Lake.

Table 3 Representative climatic features

Frost-free period (average)	90 days
Freeze-free period (average)	190 days
Precipitation total (average)	1,250 mm

Influencing water features

This ecological site is not influenced by wetland or riparian water features.

Soil features

This site is associated with the Badgerwash and Humic Haploxerands, moist lake terrace soil components.

The Badgerwash soils are moderately deep, well-drained soils that formed in mixed tephra and outwash over outwash from volcanic rocks. The surface texture is very bouldery medial loamy coarse sand, with extremely stony medial sandy loam textures in the lower horizons. There are greater than 35 percent rock fragments throughout the soil depth and a silica-cemented duripan is encountered at 20 to 40 inches. The Badgerwash soils are classified as Ashy-skeletal, mixed, frigid Humic Vitrixerands.

Humic Haploxerands, moist lake terrace soils are moderately deep to deep, moderately well-drained soils that formed in tephra over glaciolacustrine deposits from volcanic rocks. The tephra deposits are blackish in color and are approximately 7 inches deep. The tephra has developed an A/C horizon sequence over the buried soil. The A horizon has an ashy coarse sand, with a gravelly ashy coarse sand texture just below. The buried horizon has a very stony medial sandy loam texture over a very stony medial fine sandy loam. A silica-cemented duripan occurs at depths between 20 to 60 inches.

These soils have very low AWC (available water capacity) in the upper 60 inches of soil. Permeability is moderately rapid to rapid in the upper horizon, and slow to very slow through the duripan.

This ecological site is associated with the following soil components within the Lassen Volcanic National Park Soil Survey Area (CA789):

Map Unit Component /Component %

101 Badgerwash /8

101 Humic Haploxerands, moist lake terrace /2

108 Humic Haploxerands, moist lake terrace /3

117 Humic Haploxerands, moist lake terrace /90

117 Badgerwash /2

172 Badgerwash /2

173 Badgerwash /90

Table 4. Representative soil features

Family particle size	(1) Sandy
Drainage class	Moderately well drained to well drained
Permeability class	Very rapid to very slow
Soil depth	50 – 150 cm
Surface fragment cover <=3"	10 – 70 %

Surface fragment cover >3"	0 – 20 %
Available water capacity (0-101.6cm)	1.09 – 8.94 cm
Soil reaction (1:1 water) (0-101.6cm)	5.5 – 7.3
Subsurface fragment volume <=3" (Depth not specified)	0 – 100 %
Subsurface fragment volume >3" (Depth not specified)	0 – 90 %

Ecological dynamics

This ecological site is on outwash with a root restrictive layer at varying depths between 20 to 60 inches. Although the heavy dominance of lodgepole pine here is partly in response to high fire frequency, the root restrictive layer and cold air drainage may exclude other conifers from establishing on this site.

In 2009, this ecological site within Lassen Volcanic National Park, is mainly dominated by a Sierra lodgepole pine (*Pinus contorta* var. *murrayana*) forest; however, outside the park it may be expressed by any of the plant communities described within the state and transition model. The canopy is relatively open due to frequent fires of varying intensity. A recent canopy fire has left some areas almost completely bare. Nothing remains; the standing dead and downed trees have been mostly consumed. There is very little regeneration in these areas. In other areas it appears that the understory burned at a low intensity, and a large portion of the overstory Sierra lodgepole pine will survive, leaving gap openings from the dead trees. The understory is dominated with grasses and forbs, including western needlegrass (*Achnatherum occidentale*), Ross' sedge (*Carex rossii*), squirreltail (*Elymus elymoides*), goldenbush (*Ericameria* spp.), narrowleaf lupine (*Lupinus angustifolius*), and lettuce wirelettuce (*Stephanomeria lactucina*).

Sierra lodgepole pine can be long-lived exceeding 150 years old. The overstory trees cored to obtain representative site index data were between 80 to 100 years old and relatively young. Sierra lodgepole pine does not usually gain much in girth with time and older trees averaged 16 to 21 inch diameters. Trees grow tall and narrow with short branches and 1.2 to 2.4 inch needles in fascicles of two. Its thin bark and shallow roots make it susceptible to fire. Sierra lodgepole pine is the only non-serotinous lodgepole pine. Therefore it does not need fire to open its cones to release seeds. The roots of Sierra lodgepole pine are generally shallow, which enables it to grow on this site. Sierra lodgepole pine produces a taproot that may atrophy or grow horizontally in cases of high water tables or root restrictive layers.

Sierra lodgepole pine has a complex disturbance regime that includes cyclic beetle infestations and fire. Fire studies in the lodgepole pine forest of the Caribou Wilderness report a fire return interval of 67 years between 1735 and 1929. Even low intensity fires resulted in high mortality rates for the lodgepole pine (Taylor and Solem, 1995). Sierra lodgepole pine regenerates prolifically after fire and evenly aged stands are formed. The mountain pine beetle (*Dendroctonus ponderosae*) is a natural pest that can kill a significant portion of the larger trees in a stand. Infestations can last for several years and often return in 20 to 40 year cycles (Cope, 1993). After an outbreak the forest may be dominated by standing dead trees. These trees eventually fall, creating layers of overlapping logs. Fuel loads are high, but the downed logs burn slowly and at a low intensity. As noted, even low intensity fire can cause damage to live trees, and fire damaged trees are rendered more susceptible to the next beetle attack. Pine beetle infestations, wind throw and other small scale disturbances create gaps for Sierra lodgepole pine or white fir regeneration. Over time these gaps will break up the uniformity of evenly aged stands that formed after the last large fire event.

White fir is presently not common on this site. This area has been burned repeatedly in the last century, which may have reduced its presence. The root restrictive layer and/or the cold air drainage may also keep white fir from firmly establishing. There are however sufficient white fir present in the understory to indicate that, without fire and with appropriate climate conditions, white fir could establish here. White fir is a large long-lived tree in this area. It commonly reaches 300 to 400 years in age and heights of 120 to 140 feet. It produces single needles 1.2 to 2.8 inches long that are distributed along young branches. Because the female seed cones open and fall apart while still attached to the tree, cones are not often seen on the forest floor. White fir tends to develop a shallow root system that can

graft to other white fir roots and spread root rots (Zouhar, 2001). White fir is a shade-tolerant conifer and is able to establish in the understory of the Sierra lodgepole pine on this site. If it continues to grow and reproduce in the understory in the absence of disturbance, it will eventually dominate the forest. In the past, the natural fire regime kept these forests from developing into the later successional stages dominated by white or red fir (Taylor, and Solem, 2001). White fir and Sierra lodgepole pine are both relatively fire intolerant species and tend to have high mortality rates after fire.

The mountain pine beetle is the most significant forest pathogen to affect this site, but several other pathogens have the potential to cause mortality or diminish productivity. Most of these pathogens represent natural cycles of regulation and can push the closed forest types into more open forest types. Large outbreaks are often associated with drought years or overstocked forests.

There is evidence that warming temperatures are allowing mountain pine beetles to exist farther north and into upper elevations. Warmer temperatures are altering the reproductive cycles and distribution of the mountain pine beetle. It is possible that the warmer temperatures will increase mountain pine beetle infestations for several decades. The southern mountain pine beetle may move northward due to temperature change as well (Carroll et al, 2003)

Pathogens that affect Sierra lodgepole pine include other insects such as the pine engraver (*Ips pini*), the weevil (*Magdalis gentiles*), the lodgepole terminal weevil (*Pissodes terminalis*), the Warren's collar weevil (*Hylobius warreni*), the pine needle scale (*Chionaspis pinifoliae*), the black pineleaf scale (*Nuculaspis californica*), the spruce spider mite (*Oligonychus ununguis*), the lodgepole sawfly (*Neodiprion burkei*), the lodgepole needle miner (*Coleotechnites milleri*), the sugar pine tortrix (*Choristoneura lambertiana*), the pine tube moth (*Argyrotaenia pinatubana*), and the pandora moth (*Coloradia pandora*). *Ips* commonly develops in logging slash, especially slash that is shaded and does not dry quickly. Prompt slash disposal is an effective control measure. *Ips* also can build up in windthrows. Fungal diseases that affect lodgepole pine productivity include the stem cankers caused by *atropelius* canker (*Atropellis piniphilia*), comandra blister rust (*Cronartium comandrae*), and western gall rust (*Peridermium harknessii*). The honey mushroom (*Armillaria mellea*) and annosus root disease (*Heterobasidion annosum*) are sources of root rot, and wood decay is caused by such fungi as red rot (*Phellinus pini*) and red heart wood stain (*Peniophora pseudo-pini*). Dwarf mistletoe (*Arceuthobium americanum*) is a common parasite that affects large areas of lodgepole pine (Lotan and Critchfield, 1990).

Pathogens that affect white fir are the dwarf mistletoe (*Arceuthobium abietinum* f. sp. *concoloris*), Cytospora canker (*Cytospora abietis*), broom rust (*Melampsorella caryophyllacearum*), annosus root disease (*Heterobasidium annosum*), armillaria root disease (*Armillaria* sp.), trunk rot (*Echinodontium tinctorium*) and the fir engraver (*Scotylus ventralis*). The most threatening of these is the combination of the fir engraver and annosus root disease. These pathogens can kill large areas of white fir (Bohne, 2006; Laacke, 1990).

The reference state consists of the most successional advanced community phase (numbered 1.1) as well as other community phases which result from natural and human disturbances. Community phase 1.1 is deemed the phase representative of the most successional advanced pre-European plant/animal community including periodic natural surface fires that influenced its composition and production. Because this phase is determined from the oldest modern day remnant forests and/or historic literature, some speculation is necessarily involved in describing it.

All tabular data listed for a specific community phase within this ecological site description represent a summary of one or more field data collection plots taken in communities within the community phase. Although such data are valuable in understanding the phase (kinds and amounts of ground and surface materials, canopy characteristics, community phase overstory and understory species, production and composition, and growth), it typically does not represent the absolute range of characteristics nor an exhaustive listing of species for all the dynamic communities within each specific community phase.

State and transition model

Additional community tables

Table 5. Community 1.1 plant community composition

Group	Common Name	Symbol	Scientific Name	Annual Production ()	Foliar Cover (%)
Tree					
0	Tree (understory only)			3-17	
	Sierra lodgepole pine	PICOM	<i>Pinus contorta</i> var. <i>murrayana</i>	3-17	1-5
Grass/Grasslike					
0	Grass/Grasslike			16-80	
	squirreltail	ELEL5	<i>Elymus elymoides</i>	9-36	2-8
	western needlegrass	ACOC3	<i>Achnatherum occidentale</i>	7-27	2-6
	Ross' sedge	CARO5	<i>Carex rossii</i>	0-17	0-5

Forb					
0	Forb			0-68	
	narrowleaf lupine	LUAN4	<i>Lupinus angustifolius</i>	0-54	0-15
	goldenbush	ERICA2	<i>Ericameria</i>	0-13	0-2
	lettuce wirelettuce	STLA	<i>Stephanomeria lactucina</i>	0-1	0-1

Table 6. Community 1.1 forest overstory composition

Common Name	Symbol	Scientific Name	Nativity	Height M	Canopy Cover (%)	Diameter Cm	Basal Area (square M/hectare)
Tree							
Sierra lodgepole pine	PICOM	<i>Pinus contorta var. murrayana</i>	Native	15.2-24.4	35-60	30.5-43.2	23-48.2

Table 7. Community 1.1 forest understory composition

Common Name	Symbol	Scientific Name	Nativity	Height (m)	Canopy Cover (%)
Grass/grass-like (Graminoids)					
squirreltail	ELEL5	<i>Elymus elymoides</i>	Native	–	2-8
western needlegrass	ACOC3	<i>Achnatherum occidentale</i>	Native	–	2-6
Ross' sedge	CARO5	<i>Carex rossii</i>	Native	–	0-5
Forb/Herb					
narrowleaf lupine	LUAN4	<i>Lupinus angustifolius</i>	Native	–	0-15
goldenbush	ERICA2	<i>Ericameria</i>	Native	–	0-2
lettuce wirelettuce	STLA	<i>Stephanomeria lactucina</i>	Native	–	0-1
Tree					
Sierra lodgepole pine	PICOM	<i>Pinus contorta var. murrayana</i>	Native	–	1-5

Table 8. Community 1.2 plant community composition

Group	Common Name	Symbol	Scientific Name	Annual Production ()	Foliar Cover (%)
-------	-------------	--------	-----------------	----------------------	------------------

Table 9. Community 1.3 plant community composition

Group	Common Name	Symbol	Scientific Name	Annual Production ()	Foliar Cover (%)
-------	-------------	--------	-----------------	----------------------	------------------

Table 10. Community 1.4 plant community composition

Group	Common Name	Symbol	Scientific Name	Annual Production ()	Foliar Cover (%)
-------	-------------	--------	-----------------	----------------------	------------------

Table 11. Community 1.5 plant community composition

Group	Common Name	Symbol	Scientific Name	Annual Production ()	Foliar Cover (%)
-------	-------------	--------	-----------------	----------------------	------------------

Table 12. Community 1.6 plant community composition

Group	Common Name	Symbol	Scientific Name	Annual Production ()	Foliar Cover (%)
-------	-------------	--------	-----------------	----------------------	------------------

Animal community

Sierra lodgepole pine forests provide food, cover and habitat for a variety of species. There are 31 mammals and almost 50 bird species documented in Sierra lodgepole pine forests. Snags and downed logs are important for cavity-nesting birds and mammals. Other animals feed on the Sierra lodgepole pine needles and consume the seeds (Cope, 1993).

Recreational uses

This area is suitable for trails and camping.

Wood products

Sierra lodgepole pine wood is used for framing, paneling, trim, posts, and other construction products. The forests are often uniform in size, which makes harvesting easier. The wood tends to be light and straight grained with consistent texture (Cope 1993).

Other information

Alexander (1966) was used to determine forest site productivity for lodgepole pine. Low to High values of Site index and CMAI (culmination of mean annual increment) give an indication of the range of inherent productivity of this ecological site. Site index relates to height of dominant trees over a set period of time and CMAI relates to the average annual growth of wood fiber in the boles/trunks of trees. Site index and CMAI listed in the Forest Site Productivity section are in units of feet and cubic feet/acre/year, respectively. Both site index and CMAI are estimates; on-site investigation is recommended for specific forest management units for each soil classified to this ecological site. The historical and actual basal area of trees within a growing stand will greatly influence CMAI. Conifer trees appropriate for site index measurement typically occur in community phase 1.4 and older stands in 1.2 and 1.3. They are selected according to guidance listed in the site index publications.

Table 13. Representative site productivity

Common Name	Symbol	Site Index Low	Site Index High	CMAI Low	CMAI High	Age Of CMAI	Site Index Curve Code	Site Index Curve Basis	Citation
Sierra lodgepole pine	PICOM	68	89	66	102	100	520	100TA	Alexander, Robert R. 1966. Site indexes for Lodgepole pine, with corrections for stand density: instructions for field use. USDA, Forest Service. Rocky Mountain Forest and Range Experiment Station Research Paper RM-24.
Sierra lodgepole pine	PICOM	68	89	57	78	—	—	—	

Inventory data references

The following NRCS vegetation plots were used to describe this ecological site. 789110 789369- Type location 789397

Type locality

Location 1: Shasta County, CA	
Township/Range/Section	T31 N R5 E S10
UTM zone	N
UTM northing	4490710
UTM easting	634578
General legal description	The type location is about 1.6 miles west northwest from the western edge of Soap Lake in Lassen Volcanic National Park.

Other references

Agee, James K.1994. The Lodgepole Pine Series in Fire and Weather Disturbances in Terrestrial Ecosystems of the Eastern Cascades. From volume III: Assessment. USDA, Forest Service, Pacific Northwest Research Station. Gen. Tech. Report.

Alexander, Robert R. 1966. Site indexes for Lodgepole pine, with corrections for stand density: instructions for field use. USDA, Forest Service. Rocky Mountain Forest and Range Experiment Station Research Paper RM-24. NASIS ID 520

Amman, Gene D., McGregor, Mark D., Dolph Robert E. 1990. Mountain Pine Beetle: Forest Insect and Disease Leaflet 2. USDA, Forest Service, Pacific Northwest Region, Portland OR.

Beaty, Matthew and Taylor, Alan H. (2001). Spatial and Temporal Variation of Fire Regimes in a Mixed Conifer Forest Landscape, Southern Cascades, California, USA. *Journal of Biogeography*, 28, 955-966.

Bekker, Mathew F. and Taylor, Alan H. (2001). Gradient Analysis of Fire Regimes in Montane Forest of the Southern Cascade Range, Thousand Lakes Wilderness, California, USA. *Plant Ecology* 155: 15-23.

Burns, Russell M., and Barbara H. Honkala, tech. coords. 1990. *Silvics of North America: 1. Conifers; 2. Hardwoods*. Agriculture Handbook 654. U.S. Department of Agriculture, Forest Service, Washington, DC. vol.2, 877 p.

Carroll, Allan L.; Taylor, Steve W.; Régnière, Jacques; and Safranyik, Les. 2003. Effects of Climate Change on Range Expansion by the Mountain Pine Beetle in British Columbia. *Mountain Pine Beetle Symposium: Challenges and Solutions*. October 30-31, 2003, Kelowna, British Columbia. T.L. Shore, J.E. Brooks, and J.E. Stone (editors). Natural Resources Canada, Canadian Forest Service, Pacific Forestry Centre, Information Report BC-X-399, Victoria, BC. 298 p.

Cope, Amy, B. 1993. *Pinus contorta* var. *murrayana*. In: *Fire Effects Information Systems*, U.S. department of Agriculture, Forest Service, Rocky Mountain Research Station, Fire Science Laboratory (Producer). <http://www.fs.fed.us/database/feis/>

Lotan, James, E. and Critchfield, William B., 1990. *Pinus contorta*: Lodgepole Pine In: Burns, Russel M., Honkala, Barbara H. eds. *Silvics of North America*, Vol 1. Conifers.

Parker, Albert J., 1995. Comparative Gradient Structure and Forest Cover Types in Lassen Volcanic and Yosemite National Parks, California. *Bulletin of the Torrey Botanical Club*, Vol. 122, No. 1. (Jan. - Mar., 1995), pp. 58-68.

Millar, Constance I.; Westfall, Robert D.; Delany, Diane L.; King, John C.; and Graumlich, Lisa J., 2004. Response of Subalpine Conifers in the Sierra Nevada, California, U.S.A., to 20th Century Warming and Decadal Climate Variability. *Arctic, Antarctic, and Alpine Research*, Vol. 36, No. 2, 2004, pp. 181–200.

Parker, Albert J., 1991. Forest/Environment Relationships in Lassen Volcanic National Park, California, U.S.A. *Journal of Biogeography*, Vol. 18, No. 5. (Sep., 1991), pp. 543-552.

Potter, Donald; Smith, Mark; Beck, Tom; Kermeen, Brian; Hance, Wayne; and Robertson, Steve; 1992. *Ecological Characteristics of Old Growth Lodgepole Pine in California*. USDA, Forest Service.

Potter, Donald A. (1998). *Forested Communities of the Upper Montane in the Central and Southern Sierra Nevada*. U.S. Department of Agriculture, Forest Service, Pacific Southwest Research Station, General Technical Report PSW-GTR-169.

Royce, E. B. and Barbour, M. G., 2001. Mediterranean Climate Effects. I. Conifer Water Use Across a Sierra Nevada Ecotone. *American Journal of Botany* 88(5): 911–918. 2001.

Royce, E. B. and Barbour, M. G., 2001. Mediterranean Climate Effects. II. Conifer Growth Phenology Across a Sierra Nevada Ecotone. *American Journal of Botany* 88(5): 919–932. 2001.

Schumacher, Francis X. 1926. Yield, stand, and volume tables for white fir in the California pine region. University of California Agricultural Experiment Station Bulletin 407. NASIS ID 030

Taylor, Alan. H., 1990. Tree Invasion in Meadows of Lassen Volcanic National Park, California. *Professional Geographer*, 42(4), 1990, pp. 457- 470.

Taylor, Alan. H., 2000. Fire Regimes and Forest Changes in Mid and Upper Montane Forest of the Southern Cascades, Lassen Volcanic National Park, California, U.S.A. *Journal of Biogeography*, 27, 87-104.

Taylor, Alan H. and Solem, Michael N., 2001. Fire Regimes and Stand Dynamics in an Upper Montane Forest Landscape in the Southern Cascades, Caribou Wilderness, California. *Journal of the Torrey Botanical Society*, Vol. 128, No. 4. (Oct. - Dec., 2001), pp. 350-361.

Contributors

Lyn Townsend
Marchel M. Munnecke

Rangeland health reference sheet

Interpreting Indicators of Rangeland Health is a qualitative assessment protocol used to determine ecosystem condition based on benchmark characteristics described in the Reference Sheet. A suite of 17 (or more) indicators are typically considered in an assessment. The ecological site(s) representative of an assessment location must be known prior to applying the protocol and must be verified based on soils and climate. Current plant community cannot be used to identify the ecological site.

Author(s)/participant(s)	
Contact for lead author	
Date	
Approved by	
Approval date	
Composition (Indicators 10 and 12) based on	Annual Production

Indicators

1. Number and extent of rills:

2. Presence of water flow patterns:

3. Number and height of erosional pedestals or terracettes:

4. Bare ground from Ecological Site Description or other studies (rock, litter, lichen, moss, plant canopy are not bare ground):

5. Number of gullies and erosion associated with gullies:

6. Extent of wind scoured, blowouts and/or depositional areas:

7. Amount of litter movement (describe size and distance expected to travel):

8. Soil surface (top few mm) resistance to erosion (stability values are averages - most sites will show a range of values):

9. Soil surface structure and SOM content (include type of structure and A-horizon color and thickness):

10. Effect of community phase composition (relative proportion of different functional groups) and spatial distribution on infiltration and runoff:

11. Presence and thickness of compaction layer (usually none; describe soil profile features which may be mistaken for compaction on this site):

12. Functional/Structural Groups (list in order of descending dominance by above-ground annual-production or live foliar cover using symbols: >>, >, = to indicate much greater than, greater than, and equal to):

Dominant:

Sub-dominant:

Other:

Additional:

13. Amount of plant mortality and decadence (include which functional groups are expected to show mortality or decadence):

14. Average percent litter cover (%) and depth (in):

15. Expected annual annual-production (this is TOTAL above-ground annual-production, not just forage annual-production):

16. Potential invasive (including noxious) species (native and non-native). List species which BOTH characterize degraded states and have the potential to become a dominant or co-dominant species on the ecological site if their future establishment and growth is not actively controlled by management interventions. Species that become dominant for only one to several years (e.g., short-term response to drought or wildfire) are not invasive plants. Note that unlike other indicators, we are describing what is NOT expected in the reference state for the ecological site:

17. Perennial plant reproductive capability:
