

# Ecological site R023XY049NV GRANITIC SOUTH SLOPE 8-12 P.Z.

Last updated: 4/10/2025  
Accessed: 04/20/2026

---

## General information

**Provisional.** A provisional ecological site description has undergone quality control and quality assurance review. It contains a working state and transition model and enough information to identify the ecological site.

## Ecological site concept

Currently there is only a draft of the initial concept for this ecological site. The initial concept for this site places it within the Clayey Mesic Plateaus 8-14 PZ Wyoming Big Sagebrush and Thurber's Needlegrass Ecological Site Group. To view the General STM and other information available for this ESG please go to <https://edit.jornada.nmsu.edu/catalogs/esg/023X/R023XY909OR> University of Nevada Reno Researchers note that this site is slightly more productive than the group modal, ranging from 500 to 900 lb/ac. The plant community is dominated by Wyoming big sagebrush with an understory dominated by bluebunch wheatgrass and Thurber's needlegrass. The soils of this site have formed in residuum derived from granitic rocks. This site occurs on smooth to convex shoulders and sideslopes of hills and lower mountains on predominantly southerly aspects. Slopes are greater than the modal site, ranging from 15 to 75 percent, but slope gradients of 30 to 50 percent are most typical. This site has not been seen on field visits for the group report, but likely has the same STM as the modal site.

## Associated sites

<b>R023XY040NV</b>	<b>GRANITIC FAN 8-10 P.Z.</b>
--------------------	-------------------------------

## Similar sites

<b>R023XY057NV</b>	<b>GRANITIC LOAM 10-12 P.Z.</b>  ACTH7 dominant grass; less productive site
--------------------	---

**Table 1. Dominant plant species**

Tree	Not specified
Shrub	(1) <i>Artemisia tridentata</i> var. <i>wyomingensis</i>
Herbaceous	(1) <i>Pseudoroegneria spicata</i> ssp. <i>spicata</i>

## Physiographic features

This site occurs on smooth to convex shoulders and sideslopes of hills and lower mountains on predominately southerly aspects. Slopes range from 15 to 50 percent. Elevations are 4800 to about 5700 feet.

**Table 2. Representative physiographic features**

Landforms	(1) Mountain slope (2) Hill
Elevation	1,460 – 1,740 m
Slope	20 – 50 %
Aspect	S

### Climatic features

The climate associated with this site is semiarid and characterized by cool, moist winters and warm, dry summers. Average annual precipitation is 8 to 12 inches. Mean annual air temperature is 44 to 52 degrees F. The average growing season is about 90 to 120 days.

Nevada's climate is predominantly arid, with large daily ranges of temperature, infrequent severe storms, heavy snowfall in the higher mountains, and great location variations with elevation. Three basic geographical factors largely influence Nevada's climate: continentality, latitude, and elevation. Continentality is the most important factor. The strong continental effect is expressed in the form of both dryness and large temperature variations. Nevada lies on the eastern, lee side of the Sierra Nevada Range, a massive mountain barrier that markedly influences the climate of the State. The prevailing winds are from the west, and as the warm moist air from the Pacific Ocean ascend the western slopes of the Sierra Range, the air cools, condensation occurs and most of the moisture falls as precipitation. As the air descends the eastern slope, it is warmed by compression, and very little precipitation occurs. The effects of this mountain barrier are felt not only in the West but throughout the state, with the result that the lowlands of Nevada are largely desert or steppes. The temperature regime is also affected by the blocking of the inland-moving maritime air. Nevada sheltered from maritime winds, has a continental climate with well-developed seasons and the terrain responds quickly to changes in solar heating.

Nevada lies within the mid-latitude belt of prevailing westerly winds which occur most of the year. These winds bring frequent changes in weather during the late fall, winter and spring months, when most of the precipitation occurs. To the south of the mid-latitude westerlies, lies a zone of high pressure in subtropical latitudes, with a center over the Pacific Ocean. In the summer, this high-pressure belt shifts northward over the latitudes of Nevada, blocking storms from the ocean. The resulting weather is mostly clear and dry during the summer and early fall, with scattered thundershowers. The eastern portion of the state receives significant summer thunderstorms generated from monsoonal moisture pushed up from the Gulf of California, known as the North American monsoon. The monsoon system peaks in August and by October the monsoon high over the Western U.S. begins to weaken and the precipitation retreats southward towards the tropics (NOAA 2004).

Average annual precipitation is 16 to over 20 inches. Mean annual air temperature is 41 to 44 degrees F. The average growing season is about 50 to 70 days.

Mean annual precipitaion at the Bear Creek, Nevada SNOTEL station (170501020301) is 37.69 inches.

monthly mean precipitation is:

January 3.84; February 3.75; March 4.38; April 4.9;  
 May 3.99; June 2.82; July .95; August 1.66;  
 September 1.22; October 2.12;  
 November 3.67; December 4.38.

**Table 3 Representative climatic features**

Frost-free period (average)	110 days
Freeze-free period (average)	
Precipitation total (average)	250 mm

## Influencing water features

There are no influencing water features associated with this site.

## Soil features

The soils associated with this site have formed in residuum derived from granitic rocks. They are shallow and well drained. Rock fragments average 35 to 60 percent in the profile. The soils have a mollic epipedon and an argillic horizon. The soil series associated with this site include: Acrelane.

Table 4. Representative soil features

Surface texture	(1) Extremely gravelly coarse sand
Family particle size	(1) Loamy
Drainage class	Well drained
Permeability class	Moderate
Soil depth	30 – 50 cm
Surface fragment cover <=3"	50 %
Surface fragment cover >3"	10 – 20 %
Available water capacity (0-101.6cm)	2.79 – 3.05 cm
Calcium carbonate equivalent (0-101.6cm)	Not specified
Electrical conductivity (0-101.6cm)	Not specified
Sodium adsorption ratio (0-101.6cm)	Not specified
Soil reaction (1:1 water) (0-101.6cm)	5.6 – 7.8

Subsurface fragment volume <=3" (Depth not specified)	20 – 50 %
Subsurface fragment volume >3" (Depth not specified)	0 – 20 %

## Ecological dynamics

Where management results in abusive use by livestock or feral horses, unpalatable brush species increase as bluebunch wheatgrass and other desirable forage grasses decline in the understory. Cheatgrass and other annuals are plants likely to invade this site.

### Fire Ecology:

The fire return interval for Wyoming big sagebrush communities ranges from 10 to 70 years. Fire is the principal means of renewal for decadent stands of Wyoming big sagebrush. Wyoming big sagebrush is killed by fire and establishes after fire from a seedbank; from seed produced by remnant plants that escaped fire; and from plants adjacent to the burn that seed in. Ephedra generally sprouts vigorously from the roots or woody root crown after fire and rapidly produces aboveground biomass from surviving meristematic tissue. Burning bluebunch wheatgrass may remove most of the aboveground biomass but does not usually result in plant mortality. Bluebunch wheatgrass is generally favored by burning. Burning stimulates flowering and seed production. However, season of burning affects mortality. Thurber's needlegrass is classified as moderately resistant, but depending on season of burn, phenology, and fire severity, this perennial bunchgrass is moderately to severely damaged by fire. Early season burning is more damaging to this needlegrass than late season burning. Desert needlegrass has persistent dead leaf bases, which make it susceptible to burning. Fire removes the accumulation; a rapid, cool fire will not burn deep into the root crown. Most perennial grasses have root crowns that can survive wildfire. Indian ricegrass can be killed by fire, depending on severity and season of burn. Indian ricegrass reestablishes on burned sites through seed dispersed from adjacent unburned areas.

## State and transition model

### Additional community tables

Table 5. Community 1.1 plant community composition

Group	Common Name	Symbol	Scientific Name	Annual Production ()	Foliar Cover (%)
<b>Grass/Grasslike</b>					
1	<b>Primary Perennial Grasses</b>			345-549	
	bluebunch wheatgrass	PSSPS	<i>Pseudoroegneria spicata ssp. spicata</i>	235-314	–
	Thurber's needlegrass	ACTH7	<i>Achnatherum thurberianum</i>	78-157	–
	Indian ricegrass	ACHY	<i>Achnatherum hymenoides</i>	16-39	–
	desert needlegrass	ACSP12	<i>Achnatherum speciosum</i>	16-39	–
2	<b>Secondary Perennial Grasses</b>			39-78	
	squirreltail	ELEL5	<i>Elymus elymoides</i>	4-24	–
	needle and thread	HECO26	<i>Hesperostipa comata</i>	4-24	–
	Sandberg bluegrass	POSE	<i>Poa secunda</i>	4-24	–
<b>Forb</b>					
3	<b>Perennial</b>			16-78	
	milkvetch	ASTRA	<i>Astragalus</i>	4-16	–
	sulphur-flower buckwheat	ERUM	<i>Eriogonum umbellatum</i>	4-16	–
	lupine	LUPIN	<i>Lupinus</i>	4-16	–
	phlox	PHLOX	<i>Phlox</i>	4-16	–
<b>Shrub/Vine</b>					
4	<b>Primary Shrubs</b>			196-314	
	Wyoming big sagebrush	ARTRW8	<i>Artemisia tridentata ssp. wyomingensis</i>	157-235	–
	jointfir	EPHED	<i>Ephedra</i>	39-78	–

5	<b>Secondary Shrubs</b>			16-63	
	yellow rabbitbrush	CHVI8	<i>Chrysothamnus viscidiflorus</i>	8-16	–
	desert peach	PRAN2	<i>Prunus andersonii</i>	8-16	–
	antelope bitterbrush	PUTR2	<i>Purshia tridentata</i>	8-16	–
	purple sage	SADOI	<i>Salvia dorrii ssp. dorrii var. incana</i>	8-16	–

## Animal community

**Livestock Interpretations:** This site has limited value for livestock grazing due to steep slopes. Bluebunch wheatgrass is considered one of the most important forage grass species on western rangelands for livestock. Although bluebunch wheatgrass can be a crucial source of forage, it is not necessarily the most highly preferred species. Thurber's needlegrass species begin growth early in the year and remain green throughout a relatively long growing season. This pattern of development enables animals to use Thurber's needlegrass when many other grasses are unavailable. Cattle prefer Thurber's needlegrass in early spring before fruits have developed as it becomes less palatable when mature. Thurber's needlegrasses are grazed in the fall only if the fruits are softened by rain. Young desert needlegrass is palatable to all classes of livestock. Mature herbage is moderately grazed by horses and cattle, but rarely grazed by sheep. Indian ricegrass is highly palatable to all classes of livestock in both green and cured condition. It supplies a source of green feed before most other native grasses have produced much new growth. Livestock browse Wyoming big sagebrush, but may use it only lightly when palatable herbaceous species are available. Ephedra is heavily browsed by livestock on winter range but only moderately or lightly browsed during other seasons. Stocking rates vary over time depending upon season of use, climate variations, site, and previous and current management goals. A safe starting stocking rate is an estimated stocking rate that is fine tuned by the client by adaptive management through the year and from year to year. **Wildlife Interpretations:** Wyoming big sagebrush is preferred browse for wild ungulates. Pronghorn usually browse Wyoming big sagebrush heavily. Sagebrush-grassland communities provide critical sage-grouse breeding and nesting habitats. Meadows surrounded by sagebrush may be used as feeding and strutting grounds. Sagebrush is a crucial component of their diet year-round, and sage-grouse select sagebrush almost exclusively for cover. Sage-grouse prefer mountain big sagebrush and Wyoming big sagebrush communities to basin big sagebrush communities. Ephedra is an important browse species for big game animals. Ephedra is heavily used by wildlife on winter ranges. Bluebunch wheatgrass is considered one of the most important forage grass species on western rangelands for wildlife. Bluebunch wheatgrass does not generally provide sufficient cover for ungulates, however, mule deer are frequently found in bluebunch-dominated grasslands. Thurber needlegrass is valuable forage for wildlife. Young desert needlegrass is palatable to many species of wildlife. Desert needlegrass produces considerable basal foliage and is good forage while young. Indian ricegrass is eaten by pronghorn in moderate amounts whenever available. A number of heteromyid rodents inhabiting desert rangelands show preference for seed of Indian ricegrass. Indian ricegrass is an important component of jackrabbit diets in spring and summer. Indian ricegrass seed provides food for many species of birds. Doves, for example, eat large amounts of shattered Indian ricegrass seed lying on the ground.

## Hydrological functions

Runoff is high. Permeability is moderate. Hydrologic soil group is C.

## Recreational uses

Aesthetic value is derived from the diverse floral and faunal composition and the colorful flowering of wild flowers and shrubs during the spring and early summer. This site offers rewarding opportunities to photographers and for nature study. This site is used for hiking and has potential for upland and big game hunting.

## Other products

Native Americans made tea from big sagebrush leaves. They used the tea as a tonic, an antiseptic, for treating colds, diarrhea, and sore eyes and as a rinse to ward off ticks. Big sagebrush seeds were eaten raw or made into meal. Indian ricegrass was traditionally eaten by some Native Americans. The Paiutes used the seed as a reserve food source.

## Other information

Wyoming big sagebrush is used for stabilizing slopes and gullies and for restoring degraded wildlife habitat, rangelands, mine spoils and other disturbed sites. It is particularly recommended on dry upland sites where other shrubs are difficult to establish. Ephedra is listed as a successful shrub for restoring western rangeland communities and can be used to rehabilitate disturbed lands. It also has value for reducing soil erosion on both clay and sandy soils. Ephedra establishes readily through direct seeding, transplants, and stem cuttings.

## Type locality

Location 1: Washoe County, NV

Township/Range/Section	T29N R22E S7
UTM zone	N
UTM northing	283164
UTM easting	4475162
Latitude	40° 23' 55"
Longitude	119° 33' 18"
General legal description	NE 1/4 SW 1/4, Approximately ½ mile northeast of the end of the road in Rattlesnake Canyon, Fox Range, Washoe County, Nevada.

**Other references**

Fire Effects Information System (Online; <http://www.fs.fed.us/database/feis/plants/>).

USDA-NRCS Plants Database (Online; <http://www.plants.usda.gov>).

Great Basin Ecological Site Development Project: State and Transition Models for Major Land Resource Area 23, Nevada and portions of California (Online; <https://naes.agnt.unr.edu/PMS/Pubs/2019-4060.pdf>)

**Contributors**

SW  
T Stringham (UNR under contract with BLM)

**Approval**

Kendra Moseley, 4/10/2025

**Rangeland health reference sheet**

**Interpreting Indicators of Rangeland Health** is a qualitative assessment protocol used to determine ecosystem condition based on benchmark characteristics described in the Reference Sheet. A suite of 17 (or more) indicators are typically considered in an assessment. The ecological site(s) representative of an assessment location must be known prior to applying the protocol and must be verified based on soils and climate. Current plant community cannot be used to identify the ecological site.

Author(s)/participant(s)	
Contact for lead author	
Date	04/20/2026

Approved by	
Approval date	
Composition (Indicators 10 and 12) based on	Annual Production

## Indicators

**1. Number and extent of rills:**

---

**2. Presence of water flow patterns:**

---

**3. Number and height of erosional pedestals or terracettes:**

---

**4. Bare ground from Ecological Site Description or other studies (rock, litter, lichen, moss, plant canopy are not bare ground):**

---

**5. Number of gullies and erosion associated with gullies:**

---

**6. Extent of wind scoured, blowouts and/or depositional areas:**

---

**7. Amount of litter movement (describe size and distance expected to travel):**

---

**8. Soil surface (top few mm) resistance to erosion (stability values are averages - most sites will show a range of values):**

---

**9. Soil surface structure and SOM content (include type of structure and A-horizon color and thickness):**

---

**10. Effect of community phase composition (relative proportion of different functional groups) and spatial distribution on infiltration and runoff:**

---

**11. Presence and thickness of compaction layer (usually none; describe soil profile features which may be mistaken for compaction on this site):**

---

12. **Functional/Structural Groups** (list in order of descending dominance by above-ground annual-production or live foliar cover using symbols: >>, >, = to indicate much greater than, greater than, and equal to):

**Dominant:**

**Sub-dominant:**

**Other:**

**Additional:**

---

13. **Amount of plant mortality and decadence** (include which functional groups are expected to show mortality or decadence):

---

14. **Average percent litter cover (%) and depth ( in):**

---

15. **Expected annual annual-production** (this is TOTAL above-ground annual-production, not just forage annual-production):

---

16. **Potential invasive (including noxious) species (native and non-native).** List species which BOTH characterize degraded states and have the potential to become a dominant or co-dominant species on the ecological site if their future establishment and growth is not actively controlled by management interventions. Species that become dominant for only one to several years (e.g., short-term response to drought or wildfire) are not invasive plants. Note that unlike other indicators, we are describing what is NOT expected in the reference state for the ecological site:

---

17. **Perennial plant reproductive capability:**

---