

Ecological site R023XY071NV ASHY LOAM 10-12 P.Z.

Last updated: 4/10/2025
Accessed: 04/20/2026

General information

Provisional. A provisional ecological site description has undergone quality control and quality assurance review. It contains a working state and transition model and enough information to identify the ecological site.

Ecological site concept

Currently there is only a draft of the initial concept for this ecological site. The initial concept for this site places it within the Ashy or Loamy Skeletal Mod Deep 10-20 PZ High-Resilience Mountain Big Sagebrush and Idaho Fescue Ecological Site Group. To view the General STM and other information available for this ESG please go to <https://edit.jornada.nmsu.edu/catalogs/esg/023X/R023XY906NV>. The site is slightly less productive than the modal site with 1100 lbs/ac in normal years. This site is found on concave positions of upper fan piedmonts and inset fans with slope typically ranging from 2 to 4 percent. Soils on this site are moderately deep to deep and available water capacity is high in part due to the high amounts of volcanic ash and glass throughout the soil profile. This site is dominated by basin big sagebrush rather than mountain big sagebrush. Unlike the group modal site, the grass community is dominated by Idaho fescue and Thurber's needlegrass with bluebunch wheatgrass representing only a small component. This site is likely similar to the group modal site with 2 stable states, however this site was not seen during site visits for the group report.

Associated sites

R023XY020NV	LOAMY 10-12 P.Z.
R023XY031NV	CLAYPAN 10-14 P.Z.
R023XY072NV	ASHY SLOPE 10-12 P.Z.
R023XY096NV	ASHY SANDY LOAM 10-12 P.Z.

Similar sites

R023XY082NV	LOAMY FAN 10-12 P.Z. ACTH7-POA-LECI4 codominant grasses
R023XY066NV	ASHY LOAM 14-16 P.Z. ARTRV-PUTR2 codominant shrubs

R023XY094NV	ASHY SLOPE 12-14 P.Z. ARTRV dominant shrub
R023XY020NV	LOAMY 10-12 P.Z. PSSPS-ACTH7 codominant grasses
R023XY084NV	DEEP LOAMY 14-16 P.Z. PSSPS-ACTH7-FEID codominant grasses
R023XY072NV	ASHY SLOPE 10-12 P.Z. ARTRW dominant shrub
R023XY007NV	LOAMY 14-16 P.Z. ARTRV dominant shrub

Table 1. Dominant plant species

Tree	Not specified
Shrub	(1) <i>Artemisia tridentata ssp. tridentata</i>
Herbaceous	(1) <i>Festuca idahoensis</i> (2) <i>Achnatherum thurberianum</i>

Physiographic features

This site occurs on inset fans, landslides, and ash flows. Slopes range from 0 to 50 percent, but slope gradients of 0 to 4 percent are most typical. Elevations are 5700 to 6500 feet.

Table 2. Representative physiographic features

Landforms	(1) Inset fan (2) Landslide (3) Ash flow
Elevation	1,740 – 1,980 m
Slope	0 – 50 %
Aspect	Aspect is not a significant factor

Climatic features

The climate associated with this site is semiarid and characterized by cool, moist winters and warm, dry summers. Average annual precipitation is 10 to 12 inches. Mean annual air temperature is 42 to 46 degrees F. The average growing season is about 70 to 100 days.

Nevada's climate is predominantly arid, with large daily ranges of temperature, infrequent severe storms, heavy snowfall in the higher mountains, and great location variations with elevation. Three basic geographical factors largely influence Nevada's climate: continentality, latitude, and elevation. Continentality is the most important factor. The strong continental effect is expressed in the form of both dryness and large temperature variations. Nevada lies on the eastern, lee side of the Sierra Nevada Range, a massive mountain barrier that markedly influences the climate of the State. The prevailing winds are from the west, and as the warm moist air from the Pacific Ocean ascend the western slopes of the Sierra Range, the air cools, condensation occurs and most of the moisture falls as precipitation. As the air descends the eastern slope, it is warmed by compression, and very little precipitation occurs. The effects of this mountain barrier are felt not only in the West but throughout the state, with the result that the lowlands of Nevada are largely desert or steppes. The temperature regime is also affected by the blocking of the inland-moving maritime air. Nevada sheltered from maritime winds, has a continental climate with well-developed seasons and the terrain responds quickly to changes in solar heating.

Nevada lies within the mid-latitude belt of prevailing westerly winds which occur most of the year. These winds bring frequent changes in weather during the late fall, winter and spring months, when most of the precipitation occurs. To the south of the mid-latitude westerlies, lies a zone of high pressure in subtropical latitudes, with a center over the Pacific Ocean. In the summer, this high-pressure belt shifts northward over the latitudes of Nevada, blocking storms from the ocean. The resulting weather is mostly clear and dry during the summer and early fall, with scattered thundershowers. The eastern portion of the state receives significant summer thunderstorms generated from monsoonal moisture pushed up from the Gulf of California, known as the North American monsoon. The monsoon system peaks in August and by October the monsoon high over the Western U.S. begins to weaken and the precipitation retreats southward towards the tropics (NOAA 2004).

Average annual precipitation is 16 to over 20 inches. Mean annual air temperature is 41 to 44 degrees F. The average growing season is about 50 to 70 days.

Mean annual precipitation at the Bear Creek, Nevada SNOTEL station (170501020301) is 37.69 inches.

monthly mean precipitation is:

January 3.84; February 3.75; March 4.38; April 4.9;

May 3.99; June 2.82; July .95; August 1.66;

September 1.22; October 2.12;

November 3.67; December 4.38.

Table 3 Representative climatic features

Frost-free period (average)	90 days
Freeze-free period (average)	
Precipitation total (average)	280 mm

Influencing water features

There are no influencing water features associated with this site.

Soil features

The soils associated with this site are moderately deep to very deep and are formed in residuum and colluvium from volcanic rocks or mixed rocks and volcanic ash. These soils are coarse to moderately coarse textured throughout. Surface soils are neutral. Infiltration of water is rapid due to the relatively coarse textured surface soils. Available water capacity is high. There are very high amounts of vitric volcanic ash and glass throughout the soil profile which enhances the water holding capacity of these soils. Permeability is moderately slow to moderately rapid and runoff is very low. The soil series associated with this site include: Yellowhills.

Table 4. Representative soil features

Surface texture	(1) Sandy loam
Family particle size	(1) Loamy
Drainage class	Well drained
Permeability class	Moderately slow to moderately rapid
Soil depth	50 – 210 cm
Surface fragment cover ≤3"	Not specified
Surface fragment cover >3"	Not specified
Available water capacity (0-101.6cm)	13.72 – 23.11 cm
Calcium carbonate equivalent (0-101.6cm)	Not specified
Electrical conductivity (0-101.6cm)	Not specified
Sodium adsorption ratio (0-101.6cm)	Not specified
Soil reaction (1:1 water) (0-101.6cm)	6.1 – 7.8
Subsurface fragment volume ≤3" (Depth not specified)	0 – 10 %
Subsurface fragment volume >3" (Depth not specified)	Not specified

Ecological dynamics

Where management results in abusive grazing use by livestock and/or feral horses, woody plants, bottlebrush squirreltail, and Sandberg bluegrass increase as Idaho fescue, Thurber's needlegrass and other desirable forage grasses decrease. Thistle and cheatgrass are species likely to invade this site. Where this site occurs adjacent to juniper woodland communities these trees readily invade the site.

Fire Ecology:

Fire return intervals in basin big sagebrush range between 25 to 50 years. A naturally wide variation in fire frequency in this system is expected. Basin big sagebrush is readily killed when aboveground plant parts are charred by fire. Prolific seed production from nearby unburned plants coupled with high germination rates enables seedlings to establish rapidly following fire. Idaho fescue grows in a dense, fine-leaved tuft. Fires tend to burn within the accumulated fine leaves at the base of the plant and may produce temperatures sufficient to kill some of the root crown. Mature Idaho fescue plants are commonly reported to be severely damaged by fire in all seasons. Thurber's needlegrass is classified as moderately resistant, but depending on season of burn, phenology, and fire severity, this perennial bunchgrass is moderately to severely damaged by fire. Early season burning is more damaging to this needlegrass than late season burning. Burning bluebunch wheatgrass may remove most of the aboveground biomass but does not usually result in plant mortality. Bluebunch wheatgrass is generally favored by burning. Burning stimulates flowering and seed production. However, season of burning affects mortality. Basin wildrye is top-killed by fire. Older basin wildrye plants with large proportions of dead material within the perennial crown can be expected to show higher mortality due to fire than younger plants having little debris. Basin wildrye is generally tolerant of fire but may be damaged by early season fire combined with dry soil conditions. Canby's bluegrass is generally unharmed by fire. It produces little litter, and its small bunch size and sparse litter reduces the amount of heat transferred to perennating buds in the soil. Its rapid maturation in the spring also reduces fire damage, since it is dormant when most fires occur.

State and transition model

Additional community tables

Table 5. Community 1.1 plant community composition

Group	Common Name	Symbol	Scientific Name	Annual Production ()	Foliar Cover (%)
Grass/Grasslike					
1	Primary Perennial Grasses			668-1122	
	Idaho fescue	FEID	<i>Festuca idahoensis</i>	493-616	–
	Thurber's needlegrass	ACTH7	<i>Achnatherum thurberianum</i>	123-247	–
	basin wildrye	LECI4	<i>Leymus cinereus</i>	25-99	–
	bluebunch wheatgrass	PSSPS	<i>Pseudoroegneria spicata ssp. spicata</i>	25-99	–
2	Secondary Perennial Grasses			25-99	
	squirreltail	ELEL5	<i>Elymus elymoides</i>	7-25	–
	thickspike wheatgrass	ELLAL	<i>Elymus lanceolatus ssp. lanceolatus</i>	7-25	–
	needle and thread	HECO26	<i>Hesperostipa comata</i>	7-25	–
	prairie Junegrass	KOMA	<i>Koeleria macrantha</i>	7-25	–
	bluegrass	POA	<i>Poa</i>	7-25	–
Forb					
3	Perennial			62-185	
	tapertip hawksbeard	CRAC2	<i>Crepis acuminata</i>	7-37	–
	lupine	LUPIN	<i>Lupinus</i>	7-37	–
Shrub/Vine					
4	Primary Shrubs			185-308	
	basin big sagebrush	ARTRT	<i>Artemisia tridentata ssp. tridentata</i>	185-308	–
5	Secondary Shrubs			25-99	
	yellow rabbitbrush	CHVI8	<i>Chrysothamnus viscidiflorus</i>	12-37	–
	antelope bitterbrush	PUTR2	<i>Purshia tridentata</i>	12-37	–
	littleleaf horsebrush	TEGL	<i>Tetradymia glabrata</i>	12-37	–

Animal community

Livestock Interpretations: This site is suitable for livestock grazing. Grazing management should be keyed to perennial grass production. Idaho fescue provides important forage for many types of domestic livestock. The foliage cures well and is preferred by livestock in late fall and winter. Thurber's needlegrass species begin growth early in the year and remain green throughout a relatively long growing season. This pattern of development enables animals to use Thurber's needlegrass when many other grasses are unavailable. Cattle prefer Thurber's needlegrass in early spring before fruits have developed as it becomes less palatable when mature. Thurber's needlegrasses are grazed in the fall only if the fruits are softened by rain. Bluebunch wheatgrass is considered one of the most important forage grass species on western rangelands for livestock. Although bluebunch wheatgrass can be a crucial source of forage, it is not necessarily the most highly preferred species. The early growth and abundant production of basin wildrye make it a valuable source of forage for livestock. It is important forage for cattle and is readily grazed by cattle and horses in early spring and fall. Though coarse-textured during the winter, basin wildrye may be utilized more frequently by livestock and wildlife when snow has covered low shrubs and other grasses. Canby's bluegrass is a widespread forage grass. It is one of the earliest grasses in the spring and is sought by domestic livestock and several wildlife species. Canby's bluegrass is a palatable species, but its production is closely tied to weather conditions. It produces little forage in drought years, making it a less dependable food source than other perennial bunchgrasses. Basin big sagebrush may serve as emergency food during severe winter weather, but it is not usually sought out by livestock. Stocking rates vary over time depending upon season of use, climate variations, site, and previous and current management goals. A safe starting stocking rate is an estimated stocking rate that is fine tuned by the client by adaptive management through the year and from year to year. **Wildlife Interpretations:** Basin big sagebrush is the least palatable of all the subspecies of big sagebrush. Basin big sagebrush is browsed by mule deer from fall to early spring, but is not preferred. Sagebrush-grassland communities provide critical sage-grouse breeding and nesting habitats. Meadows surrounded by sagebrush may be used as feeding and strutting grounds. Sagebrush is a crucial component of their diet year-round, and sage-grouse select sagebrush almost exclusively for cover. Sage-grouse prefer mountain big sagebrush and Wyoming big sagebrush communities to basin big sagebrush communities. Idaho fescue provides important forage for several wildlife species. It is reported to be good forage for pronghorn, and deer in ranges of northern Nevada. Thurber needlegrass is valuable forage for wildlife. Bluebunch wheatgrass is considered one of the most important forage grass species on western rangelands for wildlife. Bluebunch wheatgrass does not generally provide sufficient cover for ungulates, however, mule deer are frequently found in bluebunch-dominated grasslands. Basin wildrye provides winter forage for mule deer, though use is often low compared to other native grasses. Basin wildrye provides summer forage for black-tailed jackrabbits. Because basin wildrye remains green throughout early summer, it remains available for small mammal forage for longer time than other grasses. Canby's bluegrass is desirable for pronghorn antelope and mule deer in the spring and preferable in the spring, summer, and fall for elk and desirable as part of their winter range.

Hydrological functions

Runoff is very low. Permeability is moderately slow to moderately rapid. Hydrologic soil group is A, B, and C.

Recreational uses

Aesthetic value is derived from the diverse floral and faunal composition and the colorful flowering of wild flowers and shrubs during the spring and early summer. This site offers rewarding opportunities to photographers and for nature study. This site is used for camping and hiking and has potential for upland and big game hunting.

Other products

Some Native American peoples used the bark of big sagebrush to make rope and baskets. Basin wildrye was used as bedding for various Native American ceremonies, providing a cool place for dancers to stand.

Other information

Basin big sagebrush shows high potential for range restoration and soil stabilization. Basin big sagebrush grows rapidly and spreads readily from seed. Basin wildrye is useful in mine reclamation, fire rehabilitation and stabilizing disturbed areas. Its usefulness in range seeding, however, may be limited by initially weak stand establishment.

Type locality

Location 1: Washoe County, NV	
Township/Range/Section	T43N R23E S33
UTM zone	N

UTM northing	300699
UTM easting	4607951
Latitude	41° 35' 54"
Longitude	119° 23' 29"
General legal description	NW 1/4 SW 1/4, Along road to Wall Canyon south of County Road 8A, USF&WS Sheldon Refuge, Washoe County, Nevada. This site also occurs in Humboldt County, Nevada.

Other references

Fire Effects Information System (Online; <http://www.fs.fed.us/database/feis/plants/>).

USDA-NRCS Plants Database (Online; <http://www.plants.usda.gov>).

Contributors

GKB
T Stringham (UNR under contract with BLM)

Approval

Kendra Moseley, 4/10/2025

Rangeland health reference sheet

Interpreting Indicators of Rangeland Health is a qualitative assessment protocol used to determine ecosystem condition based on benchmark characteristics described in the Reference Sheet. A suite of 17 (or more) indicators are typically considered in an assessment. The ecological site(s) representative of an assessment location must be known prior to applying the protocol and must be verified based on soils and climate. Current plant community cannot be used to identify the ecological site.

Author(s)/participant(s)	
Contact for lead author	
Date	04/20/2026
Approved by	
Approval date	

Composition (Indicators 10 and 12) based on	Annual Production
---	-------------------

Indicators

1. Number and extent of rills:

2. Presence of water flow patterns:

3. Number and height of erosional pedestals or terracettes:

4. Bare ground from Ecological Site Description or other studies (rock, litter, lichen, moss, plant canopy are not bare ground):

5. Number of gullies and erosion associated with gullies:

6. Extent of wind scoured, blowouts and/or depositional areas:

7. Amount of litter movement (describe size and distance expected to travel):

8. Soil surface (top few mm) resistance to erosion (stability values are averages - most sites will show a range of values):

9. Soil surface structure and SOM content (include type of structure and A-horizon color and thickness):

10. Effect of community phase composition (relative proportion of different functional groups) and spatial distribution on infiltration and runoff:

11. Presence and thickness of compaction layer (usually none; describe soil profile features which may be mistaken for compaction on this site):

12. Functional/Structural Groups (list in order of descending dominance by above-ground annual-production or live foliar cover using symbols: >>, >, = to indicate much greater than, greater than, and equal to):

Dominant:

Sub-dominant:

Other:

Additional:

13. Amount of plant mortality and decadence (include which functional groups are expected to show mortality or decadence):

14. Average percent litter cover (%) and depth (in):

15. Expected annual annual-production (this is TOTAL above-ground annual-production, not just forage annual-production):

16. Potential invasive (including noxious) species (native and non-native). List species which BOTH characterize degraded states and have the potential to become a dominant or co-dominant species on the ecological site if their future establishment and growth is not actively controlled by management interventions. Species that become dominant for only one to several years (e.g., short-term response to drought or wildfire) are not invasive plants. Note that unlike other indicators, we are describing what is NOT expected in the reference state for the ecological site:

17. Perennial plant reproductive capability:
