

# Ecological site R024XY004NV

## SILTY 4-8 P.Z.

Last updated: 3/07/2025  
 Accessed: 07/03/2026

### General information

**Provisional.** A provisional ecological site description has undergone quality control and quality assurance review. It contains a working state and transition model and enough information to identify the ecological site.

### MLRA notes

Major Land Resource Area (MLRA): 024X–Humboldt Basin and Range Area

Major Land Resource Area (MLRA): 024X–Humboldt Basin and Range Area Major land resource area (MLRA) 24, the Humboldt Area, covers an area of approximately 8,115,200 acres (12,680 sq. mi.). MLRA 24 is in the Great Basin Section of the Basin and Range Province of the Intermontane Plateaus. Elevations predominantly range from 3,950 to 5,900 feet (1,205 to 1,800 meters). The elevations of some mountain peaks are more than 8,850 feet (2,700 meters). A series of widely spaced north-south trending mountain ranges are separated by broad valleys filled with alluvium washed in from adjacent mountain ranges. Most valleys are drained by tributaries to the Humboldt River. Playas, however, are in lower elevation valleys with closed drainage systems. Isolated ranges are dissected, uplifted fault-block mountains. Geology is comprised of Mesozoic and Paleozoic volcanic rock and marine and continental sediments. Young andesite and basalt flows (6 to 17 million years old) are at the margins of the mountains. Dominant soil orders include Aridisols, Entisols, Inceptisols and Mollisols. Soils of the area are generally characterized by a mesic soil temperature regime, an aridic soil moisture regime and mixed geology. They are generally well drained, loamy and very deep. 75 percent of MLRA 24 is federally owned. The remainder is primarily used for farming, ranching and mining. Irrigated land comprises 3 percent of the area; most of the irrigation water is from surface water sources, such as the Humboldt River and Rye Patch Reservoir. Annual precipitation typically ranges from 6 to 12 inches (15 to 30 cm) for most of the area. In the mountains however the precipitation may be up to 40 inches (101 cm). Most of the annual precipitation is from snow in the winter. In the spring and fall, rainfall occurs as high- intensity, convective thunderstorms. Nevada is on the eastern, lee side of the Sierra Nevada Range; a massive mountain barrier that markedly influences the climate of the State. The prevailing winds are from the west. The warm moist air from the Pacific Ocean ascends the western slopes of the Sierra Range, the air cools, condenses and the moisture falls as precipitation. As the air descends the eastern slope, it is warmed by compression, and very little precipitation occurs. The effects of this mountain barrier are felt not only in the west but throughout the State. The result is the lowlands of Nevada are largely desert or steppes.

### Ecological site concept

This ecological site is on fan piedmonts. Soils associated with this site are very deep, well drained and formed in alluvium derived from mixed rocks, loess and volcanic ash. The soil profile is characterized by an ochric epipedon, a sodium free surface, and moderately to strongly sodium effected subsoil. Soil textures are dominated by silt loam, ashy very fine silt loam, and/or ashy fine sandy loam. The soil temperature regime is mesic and the soil moisture regime is typic aridic. The reference state is dominated by winterfat, bud sagebrush and Indian ricegrass. Future field work in compare the soil characteristics and abiotic factors for all winterfat dominated ESCs (024XY004NV, 024XY011NV, 024XY014NV, 024XY059NV & 024XY011OR) in MRLA 24 and determine if they are one ESC.

### Associated sites

<b>R024XY014NV</b>	<p><b>COARSE SILTY 4-8 P.Z.</b></p> <p>This ecological site occurs on fan piedmonts. Soils associated with this site are very deep, well drained and formed in alluvium derived from mixed rocks, loess and volcanic ash. The soil profile is characterized by an ochric epipedon. The reference state is dominated by winterfat (KRLA2), bud sagebrush (ARSP5), and Indian ricegrass (ACHY).</p>
--------------------	---

<b>R024XY020NV</b>	<p><b>DROUGHTY LOAM 8-10 P.Z.</b></p> <p>Important abiotic factors contributing to the presence of this site include limited available soil moisture due to texture and precipitation zone. Vegetative cover is less than 25% and is dominated by deep-rooted, cool season perennial bunchgrasses and drought tolerant shrubs. Dominant species include Thurber's needlegrass (ACTH7), Indian ricegrass (ACHY), Wyoming big sagebrush (ARTRW8), and spiny hopsage (GRSP).</p>
--------------------	---

**Similar sites**

<b>R024XY059NV</b>	<p><b>SILTY 8-10 P.Z.</b></p> <p>The soil profile is characterized by an ochric epipedon, a sodium free surface, and moderately to strongly sodium effected subsoil. More productive site Winterfat (KRLA2), Indian ricegrass (ACHY); does not occur on bolson floor.</p>
<b>R024XY060NV</b>	<p><b>SHALLOW SILTY 8-10 P.Z.</b></p> <p>Soils are dominated by fine loams and find sands and have a strong vesicular horizon and very platy structure in the surface horizon. Shadscale saltbrush (ATCO) dominant plant; Bud sagebrush (ARSP5) &amp; Greasewood (SAVE4) rare.</p>
<b>R024XY011NV</b>	<p><b>SODIC FLAT 6-8 P.Z.</b></p> <p>Important abiotic factors include crusting &amp; baking of the surface layer upon drying, inhibiting water infiltration and seedling emergence. Greasewood (SAVE4) dominant shrub; Shadscale saltbrush (ATCO) minor shrub.</p>
<b>R024XY014NV</b>	<p><b>COARSE SILTY 4-8 P.Z.</b></p> <p>The soil profile is characterized by an ochric epipedon, a sodium free surface, and moderately to strongly sodium effected subsoil. More productive site Winterfat (KRLA2), Bud sagebrush (ARSP5); greater shrub diversity.</p>
<b>R024XY067NV</b>	<p><b>SHALLOW SILTY 5-8 P.Z.</b></p> <p>The soil profile is characterized by an ochric epipedon, continuous weak cementation with in 20 inches, and coarse silty particle size control section. Shadscale saltbrush (ATCO) dominant shrub; less productive site.</p>
<b>R024XY022NV</b>	<p><b>SODIC TERRACE 8-10 P.Z.</b></p> <p>Soils are characterized by a very low infiltration, an ochric epipedon, moderate to very strong alkalinity. Greasewood (SAVE4), Big sagebrush (ARTR2) codominant; more productive site.</p>

**Table 1. Dominant plant species**

Tree	Not specified
Shrub	(1) <i>Krascheninnikovia lanata</i>
Herbaceous	(1) <i>Achnatherum hymenoides</i>

**Physiographic features**

This site is on fan piedmonts. Slopes range from 0 to 15 percent, but slope gradients of 2 to 8 are most typical. Elevations are 3900 to 5600 feet (1189 to 1707 meters).

**Table 2. Representative physiographic features**

Landforms	(1) Fan skirt (2) Lake plain (3) Inset fan
Runoff class	Low
Flooding duration	Very brief (4 to 48 hours)
Flooding frequency	Rare
Ponding frequency	None
Elevation	1,190 – 1,710 m
Slope	0 – 20 %
Water table depth	180 cm
Aspect	Aspect is not a significant factor

### **Climatic features**

The climate associated with this site is semiarid and characterized by cool, moist winters and warm, dry summers. Average annual precipitation is 4 to 8 inches (10 to 20 cm). Mean annual air temperature is 45 to 53 degrees F.

**Table 3 Representative climatic features**

Frost-free period (characteristic range)	90-90 days
Freeze-free period (characteristic range)	100-110 days
Precipitation total (characteristic range)	200-230 mm
Frost-free period (actual range)	80-90 days
Freeze-free period (actual range)	100-110 days

Precipitation total (actual range)	180-230 mm
Frost-free period (average)	90 days
Freeze-free period (average)	110 days
Precipitation total (average)	200 mm

- (1) BEOVAWE [USC00260795], Crescent Valley, NV
- (2) GOLCONDA [USC00263245], Golconda, NV

### Influencing water features

Influencing water features are not associated with this ecological site.

### Soil features

The soils associated with this site are very deep, well drained and formed in alluvium derived from mixed rocks, loess, and volcanic ash. The soil profile is characterized by an ochric epipedon, a cambic horizon, and silt loam textures throughout. The soil profile is moderately to strongly alkaline at depth, but its not salt affected at the surface.

Permeability is moderately slow and available water capacity is high. The soil temperature regime is mesic and the soil moisture regime is typic aridic.

The typical soil series associated with this site is Wholan, a coarse-silty, mixed, superactive, mesic Typic Haplocambid. Other soil series associated with this site include: Batan, Broyles, Defler, Dun Glen, Kumiva, Nomazu and Smaug.

**Table 4. Representative soil features**

Parent material	(1) Alluvium (2) Loess (3) Volcanic ash
Surface texture	(1) Very fine sandy loam (2) Sandy loam
Family particle size	(1) Loamy
Drainage class	Well drained
Permeability class	Moderately slow
Soil depth	180 – 210 cm

Surface fragment cover <=3"	0 – 10 %
Surface fragment cover >3"	Not specified
Available water capacity (0-101.6cm)	15.24 – 20.32 cm
Calcium carbonate equivalent (0-101.6cm)	0 – 20 %
Electrical conductivity (0-101.6cm)	0 – 20 mmhos/cm
Sodium adsorption ratio (0-101.6cm)	0 – 10
Soil reaction (1:1 water) (0-101.6cm)	7.4 – 9.6
Subsurface fragment volume <=3" (Depth not specified)	0 – 10 %
Subsurface fragment volume >3" (Depth not specified)	Not specified

## Ecological dynamics

An ecological site is the product of all the environmental factors responsible for its development and it has a set of key characteristics that influence a site's resilience to disturbance and resistance to invasives. Key characteristics include 1) climate (precipitation, temperature), 2) topography (aspect, slope, elevation, and landform), 3) hydrology (infiltration, runoff), 4) soils (depth, texture, structure, organic matter), 5) plant communities (functional groups, productivity), and 6) natural disturbance regime (fire, herbivory, etc.) (Caudle 2013). Biotic factors that influence resilience include site productivity, species composition and structure, and population regulation and regeneration (Chambers et al. 2013).

The plant communities of this site are dynamic in response to weather events and disturbance regimes. The reference plant community is dominated by winterfat with an herbaceous understory of Indian ricegrass and bottlebrush squirreltail. Bud sagebrush, needleandthread and Sandberg's bluegrass are also common species present in the reference plant community. This site inherently has low resistance to invasion and low resilience following invasion by non-native species. In Great Basin ecosystems, inherent resilience typically increases with elevation due to higher levels of water, nutrients and biomass production (Chambers et al. 2012). Management activities should be prioritized based on relative resilience and resistance of a specific ecological site.

Winterfat is a long-lived, drought tolerant, native shrub typically about 30 cm tall (Mozingo 1987). It has a woody base from which annual branchlets grow (Welsh et al. 1987). The most common variety is a low growing dwarf form (less than 38.1 cm), which is most commonly on desert valley floors (Stevens et al. 1977). Total winter precipitation is a primary growth driver and lower than average spring

precipitation can reverse the impact of plentiful winter precipitation. While summer rainfall has a limited impact, heavy August-September rain can cause a second flowering in winterfat (West and Gasto 1978). Winterfat reproduces from seed and primarily pollinates via wind (Stevens et al. 1977). Seed production, especially in desert regions, is dependent on precipitation (West and Gasto 1978). Seeds are abundant when summer precipitation is appreciable and browsing is minimal (Stevens et al. 1977). Winterfat has multiple dispersal mechanisms: diaspores are shed in the fall or winter, dispersed by wind, rodent-cached, or carried on animals (Majerus 2003). Diaspores take advantage of available moisture, tolerating freezing conditions as they progress from imbibed seeds to germinants to nonwoody seedlings (Booth 1989). Under some circumstances, the degree of reproduction may be dependent on mature plant density (Freeman and Emlen 1995).

Winterfat tolerates environmental stress, extremes of temperature and precipitation, and competition from other perennials but not the disturbance of fire or overgrazing (Ogle 2001). Winterfat typically germinates in late winter and early spring. Winterfat is cold tolerant and even able to germinate under high moisture stress conditions if temperatures are consistently around 40°F (Woodmansee and Potter 1971). Early germination allows seedlings to become established before other plants, especially grasses begin growing and using large quantities of soil moisture (Woodmansee and Potter 1971). Root growth of winterfat seedlings begins before above ground morphological development appear (West and Gasto 1978).

Bud sagebrush, a minor shrub on this ecological site, is a native, summer-deciduous shrub. It is a low growing, spinescent, aromatic shrub with a height of 4 to 10 inches (10 to 25 cm) and a spread of 8 to 12 inches (20 to 30 cm) (Chambers and Norton 1993).

Indian ricegrass, the dominant grass, is a hardy, cool-season, densely tufted, native perennial bunchgrass that grows from 4 to 24 inches (10 to 61 cm) in height (Blaisdell and Holmgren 1984). Indian ricegrass has deep, fibrous, extensive roots and is decidedly drought tolerant once established. It reproduces from seed. Years of good seedling establishment are infrequent, although typically observed during years with wet springs (Tirmenstein 1999).

Bottlebrush squirreltail, another cool-season, native perennial bunchgrass is common to this ecological site. Bottlebrush squirreltail has the ability to produce large numbers of highly germinable seeds, with relatively rapid germination (Young and Evans 1977) when exposed to the correct environmental cues. Early spring growth and ability to grow at low temperatures contribute to the persistence of bottlebrush squirreltail among cheatgrass dominated ranges (Hironaka and Tisdale 1972). Squirreltail generally increases in abundance when moderately grazed or protected (Hutchings and Stewart 1953). In addition, moderate trampling by livestock in big sagebrush rangelands of central Nevada enhanced bottlebrush squirreltail seedling emergence compared to untrampled conditions. Heavy trampling however significantly reduced germination sites (Eckert et al. 1987). Squirreltail is more tolerant of grazing than Indian ricegrass but all bunchgrasses are sensitive to over-utilization within the growing season.

In summary, prolonged drought and/or abusive grazing will cause a decrease in winterfat, bud sagebrush, and Indian ricegrass, while bare ground increases. Squirreltail may maintain or also decline within the community. Repeated spring and early summer grazing will have an especially detrimental effect on winterfat and bud sagebrush. Cheatgrass and other non-native annual invasive species increase with excessive grazing. Abusive grazing during the winter may lead to soil compaction and reduced infiltration. Prolonged abusive grazing during any season leads to abundant bare ground, desert pavement and active wind and water erosion. Repeated, frequent fire will promote cheatgrass dominance and elimination of the native plant community.

These communities commonly exhibit the formation of microbiotic crusts within the interspaces between shrubs. These crusts influence the soils on these sites and their ability to reduce erosion and increase infiltration; they may also alter the soil structure and possibly increase soil fertility (Fletcher and Martin 1948, Williams 1993). Finer textured soils such as silts tend to support more microbiotic cover than coarse texture soils (Anderson 1982). Disturbance such as hoof action from inappropriate grazing and cheatgrass (*Bromus tectorum*) invasion can reduce biotic crust integrity (Anderson 1982, Ponzetti et al. 2007) and increase erosion.

Drought and/or inappropriate grazing will initially favor shrubs but prolonged drought can cause a decrease in the winterfat, bud sagebrush and other shrubs, while bare ground increases. Indian ricegrass will decrease with inappropriate grazing management. Squirreltail may maintain or also decline within the community. Repeated spring and early summer grazing will have an especially detrimental effect on winterfat and bud sagebrush (*Atrémisia tridentata*). Cheatgrass and other non-native annual weeds increase with excessive grazing. Abusive grazing during the winter may lead to soil compaction and reduced infiltration. Prolonged abusive grazing during any season leads to abundant bare ground, desert pavement and active wind and water erosion. Repeated, frequent fire will promote cheatgrass dominance and elimination of the native plant community. These sites attract recreational use, primarily by off highway vehicles (OHV). Annual non-native species increase where surface soils have been disturbed. Three alternative stable states have been identified for this site.

#### Fire Ecology:

Winterfat tolerates environmental stress, extremes of temperature and precipitation, and competition from other perennials but not the disturbance of fire or overgrazing (Ogle 2001). Fire is rare within these communities due to low fuel loads. Conflicting reports are in the literature about the response of winterfat to fire. In one of the first published descriptions, Dwyer and Pieper (1967) reported that winterfat sprouts vigorously after fire. This observation was frequently cited in subsequent literature, but recent observations have suggested that winterfat can be completely killed by fire (Pellant and Reichert 1984). The response is apparently dependent on fire severity. Winterfat is able to sprout from buds near the base of the plant. However, if these buds are destroyed, winterfat will not sprout. Research has shown that winterfat seedling growth is depressed in growth by at least 90 percent when growing in the presence of cheatgrass (Hild et al. 2007). Repeated,

frequent fires will increase the likelihood of conversion to a non-native, annual plant community with trace amounts of winterfat.

Bud sagebrush (*Picrothamnus desertorum*), a minor shrub to this ecological site, is a native, summer-deciduous shrub. It is low growing, spinescent, aromatic shrub with a height of 4 to 10 inches (10 to 25 cm) and a spread of 8 to 12 inches (20 to 30 cm) (Chambers and Norton 1993). Bud sagebrush is fire intolerant and must reestablish from seed (Banner 1992, West 1994).

Indian ricegrass, the dominant grass within this site, is a hardy, cool-season, densely tufted, native perennial bunchgrass that grows from 4 to 24 inches (10 to 61 cm) in height (Blaisdell and Holmgren 1984). Indian ricegrass may reestablish on burned sites through seed dispersed from adjacent unburned areas (Young 1983). Thus the presence of surviving, seed producing plants is necessary for reestablishment of Indian ricegrass. Grazing management following fire to promote seed production and establishment of seedlings is important.

Bottlebrush squirreltail, another cool-season, native perennial bunchgrass is common to this ecological site. Bottlebrush squirreltail is considered more fire tolerant than Indian ricegrass due to its small size, coarse stems, and sparse leafy material (Britton et al. 1990). Postfire regeneration happens because of surviving root crowns and from seed sources both onsite and offsite. Bottlebrush squirreltail can produce large numbers of highly germinable seeds, with relatively rapid germination (Young and Evans 1977) when exposed to the correct environmental cues. Early spring growth and ability to grow at low temperatures contribute to the persistence of bottlebrush squirreltail among cheatgrass dominated ranges (Hironaka and Tisdale 1972).

Conflicting reports are in the literature about the response of winterfat to fire. In one of the first published descriptions, Dwyer and Pieper (1967) reported that winterfat sprouts vigorously after fire. This observation was frequently cited in subsequent literature, but recent observations have suggested that winterfat can be completely killed by fire (Pellant and Reichert 1984). The response is apparently dependent on fire severity. Winterfat is able to sprout from buds near the base of the plant. However, if these buds are destroyed, winterfat will not sprout. Research has shown that winterfat seedling growth is depressed in growth by at least 90 percent when growing in the presence of cheatgrass (Hild et al. 2007). Repeated, frequent fires will increase the likelihood of conversion to a non-native, annual plant community with trace amounts of winterfat.

Bud sagebrush is fire intolerant and must reestablish from seed (Banner 1992, West 1994). Indian ricegrass can be killed by fire, depending on severity and season of burn. Indian ricegrass reestablishes on burned sites through seed dispersed from adjacent unburned areas.

Bottlebrush squirreltail is considered more fire tolerant than Indian ricegrass due to its small size, coarse stems, and sparse leafy material (Britton et al. 1990). Postfire regeneration is from surviving root crowns and seed sources both onsite and offsite. Frequency of disturbance greatly influences postfire response of bottlebrush squirreltail. Undisturbed plants within a six-to-nine-year age class generally contain large amounts of dead material, increasing bottlebrush squirreltail's susceptibility to fire.

## State and transition model

Figure 7. T Stringham 4/2016

Figure 8. Legend

## Additional community tables

Table 5. Community 1.1 plant community composition

Group	Common Name	Symbol	Scientific Name	Annual Production ()	Foliar Cover (%)
<b>Grass/Grasslike</b>					
1	<b>Primary Perennial Grasses</b>			67-130	
	Indian ricegrass	ACHY	<i>Achnatherum hymenoides</i>	59-99	–
	squirreltail	ELEL5	<i>Elymus elymoides</i>	8-31	–
2	<b>Secondary Perennial Grasses</b>			8-31	
	needle and thread	HECO26	<i>Hesperostipa comata</i>	2-12	–
	western wheatgrass	PASM	<i>Pascopyrum smithii</i>	2-12	–
	Sandberg bluegrass	POSE	<i>Poa secunda</i>	2-12	–
<b>Forb</b>					
3	<b>Perennial Forbs</b>			8-31	
	globemallow	SPHAE	<i>Sphaeralcea</i>	2-8	–
4	<b>Annual Forbs</b>			1-20	
<b>Shrub/Vine</b>					
5	<b>Primary Shrubs</b>			84-183	

	winterfat	KRLA2	<i>Krascheninnikovia lanata</i>	76-151	-
6	<b>Secondary Shrubs</b>			8-39	
	fourwing saltbush	ATCA2	<i>Atriplex canescens</i>	4-8	-
	shadscale saltbush	ATCO	<i>Atriplex confertifolia</i>	4-8	-
	sickle saltbush	ATFA	<i>Atriplex falcata</i>	4-8	-
	yellow rabbitbrush	CHV18	<i>Chrysothamnus viscidiflorus</i>	4-8	-
	spiny hopsage	GRSP	<i>Grayia spinosa</i>	4-8	-

Table 6. Community 1.2 plant community composition

Group	Common Name	Symbol	Scientific Name	Annual Production ()	Foliar Cover (%)
-------	-------------	--------	-----------------	----------------------	------------------

Table 7. Community 2.1 plant community composition

Group	Common Name	Symbol	Scientific Name	Annual Production ()	Foliar Cover (%)
-------	-------------	--------	-----------------	----------------------	------------------

Table 8. Community 2.2 plant community composition

Group	Common Name	Symbol	Scientific Name	Annual Production ()	Foliar Cover (%)
-------	-------------	--------	-----------------	----------------------	------------------

Table 9. Community 3.1 plant community composition

Group	Common Name	Symbol	Scientific Name	Annual Production ()	Foliar Cover (%)
-------	-------------	--------	-----------------	----------------------	------------------

Table 10. Community 4.1 plant community composition

Group	Common Name	Symbol	Scientific Name	Annual Production ()	Foliar Cover (%)
-------	-------------	--------	-----------------	----------------------	------------------

Table 11. Community 4.2 plant community composition

Group	Common Name	Symbol	Scientific Name	Annual Production ()	Foliar Cover (%)
-------	-------------	--------	-----------------	----------------------	------------------

## Animal community

Livestock Interpretations: This site has value for livestock grazing. Grazing management should be keyed to dominant grasses or palatable shrub production. Winterfat is a valuable forage species with an average of 10 percent crude protein during winter when few nutritious options exist for livestock and wildlife (Welch 1989). However, excessive grazing throughout the West has negatively impacted survival of winterfat stands (Hilton 1941; Statler 1967; Stevens et al. 1977). Time of grazing is critical for winterfat with the active growing period being most critical (Romo 1995). Stevens et al. (1977) observed that both vigor and reproduction of winterfat were reduced in Steptoe Valley, Nevada by improper season of use, and recommended no more than 25 percent utilization during periods of active growth and up to 75 percent utilization during dormant season use. Rasmussen and Brotherson (1986) observed significantly greater foliar cover and density of winterfat in areas ungrazed for 26 years versus winter grazed areas in Utah. In exclosures protected from grazing, Rice and Westoby (1978) observed that winterfat increased in foliar cover but not in density where it was dominant, and in both foliar cover and density in shadscale-perennial grass communities where it was not dominant. Bud sagebrush is also a palatable, nutritious forage for upland game birds, small game, big game and domestic sheep in winter, particularly late winter (Johnson 1978), however it can be poisonous or fatal to calves when eaten in quantity (Stubbendieck et al. 1992). Budsage is highly susceptible to effects of browsing. It decreases under browsing due to year-long palatability of its buds and is particularly susceptible to browsing in the spring when it is physiologically most active (Chambers and Norton 1993). Heavy browsing (>50 percent) may kill budsage rapidly (Wood and Brotherson 1986). Grazing management following fire to promote Indian ricegrass seed production and establishment of seedlings is important. Heavy spring grazing reduced the vigor of Indian ricegrass and decrease the stand (Cook and Child 1971). In eastern Idaho, productivity of Indian ricegrass was at least 10 times greater in undisturbed plots than in heavily grazed ones (Pearson 1976). Cook and Child (1971) observed significant reduction in plant cover after 7 years of rest from heavy (90 percent) and moderate (60 percent) spring use. The seed crop may be reduced where grazing is heavy (Bich et al. 1995). Tolerance to grazing increases after May thus spring deferment may be necessary for stand enhancement (Pearson 1964, Cook and Child 1971); however, utilization of less than 60 percent is recommended. Bottlebrush squirreltail is very palatable winter forage for domestic sheep of Intermountain ranges. Overall, bottlebrush squirreltail is considered moderately palatable to livestock. Stocking rates vary over time depending upon season of use, climate variations, site, and previous and current management goals. A safe starting stocking rate is an estimated stocking rate that is fine tuned by the client by adaptive management through the year and from year to year. Wildlife Interpretations: In addition to grazing by cattle, winterfat is browsed by rabbits, antelope, and other wildlife species (Stevens et al. 1977, Ogle et al. 2001). Winterfat and perennial grasses average 80 percent of jackrabbits' diet in southeastern Idaho, with shrubs being grazed in fall and winter particularly (Johnson and Anderson 1984). Pronghorn and rabbits browse stems, leaves, and seed stalks of winterfat year round, especially during periods of active growth (Stevens et al. 1977). Management of wildlife browse is difficult and browse may be harmful to winterfat reestablishment as seed production and regrowth are curtailed if the plant is grazed at the onset of growth (Eckert 1954). Winterfat is used for cover by

rodents. It is potential nesting cover for upland game birds, especially when grasses grow up through its crown. Budsage is palatable, nutritious forage for upland game birds, small game and big game in winter. Budsage is browsed by mule deer in Nevada in winter and is utilized by bighorn sheep in summer, but the importance of budsage in the diet of bighorns is not known. Bud sage comprises 18 to 35 percent of a pronghorn's diet during the spring where it is available. Chukar will utilize the leaves and seeds of bud sage. Budsage is highly susceptible to effects of browsing. It decreases under browsing due to year-long palatability of its buds and is particularly susceptible to browsing in the spring when it is physiologically most active. Indian ricegrass is eaten by pronghorn in moderate amounts whenever available. In Nevada, it is consumed by desert bighorns. A number of heteromyid rodents inhabiting desert rangelands show preference for seed of Indian ricegrass. Indian ricegrass is an important component of jackrabbit diets in spring and summer. In Nevada, Indian ricegrass may even dominate jackrabbit diets during the spring through early summer months. Indian ricegrass seed provides food for many species of birds. Doves, for example, eat large amounts of shattered Indian ricegrass seed lying on the ground. Bottlebrush squirreltail is a dietary component of several wildlife species. Bottlebrush squirreltail may provide forage for mule deer and pronghorn.

### Hydrological functions

Runoff is low to medium. Permeability is moderately slow to moderately rapid. Hydrologic soil groups are A and B. Rills are none. Water flow patterns are rare to common depending on site location relative to major inflow areas. Pedestals are none. Gullies are none. Shrubs and deep-rooted perennial herbaceous bunchgrasses and/or rhizomatous grasses aid in infiltration. Shrub canopy and associated litter break raindrop impact and provide opportunity for snow catch and accumulation on site.

### Recreational uses

Aesthetic value is derived from the diverse floral and faunal composition and the colorful flowering of wild flowers and shrubs during the spring and early summer. This site offers rewarding opportunities to photographers and for nature study. This site has potential for upland bird and big game hunting.

### Other products

Indian ricegrass is traditionally eaten by some Native Americans. The Paiutes use seeds as a reserve food source.

### Other information

Winterfat adapts well to most site conditions, and its extensive root system stabilizes soil. However, winterfat is intolerant of flooding, excess water, and acidic soils. Bottlebrush squirreltail is tolerant of disturbance and is a suitable species for revegetation.

### Inventory data references

NASIS soil mapunit data was used for correlated soil components.

### Type locality

Location 1: Pershing County, NV	
Township/Range/Section	T33N R35E S26
UTM zone	N
UTM northing	4506603
UTM easting	414456
Latitude	40° 42' 21"
Longitude	118° 0' 45"

General legal description	SE¼NW¼ About 2 miles southeast of Dun Glen-Mill City exit off I-80 west of Winnemucca, Dun Glen Flat area, Pershing County, Nevada.  This site also occurs in Elko, Eureka, Humboldt, Lander, and Washoe Counties, Nevada.
---------------------------	--

### Other references

Banner, R.E. 1992. Vegetation types of Utah. *Journal of Range Management* 14(2):109-114.

Bich, B.S., J.L. Butler, and C.A. Schmidt. 1995. Effects of differential livestock use of key plant species and rodent populations within selected *Oryzopsis hymenoides/Hilaria jamesii* communities in Glen Canyon National Recreation Area. *The Southwestern Naturalist* 40(3):281-287.

Blaisdell, J.P. and R.C. Holmgren. 1984. Managing Intermountain Rangelands – Salt-desert shrub ranges. USDA-FS General Technical Report INT-163. 52 p.

Booth, D.T. 1989. A model of freeze tolerance in winterfat germinants. In: *Proceedings--Symposium on shrub ecophysiology and biotechnology*; 1987 June 30-July 2; Logan, UT. Gen. Tech. Rep. INT-256. Ogden, UT. U.S. Department of Agriculture, Forest Service, Intermountain Research Station: Pgs 83-89.

Britton, C.M., G.R. McPherson, and F.A. Sneva. 1990. Effects of burning and clipping on five bunchgrasses in eastern Oregon. *The Great Basin Naturalist* 50(2):115-120.

Chambers, J.C. and B.E. Norton. 1993. Effects of grazing and drought on population dynamics of salt desert species on the Desert Experimental Range, Utah. *Journal of Arid Environments* 24:261-275.

Clark, L.D. and N.E. West. 1971. Further studies of *Eurotia lanata* germination in relation to salinity. *The Southwestern Naturalist* 15(3):371-375.

Cook, C.W. and R.D. Child. 1971. Recovery of desert plants in various states of vigor. *Journal of Range Management* 24(5):339-343.

Dwyer, D.D. and R.D. Pieper. 1967. Fire effects on blue grama--pinyon-juniper rangeland in New Mexico. *Journal of Range Management* 20:359-362.

Eckert, R.E., Jr. 1954. A study of competition between whitesage and halogeton in Nevada. *Journal of Range Management* 7:223-225.

Eckert, R.E., Jr., F.F. Peterson, and F.L. Emmerich. 1987. A study of factors influencing secondary succession in the sagebrush [*Artemisia* spp. L.] type. In: Frasier, G.W. and R.A. Evans, (eds.). *Proceedings of the symposium: "Seed and seedbed ecology of rangeland plants"*; 1987 April 21-23; Tucson, AZ. Washington, DC: U.S. Department of Agriculture, Agricultural Research Service: Pgs 149-168.

Freeman, D.C. and J.M. Emlen. 1995. Assessment of interspecific interactions in plant communities: an illustration from the cold desert saltbush grasslands of North America. *Journal of Arid Environments* 31:179-198.

Hild, A.L., J.M. Muscha, and N.L. Shaw. 2007. Emergence and growth of four winterfat accessions in the presence of the exotic annual cheatgrass. In: Sosebee, R.E., D.B. Wester, C.M. Britton, E.D. McArthur, and S.G. Kitchen (compilers). *Proceedings: Shrubland dynamics—fire and water*; 2004 Aug 10-12; Lubbock, TX. RMRS-P-47. Fort Collins, CO: U.S. Department of Agriculture, Forest Service, Rocky Mountain Research Station: Pgs 147-152.

Hilton, J.W. 1941. Effects of certain micro-ecological factors on the germinability and early development of *Eurotia lanata*. *Northwest Science* 15:86-92.

Hironaka, M. and E.W. Tisdale. 1972. Growth and development of *Sitanion hystrix* and *Poa sandbergii*. Research Memorandum RM 72-24. U.S. International Biological Program, Desert Biome. 15 p.

Hutchings, S.S. and G. Stewart. 1953. Increasing forage yields and sheep production on Intermountain winter ranges. Circular No. 925. Washington, DC: U.S. Department of Agriculture. 63 p.

Johnson, K.L. 1978. Wyoming shrublands: Proceedings, 7th Wyoming shrub ecology workshop; 1978 May 31-June 1; Rock Springs, WY. Laramie, WY: University of Wyoming, Agricultural Extension Service. 58 p.

- Johnson, R.D. and J.E. Anderson. 1984. Diets of black-tailed jackrabbits in relation to population density and vegetation. *Journal of Range Management* 37:79-83.
- Majerus, M. 2003. Production and conditioning of winterfat seeds (*Krascheninnikovia lanata*). *Native Plants Journal* 4(1):10-17.
- Mozingo, H.N. 1987. *Shrubs of the Great Basin: A Natural History*. University of Nevada Press, Reno, NV. 342 p.
- Ogle, D.G., L. St. John, and L. Holzworth. 2001. Plant guide management and use of winterfat. Boise (ID): USDA-NRCS. 4 p.
- Pearson, L.C. 1964. Effect of harvest date on recovery of range grasses and shrubs. *Agronomy Journal* 56:80-82.
- Pearson, L.C. 1976. Primary production in grazed and ungrazed desert communities of eastern Idaho. *Ecology* 46(3):278-285.
- Pellant, M. and L. Reichert. 1984. Management and rehabilitation of a burned winterfat community in southwestern Idaho. In: Tiedemann, Arthur R.; McArthur, E. Durant; Stutz, Howard C.; [and others], compilers. *Proceedings--symposium on the biology of Atriplex and related chenopods; 1983 May 2-6; Provo, UT. Gen. Tech. Rep. INT-172*. Ogden, UT: U.S. Department of Agriculture, Forest Service, Intermountain Forest and Range Experiment Station: Pgs 281-285.
- Rasmussen, L.L. and J.D. Brotherson. 1986. Response of winterfat communities to release from grazing pressure. *Great Basin Naturalist* 46:148-156.
- Rice, B. and M. Westoby. 1978. Vegetative responses of some Great Basin shrub-communities protected against jackrabbits or domestic stock. *Journal of Range Management* 31:28-34.
- Romo, J.T., R.E. Redmann, B.L. Kowalenko, and A.R. Nicholson. 1995. Growth of winterfat following defoliation in northern mixed prairie of Saskatchewan. *Journal of Range Management* 48:240-245.
- Statler, G.D. 1967. *Eurotia lanata* establishment trials. *Journal of Range Management* 20:253-255.
- Stevens, R., B.C. Giunta, K.R. Jorgensen, and A.P. Plummer. Winterfat (*Ceratoides lanata*). Publ. No. 77-2. Salt Lake City, UT: Utah State Division of Wildlife Resources. 41 p.
- Stubbendieck, J., S.L. Hatch, and C.H. Butterfield. 1992. *North American Range Plants*. 4th ed. Lincoln, NE: University of Nebraska Press. 493 p.
- Tirmenstein, D. 1999. *Achnatherum hymenoides*. In: *Fire Effects Information System*, [Online]. U.S. Department of Agriculture, Forest Service, Rocky Mountain Research Station, Fire Sciences Laboratory (Producer). Available: <http://www.fs.fed.us/database/feis/>
- Welch, B.L. 1989. Nutritive value of shrubs. In: McKell, C.M. (ed.). *The biology and utilization of shrubs*. San Diego, CA: Academic Press, Inc. Pgs 405-424.
- Welsh, S.L., N.D. Atwood, S. Goodrich, L.C. Higgins, (eds.). 1987. *A Utah Flora. The Great Basin Naturalist Memoir No. 9*. Provo, UT: Brigham Young University. 894 p.
- West, N.E. 1994. Effects of fire on salt-desert shrub rangelands. In: Monsen, S.B. and S.G. Kitchen (compilers). *Proceedings--ecology and management of annual rangelands; 1992 May 18-22; Boise, ID. Gen. Tech. Rep. INT-GTR-313*. Ogden, UT: U.S. Department of Agriculture, Forest Service, Intermountain Research Station: Pgs 71-74.
- West, N.E. and J. Gasto. 1978. Phenology of the aerial portions of shadscale and winterfat in Curlew Valley, Utah. *Journal of Range Management* 31(1):43-45.
- Wood, B.W. and J.D. Brotherson. 1986. Ecological adaptation and grazing response of budsage (*Artemisia spinescens*). In: McArthur, E.D. and B.L. Welch (compilers). *Proceedings--symposium on the biology of Artemisia and Chrysothamnus; 1984 July 9-13; Provo, UT. Gen. Tech. Rep. INT-200*. Ogden, UT: U.S. Department of Agriculture, Forest Service, Intermountain Research Station: 75-92.
- Woodmansee, R.G. and L.D. Potter. 1971. Natural reproduction of winterfat (*Eurotia lanata*) in New Mexico. *Journal of Range management* 24(1):24-30.
- Workman, J.P. and N.E. West. 1967. Germination of *Eurotia lanata* in relation to temperature and salinity. *Ecology* 48(4):659-661.
- Young, R.P. 1983. Fire as a vegetation management tool in rangelands of the Intermountain Region. In: Monsen, S.B. and N. Shaw (compilers). *Managing Intermountain rangelands--improvement of range and wildlife habitats: Proceedings; 1981 September 15-17; Twin Falls, ID; 1982 June 22-24; Elko, NV. Gen. Tech. Rep. INT-157*. Ogden, UT: U.S. Department of Agriculture, Forest Service, Intermountain Forest and Range Experiment Station: Pgs 18-31.

### Contributors

CP/GKB  
TK Stringham  
P NovakEchenique

### Approval

Kendra Moseley, 3/07/2025

### Rangeland health reference sheet

**Interpreting Indicators of Rangeland Health** is a qualitative assessment protocol used to determine ecosystem condition based on benchmark characteristics described in the Reference Sheet. A suite of 17 (or more) indicators are typically considered in an assessment. The ecological site(s) representative of an assessment location must be known prior to applying the protocol and must be verified based on soils and climate. Current plant community cannot be used to identify the ecological site.

Author(s)/participant(s)	Patti Novak-Echenique
Contact for lead author	State Rangeland Management Specialist
Date	12/17/2009
Approved by	
Approval date	
Composition (Indicators 10 and 12) based on	Annual Production

### Indicators

1. **Number and extent of rills:** Rills are not typical.

---

2. **Presence of water flow patterns:** Water flow patterns are rare to common depending on site location relative to major inflow areas.

---

3. **Number and height of erosional pedestals or terracettes:** Pedestals are none.

---

4. **Bare ground from Ecological Site Description or other studies (rock, litter, lichen, moss, plant canopy are not bare ground):**  
Bare Ground  $\pm$  80 percent.

---

5. Number of gullies and erosion associated with gullies: None

---

6. Extent of wind scoured, blowouts and/or depositional areas: None

---

7. Amount of litter movement (describe size and distance expected to travel): Fine litter (foliage of grasses and annual & perennial forbs) expected to move distance of slope length during periods of intense summer convection storms or run in of early spring snow melt flows. Persistent litter (large woody material) will remain in place except during unusual flooding (ponding) events.

---

8. Soil surface (top few mm) resistance to erosion (stability values are averages - most sites will show a range of values): Soil stability values will range from 3 to 5.

---

9. Soil surface structure and SOM content (include type of structure and A-horizon color and thickness): Structure of soil surface is thin to medium platy. Soil surface colors are very light and soils are typified by an ochric epipedon. Organic matter is typically less than 0.9 percent (OM values taken from lab characterization data).

---

10. Effect of community phase composition (relative proportion of different functional groups) and spatial distribution on infiltration and runoff: Shrubs and deep-rooted perennial herbaceous bunchgrasses, rhizomatous grasses or both aid in infiltration. Shrub canopy and associated litter break raindrop impact and provide opportunity for snow catch and accumulation on site.

---

11. Presence and thickness of compaction layer (usually none; describe soil profile features which may be mistaken for compaction on this site): Compacted layers are not typical. Platy, subangular blocky, or massive subsurface layers are normal for this site and are not to be interpreted as compaction.

---

12. Functional/Structural Groups (list in order of descending dominance by above-ground annual-production or live foliar cover using symbols: >>, >, = to indicate much greater than, greater than, and equal to):

Dominant: Low statured or half shrubs (winterfat & budsage)

Sub-dominant: deep-rooted, cool season, perennial bunchgrasses &gt; shallow-rooted cool season, perennial bunchgrasses &gt; associated shrubs &gt; cool season, rhizomatous grasses = deep-rooted, cool season, perennial forbs = fibrous, shallow-rooted, cool season, perennial and annual forbs

Other: Microbiotic crusts

Additional:

---

13. Amount of plant mortality and decadence (include which functional groups are expected to show mortality or decadence): Dead branches within individual shrubs common and standing dead shrub canopy material may be as much as 35% of total woody canopy.

---

14. **Average percent litter cover (%) and depth ( in):** Between plant interspaces (< 5%) and depth ( $\pm \frac{1}{4}$  in.)

---

15. **Expected annual annual-production (this is TOTAL above-ground annual-production, not just forage annual-production):**  
For normal or average growing season (March thru May)  $\pm$  350 lbs/ac.

---

16. **Potential invasive (including noxious) species (native and non-native).** List species which BOTH characterize degraded states and have the potential to become a dominant or co-dominant species on the ecological site if their future establishment and growth is not actively controlled by management interventions. Species that become dominant for only one to several years (e.g., short-term response to drought or wildfire) are not invasive plants. Note that unlike other indicators, we are describing what is NOT expected in the reference state for the ecological site: Invaders include cheatgrass, annual mustards, annual kochia, Russian thistle, halogeton, and knapweeds

---

17. **Perennial plant reproductive capability:** All functional groups should reproduce in average (or normal) and above average growing season years

---