

Ecological site R024XY007NV

SALINE BOTTOM

Last updated: 3/07/2025
 Accessed: 05/22/2026

General information

Provisional. A provisional ecological site description has undergone quality control and quality assurance review. It contains a working state and transition model and enough information to identify the ecological site.

MLRA notes

Major Land Resource Area (MLRA): 024X–Humboldt Basin and Range Area

Major Land Resource Area (MLRA): 024X–Humboldt Basin and Range Area Major land resource area (MLRA) 24, the Humboldt Area, covers an area of approximately 8,115,200 acres (12,680 sq. mi.). MLRA 24 is in the Great Basin Section of the Basin and Range Province of the Intermontane Plateaus. Elevations predominantly range from 3,950 to 5,900 feet (1,205 to 1,800 meters). The elevations of some mountain peaks are more than 8,850 feet (2,700 meters). A series of widely spaced north-south trending mountain ranges are separated by broad valleys filled with alluvium washed in from adjacent mountain ranges. Most valleys are drained by tributaries to the Humboldt River. Playas, however, occur in lower elevation valleys with closed drainage systems. Isolated ranges are dissected, uplifted fault-block mountains. Geology is comprised of Mesozoic and Paleozoic volcanic rock and marine and continental sediments. Young andesite and basalt flows (6 to 17 million years old) are at the margins of the mountains. Dominant soil orders include Aridisols, Entisols, Inceptisols and Mollisols. Soils of the area are generally characterized by a mesic soil temperature regime, an aridic soil moisture regime and mixed geology. They are generally well drained, loamy and very deep. 75 percent of MLRA 24 is federally owned. The remainder is primarily used for farming, ranching and mining. Irrigated land comprises 3 percent of the area; most of the irrigation water is from surface water sources, such as the Humboldt River and Rye Patch Reservoir. Annual precipitation typically ranges from 6 to 12 inches (15 to 30 cm) for most of the area. In the mountains however the precipitation may be up to 40 inches (101 cm). Most of the annual precipitation is from snow in the winter. In the spring and fall, rainfall occurs as high- intensity, convective thunderstorms. Nevada is on the eastern, lee side of the Sierra Nevada Range; a massive mountain barrier that markedly influences the climate of the State. The prevailing winds are from the west. The warm moist air from the Pacific Ocean ascends the western slopes of the Sierra Range, the air cools, condenses and the moisture falls as precipitation. As the air descends the eastern slope, it is warmed by compression, and very little precipitation occurs. The effects of this mountain barrier are felt not only in the west but throughout the State. The result is the lowlands of Nevada are largely desert or steppes.

Ecological site concept

Saline Bottom ecological site is on alluvial flats, stream terraces and flood plains. Soils are very deep, somewhat poorly drained and formed in alluvium derived from mixed alluvium, loess and volcanic ash. The soil profile is characterized by an ochric epipedon, strong to moderate salinity throughout and a high water table between 70-100cm at some time during the year. Sodicity (SAR) is 13-99 in the upper 50cm and decreases with depth. The soil temperature regime is mesic and an aridic moisture regime exists during the growing season in normal years after ground water levels drop below the moisture control section. Important abiotic factors contributing to this ecological site include sodicity, high amounts of soluble salts in the profile, increased available soil moisture to landscape position and seasonally available ground water.

Associated sites

R024XY064NV	<p>SODIC BOTTOM</p> <p>The soils are very deep and calcareous. Surface soils are less than 10 inches thick and are medium to moderately fine textured.</p>
--------------------	---

R024XY011NV	<p>SODIC FLAT 6-8 P.Z.</p> <p>This ecological site is on alluvial flats. Soils are deep, poorly to somewhat poorly drained, and formed in alluvium derived from mixed rocks with a component of volcanic ash.</p>
R024XY003NV	<p>SODIC TERRACE 6-8 P.Z.</p> <p>Soils are very deep, well drained and formed in a thin layer of loess and alluvium derived from mixed parent material influenced by volcanic ash over lacustrine sediments and characterized by a very low infiltration.</p>
R024XY006NV	<p>DRY FLOODPLAIN</p> <p>This ecological site is on stream terraces and on fan skirts along intermittent drainageways. Soils are very deep, moderately well drained and formed in alluvium derived from mixed rocks with components of loess and volcanic ash.</p>
R024XY009NV	<p>SALINE MEADOW</p> <p>This ecological occurs on floodplains and inset fans. Soils are very deep, poorly drained and formed in alluvium derived from mixed parent material. The soil profile is characterized by a fine sand surface texture.</p>
R024XY063NV	<p>SALINE FLOODPLAIN</p> <p>This ecological site is on alluvial flats, stream terraces and flood plains. Soils are very deep, somewhat poorly drained and formed in alluvium derived from mixed alluvium, loess and volcanic ash.</p>

Similar sites

R025XY003NV	<p>LOAMY BOTTOM 8-14 P.Z.</p> <p>More productive site; Greasewood (SAVE4) and Saltgrass (DISP) absent; soils not saline-alkali affected.</p>
R024XY008NV	<p>SODIC FLAT 8-10 P.Z.</p> <p>Much less productive site Greasewood (SAVE4) and Basin wildrye (LECI4)</p>
R024XY064NV	<p>SODIC BOTTOM</p> <p>Silver buffaloberry (SHAR) - Greasewood (SAVE4)- Shadscale saltbrush (ATCO) codominant shrubs.</p>
R024XY011NV	<p>SODIC FLAT 6-8 P.Z.</p> <p>Much less productive site. Greasewood (SAVE4), Basin wildrye (LECI4)</p>
R024XY006NV	<p>DRY FLOODPLAIN</p> <p>Soils are moderately well drained. Basin Big sagebrush (ARTRT) dominant shrub; Greasewood (SAVE4) minor shrub, or absent.</p>

R024XY063NV	SALINE FLOODPLAIN SHAR and Basin Big sagebrush (ARTRT) codominant shrubs; Greasewood (SAVE4) minor shrub, to absent.
--------------------	--

Table 1. Dominant plant species

Tree	Not specified
Shrub	(1) <i>Sarcobatus vermiculatus</i>
Herbaceous	(1) <i>Leymus cinereus</i>

Physiographic features

This ecological site is on alluvial flats, stream terraces and flood plains. Slopes range from 0 to 4 percent but are typically less than 2 percent. Elevations are from 3900 to 6000 (1189 to 1829 meters) feet.

Table 2. Representative physiographic features

Landforms	(1) Alluvial flat (2) Stream terrace (3) Flood plain
Runoff class	Medium to very high
Flooding duration	Very brief (4 to 48 hours) to brief (2 to 7 days)
Flooding frequency	None to occasional
Ponding frequency	None
Elevation	1,190 – 1,830 m
Slope	0 %
Water table depth	30 – 180 cm
Aspect	Aspect is not a significant factor

Climatic features

The climate is semiarid, characterized by cold, moist winters, and warm, somewhat dry summers. Average annual precipitation is 6 to 10 inches (15 to 25 cm). Mean annual air temperature is 45 to 53 degrees F. The average growing season is about 90 to 130 days.

Table 3 Representative climatic features

Frost-free period (characteristic range)	80-80 days
Freeze-free period (characteristic range)	100-110 days
Precipitation total (characteristic range)	230-250 mm
Frost-free period (actual range)	80-80 days
Freeze-free period (actual range)	100-110 days
Precipitation total (actual range)	200-250 mm
Frost-free period (average)	80 days
Freeze-free period (average)	110 days
Precipitation total (average)	230 mm

- (1) WINNEMUCCA MUNI AP [USW00024128], Winnemucca, NV
- (2) OROVADA 3 W [USC00265818], Orovada, NV

Influencing water features

This site is associated with perennial streams and dry lake beds.

Soil features

The soils associated with this site are very deep, somewhat poorly drained and formed in alluvium derived from mixed parent material with loess and volcanic ash.

These soils are characterized by ochric epipedon and are strongly salt and sodium affected in their upper profile with SAR ranging from 13-99. Salt and sodium concentrations typically decrease with depth.

A seasonal high water table exists at depths of 20 to 60 inches (51 to 152 cm). Additional moisture is received on this site during the winter and spring months as run-in from higher landscapes or occasional brief overflow from adjacent streams. Wetting of these soils dilutes their salt and sodium concentrations and the degree of salinity and alkalinity fluctuates widely through the year. Permeability is slow to moderately slow. Seed viability, germination, and available water capacity is reduced due to the saline condition of these soils. The surface layer of these soils will crust and bake upon drying, inhibiting water infiltration and seedling emergence.

A representative soil series associated with this ecological site is Wendane, a fine-silty, mixed, superactive, calcareous, mesic Aquandic Halaquepts.

Soil series associated with this site includes: Argenta, Bloor, Cluro, Delvada, Dunphy, Humbolt, Hussa, Ocala, paranat, Rixie, Rose Creek, Sonoma, Umlerland, Wendane, and Yobe.

Table 4. Representative soil features

Parent material	(1) Alluvium (2) Loess (3) Volcanic ash
Surface texture	(1) Silt loam (2) Very fine sandy loam (3) Silty clay loam
Family particle size	(1) Loamy
Drainage class	Somewhat poorly drained to poorly drained
Permeability class	Slow to moderately slow
Soil depth	180 – 210 cm
Surface fragment cover <=3"	Not specified
Surface fragment cover >3"	Not specified
Available water capacity (0-101.6cm)	15.24 – 21.08 cm
Calcium carbonate equivalent (0-101.6cm)	10 – 20 %
Electrical conductivity (0-101.6cm)	20 – 50 mmhos/cm
Sodium adsorption ratio (0-101.6cm)	0 – 100
Soil reaction (1:1 water) (0-101.6cm)	7.4 – 10

Subsurface fragment volume <=3" (Depth not specified)	0 – 10 %
Subsurface fragment volume >3" (Depth not specified)	Not specified

Ecological dynamics

An ecological site is the product of all the environmental factors responsible for its development and it has a set of key characteristics that influence a site's resilience to disturbance and resistance to invasives. Key characteristics include 1) climate (precipitation, temperature), 2) topography (aspect, slope, elevation, and landform), 3) hydrology (infiltration, runoff), 4) soils (depth, texture, structure, organic matter), 5) plant communities (functional groups, productivity), and 6) natural disturbance regime (fire, herbivory, etc.) (Caudle et al 2013). Biotic factors that influence resilience include site productivity, species composition and structure, and population regulation and regeneration (Chambers et al. 2013).

The Great Basin shrub communities have high spatial and temporal variability in precipitation, both among years and within growing seasons. Nutrient availability is typically low but increases with elevation and closely follows moisture availability. The moisture resource supporting

the greatest amount of plant growth is typically the water stored in the soil profile during the winter. The invasibility of plant communities is commonly linked to resource availability. Disturbance can decrease resource uptake due to damage or mortality of the native species and depressed competition or can increase resource pools by the decomposition of dead plant material following disturbance.

Black greasewood is classified as a phreatophyte (Eddleman 2002), and its distribution is well correlated with the distribution of groundwater (Mozingo 1987). Meinzer (1927) discovered that the taproots of black greasewood could penetrate from 20 to 57 feet (6 to 17 meters) below the surface.

Romo (1984) observed water tables ranging from 3.5 to 15 m under black greasewood dominated communities in Oregon. Black greasewood stands develop best where moisture is readily available, either from surface or subsurface runoff (Brown 1971). It is commonly on floodplains that are either subject to periodic flooding, have a high water table at least part of the year, or have a water table less than 34 feet (10 meters) deep (Harr and Price 1972, Blauer et al. 1976, Branson et al. 1976, Blaisdell and Holmgren 1984, Eddleman 2002). Ganskopp (1986) reported that water tables within 9.8 to 11.8 inches (25 to 30 cm) (of the surface had no effect on black greasewood in Oregon. However, a study, conducted in California, observed black greasewood to not survive six months of continuous flooding (Groeneveld and Crowley 1988, Groeneveld 1990). Black greasewood is typically a deep-rooted shrub but has some shallow roots near the soil surface; the maximum rooting depth can be determined by the depth to a saturated zone (Harr and Price 1972). The perennial bunchgrasses generally have somewhat shallower root systems than the shrubs, but root densities are commonly as high as or higher than those of shrubs in the upper 0.5 m but taper off more rapidly than shrubs. However, basin wildrye is weakly rhizomatous and has root depths of up to 2 meters. Basin wildrye exhibits greater lateral root spread than many other grass species (Abbott et al. 1991, Reynolds and Fraley 1989).

Drought will initially cause a decline in bunchgrasses, but prolonged drought will eventually cause a decline in shrubs, including black greasewood. As site condition deteriorates, these sites may become a pure stand of black greasewood or a pure stand with an annual understory. Marcum and Kopec (1997) observed inland saltgrass more tolerant of increased levels of salinity than alkali sacaton therefore dewatering and/or long term drought causing increased levels of salinity would create environmental conditions more favorable to inland saltgrass over alkali sacaton. Alkali sacaton is considered a facultative wet species in this region; therefore it is not drought tolerant. A lowering of the water table can occur with ground water pumping in these sites. This may contribute to the loss of deep-rooted species such as greasewood and basin wildrye and an increase in rabbitbrush, shadscale and other species with the absence of drought.

Basin wildrye is a large, cool-season perennial bunchgrass with an extensive deep coarse fibrous root system (Reynolds and Fraley 1989). Clumps may reach up to six feet (1.8 meters) in height (Ogle et al 2012b). Basin wildrye does not tolerate long periods of inundation; it prefers cycles of wet winters and dry summers and is most commonly in deep soils with high water holding capacities or seasonally high water tables (Ogle et al 2012b, Perryman and Skinner 2007).

Seasonally high water tables are necessary for maintenance of site productivity and reestablishment of basin wildrye stands following disturbances such as fire, drought or excessive herbivory (Eckert et al. 1973). The sensitivity of basin wildrye seedling establishment to reduced soil water availability is increased as soil pH increases (Stuart et al. 1971). Lowering of the water table through extended drought, channel incision or water pumping will decrease basin wildrye production and establishment, while sagebrush, black greasewood, rabbitbrush, and invasive weeds increase. Farming and abandonment may facilitate the creation of surface vesicular crust, increased surface ponding, and decreased infiltration; which leads to dominance by sprouting shrubs and an annual understory.

This ecological site has moderate resilience to disturbance and resistance to invasion. Primary disturbances on these ecological sites is drought, fire, flooding, Aroga infestation (Aroga websteri), and channel incision or other disturbance leading to a lowered seasonal water table. This facilitates an increase in shrubs and a decrease in basin wildrye. The introduction of annual weedy species, like cheatgrass (*Bromus tectorum*), may cause an increase in fire frequency and eventually lead to an annual state or a state dominated by rabbitbrush. Other troublesome non-native weeds such as broadleaved pepperweed or tall whitetop (*Lepidium latifolium*), hoary cress or whitetop (*Cardaria draba*), scotch cottonthistle (*Onopordum acanthium*) or bull thistle (*Cirsium vulgare*) are potential invaders on this site. Two

possible alternative stable states have been identified for this site.

Fire Ecology:

Natural fire return intervals are estimated to vary between less than 35 years up to 100 years in salt desert ecosystems with basin wildrye (Paysen et al. 2000). Higher production sites would have experienced fire more frequently than lower production sites. Basin wildrye is relatively resistant to fire, particularly dormant season fire, as plants sprout from surviving root crowns and rhizomes (Zschaechner 1985). Fire maintained the grass dominance of these ecosystems. Increases in the fire return interval leads to increases in the shrub component of the plant community, potentially facilitating increases in bare ground, inland saltgrass and invasive weeds. Lack of fire combined with excessive herbivory converts these sites to black greasewood dominance.

Basin wildrye is relatively resistant to fire, particularly dormant season fire, as plants sprout from surviving root crowns and rhizomes (Zschaechner 1985). Fire maintained the grass dominance of these ecosystems, therefore increases in the fire return interval favors increases in the shrub component of the plant community. The reduction of grasses potentially facilitates increases in bare ground, inland salt grass, and invasive weeds. Lack of fire combined with excessive herbivory converts these sites to sagebrush, black greasewood, and rabbitbrush dominance.

Historically, black greasewood-saltbush communities had sparse understories and bare soil in intershrub spaces, making these communities somewhat resistant to fire (Young 1983, Paysen et al. 2000). They may burn only during high fire hazard conditions; for example, years with high precipitation can result in almost continuous fine fuels, increasing fire hazard (West 1994, Paysen et al. 2000). Black greasewood may be killed by severe fires, but can resprout after low to moderate severity fires (Robertson 1983, West 1994). Bentz et al. (2008) reported that

following a Nevada wildfire, black greasewood sprouts reached approximately 2.5 feet (76 cm) within 3 years. Grazing and other disturbance may result in increased biomass production due to sprouting and increased seed production, also leading to greater fuel loads (Sanderson and Stutz 1994, Paysen et al. 2000).

During settlement, many of the cattle in the Great Basin were wintered on extensive basin wildrye stands, however due to sensitivity to spring use many stands were decimated by early in the 20th century (Young et al. 1975). Less palatable species such as black greasewood, rabbitbrush and inland saltgrass increased in dominance along with invasive non-native species such as fivehook bassia, mustards, halogeton and cheatgrass (Roundy 1985). Spring defoliation of basin wildrye and/or consistent, heavy grazing during the growing season may significantly reduce basin wildrye production and density (Krall et al. 1971). Thus, inadequate rest and recovery from defoliation can cause a decrease in basin wildrye and an increase in rabbitbrush and black greasewood, along with inland saltgrass and non-native weeds (Young et al. 1975, Roundy 1985). Additionally, natural basin wildrye seed viability may be low and seedlings lack vigor (Young and Evans 1981). Roundy (1985) observed that although basin wildrye is adapted to seasonally dry saline soils, high and frequent spring precipitation is necessary to establish it from seed suggesting that establishment of natural basin wildrye seedlings happens only during years of unusually high precipitation. Therefore, reestablishment of a stand that has been decimated by grazing may be episodic.

Hydrology of these basin wildrye dominated sites is also critical for site function and maintenance. Seasonally high water tables have been necessary for maintenance of site productivity and reestablishment of basin wildrye stands following disturbances such as fire, drought or excessive herbivory (Eckert et al. 1973). The sensitivity of basin wildrye seedling establishment to reduced soil water availability is increased as soil pH increases (Stuart et al. 1971). Lowering of the water table through extended drought or water pumping will decrease basin wildrye production and establishment while black greasewood, rabbitbrush, inland saltgrass and invasive weeds increase. Farming of saline-sodic soils may cause an increase in soil pH in the surface horizons leading to soil surface sealing, increased ponding and reduced infiltration. Additionally farming may facilitate the creation of a cemented layer (plow pan) causing a reduction in root penetration of seedlings. Cessation of farming leads to rabbitbrush dominance, increased bare ground and the potential for weed invasion.

As ecological condition declines and where management results in abusive grazing use by livestock, rabbitbrush and black greasewood increase and become the dominant vegetation in lower condition classes. Inland saltgrass increases as condition declines and typically dominates the understory when this site is in low ecological condition. Fivehook bassia and annual mustards are species likely to invade this site.

Fire Ecology:

Black greasewood communities have been historically subject to stand-replacing fire regimes with intervals of

State and transition model

Figure 7. T. Stringham 8/2016

Figure 8. T Stringham 8/2016

Additional community tables

Table 5. Community 1.1 plant community composition

Group	Common Name	Symbol	Scientific Name	Annual Production ()	Foliar Cover (%)
Grass/Grasslike					
1	Primary Perennial Grasses			894-1459	
	basin wildrye	LECI4	<i>Leymus cinereus</i>	785-942	–
	alkali sacaton	SPAI	<i>Sporobolus airoides</i>	78-392	–
	saltgrass	DISP	<i>Distichlis spicata</i>	31-126	–
2	Secondary Perennial Grasses			31-126	
	squirreltail	ELEL5	<i>Elymus elymoides</i>	8-78	–
	bluegrass	POA	<i>Poa</i>	8-78	–
	alkaligrass	PUCCI	<i>Puccinellia</i>	8-78	–
Forb					
3	Perennial Forbs			31-126	
	basin wildrye	LECI4	<i>Leymus cinereus</i>	785-942	–
	milkvetch	ASTRA	<i>Astragalus</i>	8-31	–
	povertyweed	IVAX	<i>Iva axillaris</i>	8-31	–
	niterwort	NITRO	<i>Nitrophila</i>	8-31	–
	thelypody	THELY	<i>Thelypodium</i>	8-31	–
Shrub/Vine					
4	Primary Shrubs			94-314	
	greasewood	SAVE4	<i>Sarcobatus vermiculatus</i>	78-235	–
	rubber rabbitbrush	ERNAN5	<i>Ericameria nauseosa ssp. nauseosa var. nauseosa</i>	16-78	–
5	Secondary Shrubs			25-99	
	alkali sacaton	SPAI	<i>Sporobolus airoides</i>	78-392	–
	shadscale saltbush	ATCO	<i>Atriplex confertifolia</i>	16-47	–
	sickle saltbush	ATFA	<i>Atriplex falcata</i>	16-47	–
	Torrey's saltbush	ATTO	<i>Atriplex torreyi</i>	16-47	–
	silver buffaloberry	SHAR	<i>Shepherdia argentea</i>	16-47	–
	seepweed	SUAED	<i>Suaeda</i>	16-47	–

Table 6. Community 1.2 plant community composition

Group	Common Name	Symbol	Scientific Name	Annual Production ()	Foliar Cover (%)
-------	-------------	--------	-----------------	----------------------	------------------

Table 7. Community 1.3 plant community composition

Group	Common Name	Symbol	Scientific Name	Annual Production ()	Foliar Cover (%)
-------	-------------	--------	-----------------	----------------------	------------------

Table 8. Community 2.1 plant community composition

Group	Common Name	Symbol	Scientific Name	Annual Production ()	Foliar Cover (%)
-------	-------------	--------	-----------------	----------------------	------------------

Table 9. Community 2.2 plant community composition

Group	Common Name	Symbol	Scientific Name	Annual Production ()	Foliar Cover (%)
-------	-------------	--------	-----------------	----------------------	------------------

Table 10. Community 2.3 plant community composition

Group	Common Name	Symbol	Scientific Name	Annual Production ()	Foliar Cover (%)
-------	-------------	--------	-----------------	----------------------	------------------

Table 11. Community 3.1 plant community composition

Group	Common Name	Symbol	Scientific Name	Annual Production ()	Foliar Cover (%)
-------	-------------	--------	-----------------	----------------------	------------------

Table 12. Community 3.2 plant community composition

Group	Common Name	Symbol	Scientific Name	Annual Production ()	Foliar Cover (%)
-------	-------------	--------	-----------------	----------------------	------------------

Animal community

Livestock Interpretations: This site is suitable for livestock grazing. Grazing management should be keyed to dominant grasses and palatable shrubs production. The early growth and abundant production of basin wildrye make it a valuable source of forage for livestock. It is important forage for cattle and is readily grazed by cattle and horses in early spring and fall. Though coarse textured during the winter, basin wildrye may be utilized more frequently by livestock and wildlife when snow has covered low shrubs and other grasses. Alkali sacaton is a valuable forage species in arid and semiarid regions. Plants are tolerant to moderate grazing and can produce abundant herbage utilized by livestock. Saltgrass's value as forage depends primarily on the relative availability of other grasses of higher nutritional value and palatability. It can be an especially important late summer grass in arid environments after other forage grasses have decreased. Saltgrass is rated as a fair to good forage species only because it stays green after most other grasses dry. Livestock generally avoid saltgrass due to its coarse foliage. Saltgrass is described as an increaser under grazing pressure. Black greasewood is an important winter browse plant for domestic sheep and cattle. It also receives light to moderate use by domestic sheep and cattle during spring and summer months. Black greasewood contains soluble sodium and potassium oxalates that may cause poisoning and death in domestic sheep and cattle if large amounts are consumed in a short time. In general, livestock forage only lightly on rubber rabbitbrush during the summer, but winter use can be heavy in some locations. Fall use is variable, but flowers are commonly used by livestock. A few leaves and the more tender stems may also be used. Stocking rates vary over time depending upon season of use, climate variations, site, and previous and current management goals. A safe starting stocking rate is an estimated stocking rate that is fine tuned by the client by adaptive management through the year and from year to year. **Wildlife Interpretations:** Basin wildrye provides winter forage for mule deer, though use is commonly low compared to other native grasses. Basin wildrye provides summer forage for black-tailed jackrabbits. Because basin wildrye remains green throughout early summer, it remains available for small mammal forage for longer time than other grasses. The western salt desert shrub and grassland communities where alkali sacaton is common support an abundance of mule deer, pronghorn, carnivores, small mammals, birds, amphibians, and reptiles. Saltgrass provides cover for a variety of bird species, small mammals, and arthropods and is on occasion used as forage for several big game wildlife species. Black greasewood is an important winter browse plant for big game animals and a food source for many other wildlife species. It also receives light to moderate use by mule deer and pronghorn during spring and summer months. Wildlife forage only lightly on rubber rabbitbrush during the summer, but winter use can be heavy in some locations. Fall use is variable, but flowers are commonly used by wildlife. A few leaves and the more tender stems may also be used. The forage value of rubber rabbitbrush varies greatly among subspecies and ecotypes.

Hydrological functions

Runoff is low to very high. Permeability is slow to moderately slow. Hydrologic soil groups are B, C, and D.

Recreational uses

Aesthetic value is derived from the diverse floral and faunal composition and the colorful flowering of wild flowers and shrubs during the spring and early summer. This site offers rewarding opportunities to photographers and for nature study. This site has potential for upland bird and big game hunting.

Other products

Basin wildrye is used as bedding for various Native American ceremonies, providing a cool place for dancers to stand. The leaves, seeds and stems of black greasewood are edible.

Other information

Basin wildrye is useful in mine reclamation, fire rehabilitation and stabilizing disturbed areas. Its usefulness in range seeding, however, may be limited by initially weak stand establishment. Alkali sacaton is one of the most used species for seeding and stabilizing disturbed lands. Due to alkali sacaton's salt tolerance, is recommended for native grass seeding on subirrigated saline sites. Given its extensive system of rhizomes and roots which form a dense sod, saltgrass is considered a suitable species for controlling wind and water erosion. Black greasewood is useful for stabilizing soil on wind-blown areas. It successfully revegetates eroded areas and sites too saline for most plant species.

Inventory data references

NASIS soil component data.

Type locality

Location 1: Lander County, NV

Township/Range/Section	T31N R45E S17
UTM zone	N
UTM northing	4490010
UTM easting	506009
Latitude	40° 33'39"
Longitude	116° 55'44"
General legal description	S½, Approximately 6 miles south of Battle Mountain, along east side of Marvel Ranch Road (Old 8A), Lander County, Nevada. This site also occurs in Elko, Eureka, Humboldt and Pershing Counties, Nevada.

Other references

Fire Effects Information System (Online; <http://www.fs.fed.us/database/feis/plants/>).

USDA-NRCS Plants Database (Online; <http://www.plants.usda.gov>).

Eckert, R.E., Jr., A.D. Bruner, and G.J. Klomp. 1973. Productivity of tall wheatgrass and great basin wildrye under irrigation on a greasewood – rabbitbrush range site. *Journal of Range Management* 26(4):286-288.

Krall, J.L., J.R. Stroh, C.S. Cooper, and S.R. Chapman. 1971. Effect of time and extent of harvesting on basin wildrye. *Journal of Range Management* 24:414-418.

Paysen, T.E., R.J. Ansley, and J.K. Brown. 2000. Fire in western shrubland, woodland, and grassland ecosystems. In: Brown, J.K. and J.K. Smith (eds). *Wildland fire in ecosystems: Effects of fire on flora*. Gen. Tech. Rep. RMRS-GTR-42-Volume 2. Ogden, UT. U.S. Department of Agriculture, Forest Service, Rocky Mountain Research Station. Pgs 121-159.

Roundy, B.A. 1985. Emergence and establishment of basin wildrye and tall wheatgrass in relation to moisture and salinity. *Journal of Range Management* 38(2):126-131.

Stuart, D.M., G.E. Schuman, and A.S. Dylla. 1971. Chemical characteristics of the coppice dune soils in Paradise Valley, Nevada. *Proceedings of the Soil Science Society of America* 34:607-611.

Young, J.A., R.A. Evans, and P.T. Tueller. 1975. Great Basin plant communities – pristine and grazed. Pgs 187-212, In: R. Elston (ed.). *Holocene climate in the Great Basin, Occasional Papers, Nevada Archeological Survey, Reno NV*.

Young, J.A. and R.A. Evans. 1981. Germination of Great Basin wildrye seeds collected from native stands. *Agronomy Journal* 73:917-920.

Zschaechner, G.A. 1985. Studying rangeland fire effects: a case study in Nevada. In: Sanders, K. and J. Durham (eds). *Rangeland fire effects. Proceedings of the symposium. 1984 November 27-29; Boise, ID. Boise, ID. U.S. Department of the Interior, Bureau of Land Management, Idaho State Office. Pgs 66-84.*

Contributors

CP/GKB
E.Hourihan

Approval

Kendra Moseley, 3/07/2025

Rangeland health reference sheet

Interpreting Indicators of Rangeland Health is a qualitative assessment protocol used to determine ecosystem condition based on benchmark characteristics described in the Reference Sheet. A suite of 17 (or more) indicators are typically considered in an assessment. The ecological site(s) representative of an assessment location must be known prior to applying the protocol and must be verified based on soils and climate. Current plant community cannot be used to identify the ecological site.

Author(s)/participant(s)	Patti Novak-Echenique
Contact for lead author	State Rangeland Management Specialist
Date	09/24/2009
Approved by	
Approval date	
Composition (Indicators 10 and 12) based on	Annual Production

Indicators

1. **Number and extent of rills:** Rills are typically none.

2. **Presence of water flow patterns:** Water flow patterns are rare to common. Moderately fine to fine surface textures result in limited infiltration rates. Concentrations of surface salts and sodium result in chemical crusts which also impedes precipitation infiltration. Water flow patterns are typically short, ending in depressional areas.

3. **Number and height of erosional pedestals or terracettes:** Pedestals are none.

4. **Bare ground from Ecological Site Description or other studies (rock, litter, lichen, moss, plant canopy are not bare ground):**
Bare Ground \pm 65 percent.

5. **Number of gullies and erosion associated with gullies:** None

6. **Extent of wind scoured, blowouts and/or depositional areas:** None

7. **Amount of litter movement (describe size and distance expected to travel):** Fine litter (foliage of grasses and annual & perennial forbs) only expected to move during periods of flooding by adjacent streams. Persistent litter (large woody material) will remain in place except during major flooding events.

8. **Soil surface (top few mm) resistance to erosion (stability values are averages - most sites will show a range of values):** Soil stability values will range from 1 to 4 (To be field tested).

9. **Soil surface structure and SOM content (include type of structure and A-horizon color and thickness):** Structure of soil surface will be platy or massive. Soil surface colors are light and soils are typified by an ochric epipedon. Organic matter can range from 1.5 to 4.5 percent (OM values taken from lab characterization data).

10. **Effect of community phase composition (relative proportion of different functional groups) and spatial distribution on infiltration and runoff:** Deep-rooted perennial grasses (basin wildrye and alkali sacaton] slow runoff and increase infiltration. Tall stature and relatively coarse foliage of basin wildrye and associated litter break raindrop impact and provide opportunity for snow catch and moisture accumulation on site.

11. **Presence and thickness of compaction layer (usually none; describe soil profile features which may be mistaken for compaction on this site):** Compacted layers are not typical. Platy or massive subsurface layers are normal for this site and are not to be interpreted as compaction.

12. **Functional/Structural Groups (list in order of descending dominance by above-ground annual-production or live foliar cover using symbols: >>, >, = to indicate much greater than, greater than, and equal to):**

Dominant: Tall-statured, deep-rooted, cool season, perennial bunchgrasses

Sub-dominant: short-statured rhizomatous grasses & associated perennial grasses and grass-like plants & tall shrubs & deep-rooted, cool season, perennial forbs = fibrous, shallow-rooted, cool season, perennial and annual forbs

Other:

Additional:

13. **Amount of plant mortality and decadence (include which functional groups are expected to show mortality or decadence):** Dead branches within individual shrubs common and standing dead shrub canopy material may be as much as 25 percent of total woody canopy.

14. **Average percent litter cover (%) and depth (in):** Within plant interspaces (\pm 35 percent) and depth of litter \pm 1/2 inch.

15. **Expected annual annual-production (this is TOTAL above-ground annual-production, not just forage annual-production):**
For normal or average growing season (through end of May) \pm 1400 lbs/ac; Winter moisture significantly affects total production
-

16. **Potential invasive (including noxious) species (native and non-native). List species which BOTH characterize degraded states and have the potential to become a dominant or co-dominant species on the ecological site if their future establishment and growth is not actively controlled by management interventions. Species that become dominant for only one to several years (e.g., short-term response to drought or wildfire) are not invasive plants. Note that unlike other indicators, we are describing what is NOT expected in the reference state for the ecological site: Increasesers include rubber rabbitbrush. Potential invaders include annual mustards, povertyweed, annual kochia, pigweed, and tall whitetop (perennial pepperweed).**
-

17. **Perennial plant reproductive capability: All functional groups should reproduce in average (or normal) and above average growing season years.**
-