

Ecological site R024XY067NV SHALLOW SILTY 5-8 P.Z.

Last updated: 3/06/2025
Accessed: 05/29/2026

General information

Provisional. A provisional ecological site description has undergone quality control and quality assurance review. It contains a working state and transition model and enough information to identify the ecological site.

MLRA notes

Major Land Resource Area (MLRA): 024X–Humboldt Basin and Range Area

Major land resource area (MLRA) 24, the Humboldt Area, covers an area of approximately 8,115,200 acres (12,680 sq. mi.). It is found in the Great Basin Section of the Basin and Range Province of the Intermontane Plateaus. Elevations range from 3,950 to 5,900 feet (1,205 to 1,800 meters) in most of the area, some mountain peaks are more than 8,850 feet (2,700 meters). A series of widely spaced north-south trending mountain ranges are separated by broad valleys filled with alluvium washed in from adjacent mountain ranges. Most valleys are drained by tributaries to the Humboldt River. However, playas occur in lower elevation valleys with closed drainage systems. Isolated ranges are dissected, uplifted fault-block mountains. Geology is comprised of Mesozoic and Paleozoic volcanic rock and marine and continental sediments. Occasional young andesite and basalt flows (6 to 17 million years old) occur at the margins of the mountains. Dominant soil orders include Aridisols, Entisols, Inceptisols and Mollisols. Soils of the area are generally characterized by a mesic soil temperature regime, an aridic soil moisture regime and mixed geology. They are generally well drained, loamy and very deep. Approximately 75 percent of MLRA 24 is federally owned, the remainder is primarily used for farming, ranching and mining. Irrigated land makes up about 3 percent of the area; the majority of irrigation water is from surface water sources, such as the Humboldt River and Rye Patch Reservoir. Annual precipitation ranges from 6 to 12 inches (15 to 30 cm) for most of the area, but can be as much as 40 inches (101 cm) in the mountain ranges. The majority of annual precipitation occurs as snow in the winter. Rainfall occurs as high-intensity, convective thunderstorms in the spring and fall. Nevada lies on the eastern, lee side of the Sierra Nevada Range, a massive mountain barrier that markedly influences the climate of the State. The prevailing winds are from the west, and as the warm moist air from the Pacific Ocean ascends the western slopes of the Sierra Range, the air cools, condensation takes place and most of the moisture falls as precipitation. As the air descends the eastern slope, it is warmed by compression, and very little precipitation occurs. The effects of this mountain barrier are felt not only in the west but throughout the State, with the result that the lowlands of Nevada are largely desert or steppes.

Ecological site concept

This site is on lake plains. Soils are very deep, well drained and formed in alluvium. The soil profile is characterized by an ochric epipedon, continuous weak cementation with in 20 inches (51cm), and coarse silty particle size control section. This site will be field checked for possible correlation to 024XY003NV.

Associated sites

R024XY008NV	<p>SODIC FLAT 8-10 P.Z.</p> <p>This site presents crusting & baking of the surface layer upon drying, inhibiting water infiltration and seedling emergence. High salt concentrations reduce seed viability, germination and the available water capacity of these soils. Black greasewood (SAVE4) dominant shrub; Shadscale (ATCO) minor shrub; Basin wildrye (LECI4) dominant grass.</p>
R024XY011NV	<p>SODIC FLAT 6-8 P.Z.</p> <p>The soil profile is characterized by an ochric epipedon, a pH greater than 8.0 throughout, sodicity (SAR) greater than 30 and a water table between 35 to 59 inches (90 to 150cm) at some point during the spring. Dominant plant species are Black greasewood (SAVE4), Basin wildrye (LECI4), and saltgrass (DISP).</p>

R024XY003NV	<p>SODIC TERRACE 6-8 P.Z.</p> <p>This ecological site is on lake plains and basin floor remnants. Soil are very deep, well drained and formed in a thin layer of loess and alluvium derived from mixed parent material influenced by volcanic ash over lacustrine sediments. Soils are characterized by a very low infiltration. The reference state is characterized by the mixing of shadscale (ATCO) and black greasewood (SAVE4) and approximate canopy cover is less than 15 percent.</p>
--------------------	---

Similar sites

R024XY025NV	<p>LOAMY SLOPE 5-8 P.Z.</p> <p>Indian ricegrass (ACHY) is a minor component; Bud sagebrush (ARSP5) is a major shrub.</p>
R024XY060NV	<p>SHALLOW SILTY 8-10 P.Z.</p> <p>Wyoming sagebrush (ARTRW8) codominant shrub; more productive site.</p>

Table 1. Dominant plant species

Tree	Not specified
Shrub	(1) <i>Atriplex confertifolia</i>
Herbaceous	(1) <i>Elymus elymoides</i>

Physiographic features

This site is on alluvial flats, lake terraces, basin-floor remnants, fan skirts and lake plains. The site is commonly associated with playas and is dissected by shallow intermittent drainageways. Slope gradients of 0 to 2 percent are typical. Elevations are 4000 to 4600 feet (1219 to 1402 m).

Table 2. Representative physiographic features

Landforms	(1) Lake terrace (2) Basin-floor remnant (3) Lake plain
Runoff class	Low to medium
Flooding frequency	None
Ponding frequency	None
Elevation	1,220 – 1,400 m

Slope	0 %
Water table depth	180 cm
Aspect	Aspect is not a significant factor

Climatic features

The climate associated with this site is semiarid and characterized by cool, moist winters and warm, dry summers. Average annual precipitation is 4 to 8 inches (10 to 15cm). Mean annual air temperature is 52 to 56 degrees F. The average growing season is about 100 to 130 days.

Table 3 Representative climatic features

Frost-free period (average)	130 days
Freeze-free period (average)	
Precipitation total (average)	200 mm

Influencing water features

There are no influencing water features associated with this site.

Soil features

The soils associated with this site are very deep, well drained and formed in alluvium from mixed parent material and lacustrine deposits. The soil profile is characterized by an ochric epipedon, continuous weak cementation with in 20 inches (51cm), and coarse silty particle size control section.

The silty surface texture results in plating or surface crusting that decreases infiltration of water and restricts seed germination and establishment. Available water capacity is moderate. The potential for sheet and rill erosion is slight. The soil series associated with this site include: Cresal and Mazuma.

Table 4. Representative soil features

Parent material	(1) Alluvium (2) Lacustrine deposits
Surface texture	(1) Silt loam
Family particle size	(1) Loamy
Drainage class	Well drained
Permeability class	Moderately slow

Soil depth	180 – 210 cm
Surface fragment cover <=3"	Not specified
Surface fragment cover >3"	Not specified
Available water capacity (0-101.6cm)	12.95 – 18.8 cm
Calcium carbonate equivalent (0-101.6cm)	0 – 20 %
Electrical conductivity (0-101.6cm)	0 – 30 mmhos/cm
Sodium adsorption ratio (0-101.6cm)	0 – 100
Soil reaction (1:1 water) (0-101.6cm)	7.9 – 9.6
Subsurface fragment volume <=3" (Depth not specified)	0 – 10 %
Subsurface fragment volume >3" (Depth not specified)	Not specified

Ecological dynamics

An ecological site is the product of all the environmental factors responsible for its development and it has a set of key characteristics that influence a site's resilience to disturbance and resistance to invasives. Key characteristics include 1) climate (precipitation, temperature), 2) topography (aspect, slope, elevation, and landform), 3) hydrology (infiltration, runoff), 4) soils (depth, texture, structure, organic matter), 5) plant communities (functional groups, productivity), and 6) natural disturbance regime (fire, herbivory, etc.) (Caudle 2013). Biotic factors that influence resilience include site productivity, species composition and structure, and population regulation and regeneration (Chambers et al. 2013).

Shadscale is a densely clumped, rounded, compact native shrub. It generally attains heights of 8 to 32 inches and widths of 12 to 68 inches (Blaisdell and Holmgren 1984). Shadscale is considered an evergreen to partially deciduous shrub, since a small percentage of leaves are dropped in the winter (Smith and Nobel 1986). Shadscale possesses wider ecological amplitude than most *Atriplex* species (Crofts and Van Epps 1975), and shows ploidy levels from diploid (2x) to decaploid (10x). The extensive polyploidy of shadscale is an important consideration when implementing revegetation projects because ploidy levels are usually associated with distinct habitats (Sanderson et al. 1990). Diploid individuals are unlikely to perform as well in areas where tetraploids are more common. Diploid individuals generally occur above Pleistocene lake levels, whereas lake floors are usually occupied by autotetraploids. Overall, tetraploids are the most widespread throughout its range (Carlson 1984). Thus, the shadscale most associated with this site is a tetraploid. Bud

sagebrush, a common shrub to this ecological site, is a native, summer-deciduous shrub. It is low growing, spinescent, aromatic shrub with a height of 4 to 10 inches and a spread of 8 to 12 inches (Chambers and Norton 1993). Indian ricegrass, the dominant grass within this site, is a hardy, cool-season, densely tufted, native perennial bunchgrass that grows from 4 to 24 inches in height (Blaisdell and Holmgren 1984). Squirreltail, another native, cool-season perennial bunchgrass, is often a co-dominant with Indian ricegrass.

Shadscale has experienced widespread mortality during periods of above average precipitation (Nelson et al. 1990a). The roots of desert shrubs are sensitive to the level of soil oxygen, waterlogging reduces soil oxygen. Waterlogging causes physiological changes in plants increasing susceptibility to parasite and disease where prolonged period of high soil moisture occur (Nelson et al. 1990a). Periods of elevated precipitation result in increased soil moisture and salinity, which predisposes the roots of the shrubs to pathogenic root rot organisms (Weber et al. 1990). Shadscale occurs in widespread genetically uniform populations on the edaphically consistent soil of the Pleistocene lake bottoms, setting the stage for extensive areas of plant death (Nelson et al. 1990a). Valley bottoms and upland depressions typically exhibit the greatest concentration of die back, due to ponding and run-in moisture. Shadscale is also susceptible to insect attack. Scale insects and mealy bugs have been found in the crown and upper root zone of shadscale plants during periods of dieoff (Nelson et al. 1990b).

Historically, shadscale dominant salt-desert shrub communities were free of exotic invaders; however, excessive grazing pressure during settlement and into the 20th century has increased the overall presence of cheatgrass, halogeton, Russian thistle and weedy mustard species (Peters and Bunting 1994). The lack of continuous fuels to carry fires made fire rare to non-existent in shadscale communities (Young and Tipton 1990), thus it is not surprising that shadscale and bud sagebrush are both fire intolerant (Banner 1992, West 1994). Shadscale does not readily recover from fire, except for establishment through seed (West 1994). The slow reestablishment allows for easy invasion by cheatgrass and other non-native weedy species (Sanderson et al. 1990). The increased presence of exotic annual grasses has greatly altered fire regimes in areas of the Intermountain West where shadscale is a major vegetational component. Exotic annuals increase fire frequency under wet to near-normal summer moisture conditions and repeated, frequent fire has converted large expanses of shadscale rangeland to annual non-native plant communities (Knapp 1998). Grazing exclusion for 2 or more years is beneficial for revegetation of postfire shadscale communities as first year shadscale seedlings lack spines and are highly susceptible to browsing. Spines develop in the second year (Zielinski 1994).

Fire Ecology:

Shadscale does not readily recover from fire, except for establishment through seed (West 1994). The slow reestablishment allows for easy invasion by cheatgrass and other non-native weedy species (Sanderson et al. 1990). The increased presence of exotic annual grasses has greatly altered fire regimes in areas of the Intermountain West where shadscale is a major vegetation component. Exotic annuals increase fire frequency under wet to near-normal summer moisture conditions and repeated, frequent fire has converted large expanses of shadscale rangeland to annual non-native plant communities (Knapp 1998).

The effect of fire on bunchgrasses relates to culm density, culm-leaf morphology, and the size of the plant. The initial condition of bunchgrasses within the site along with seasonality and intensity of the fire all factor into the individual species response. For most forbs and grasses the growing points are located at or below the soil surface providing relative protection from disturbances which decrease above ground biomass, such as grazing or fire. Thus, fire mortality is more correlated to duration and intensity of heat which is related to culm density, culm-leaf morphology, size of plant and abundance of old growth (Wright 1971, Young 1983). However, season and severity of the fire and post-fire soil moisture availability will influence plant response.

Indian ricegrass is a deep-rooted, cool season perennial bunchgrass that is adapted primarily to sandy soils. A prominent grass on this site, it is fairly fire tolerant (Wright 1985), which is likely due to its low culm density and below ground plant crowns. Vallentine (1989) cites several studies in the sagebrush zone that classified Indian ricegrass as being slightly damaged from late summer burning. Indian ricegrass has also been found to reestablish on burned sites through seed dispersed from adjacent unburned areas (Young 1983, West 1994). Thus the presence of surviving, seed producing plants facilitates the reestablishment of Indian ricegrass. Grazing management following fire to promote seed production and establishment of seedlings is important.

Bottlebrush squirreltail is considered more fire tolerant than Indian ricegrass due to its small size, coarse stems, and sparse leafy material (Britton et al. 1990). Postfire regeneration occurs from surviving root crowns and from on- and off-site seed sources. Bottlebrush squirreltail has the ability to produce large numbers of highly germinable seeds, with relatively rapid germination (Young and Evans 1977) when exposed to the correct environmental cues. Early spring growth and ability to grow at low temperatures contribute to the persistence of bottlebrush squirreltail among cheatgrass dominated ranges (Hironaka and Tisdale 1972).

Rehabilitation following fire will have limited success. Observations from one hundred and seven separate plantings within the shadscale zone in Utah and Nevada indicate a very low success rate (Bleak et al. 1965). Seed from 148 native and non-native grasses, forbs and shrubs were planted from 1937 to 1962 across ten locations. Good seedling stands were obtained with introduced wheatgrasses, but most perished during the first summer. A few plantings of crested wheatgrass (*Agropyron cristatum*), fairway and Siberian wheatgrass (*Agropyron fragile*) along with Russian wildrye (*Psathyrostachys juncea*) maintained stands for 10 or more years but eventually declined to very few plants (Bleak et al. 1965). The primary cause of seeding failures appeared to be the arid climate.

State and transition model

Additional community tables

Table 5. Community 1.1 plant community composition

Group	Common Name	Symbol	Scientific Name	Annual Production ()	Foliar Cover (%)
Grass/Grasslike					

1	Primary Perennial Grasses			4-22	
	squirreltail	ELEL5	<i>Elymus elymoides</i>	4-22	-
2	Secondary Perennial Grasses			4-11	
	Indian ricegrass	ACHY	<i>Achnatherum hymenoides</i>	1-7	-
	saltgrass	DISP	<i>Distichlis spicata</i>	1-7	-
	basin wildrye	LECI4	<i>Leymus cinereus</i>	1-7	-
	Sandberg bluegrass	POSE	<i>Poa secunda</i>	1-7	-
Forb					
3	Perennial Forbs			4-18	
Shrub/Vine					
4	Primary Shrubs			168-191	
	shadscale saltbush	ATCO	<i>Atriplex confertifolia</i>	168-191	-
5	Secondary Shrubs			4-22	
	spiny hopsage	GRSP	<i>Grayia spinosa</i>	2-4	-
	greasewood	SAVE4	<i>Sarcobatus vermiculatus</i>	2-4	-
	seepweed	SUAED	<i>Suaeda</i>	2-4	-
	horsebrush	TETRA3	<i>Tetradymia</i>	2-4	-

Table 6. Community 1.2 plant community composition

Group	Common Name	Symbol	Scientific Name	Annual Production ()	Foliar Cover (%)
-------	-------------	--------	-----------------	----------------------	------------------

Table 7. Community 2.1 plant community composition

Group	Common Name	Symbol	Scientific Name	Annual Production ()	Foliar Cover (%)
-------	-------------	--------	-----------------	----------------------	------------------

Table 8. Community 2.2 plant community composition

Group	Common Name	Symbol	Scientific Name	Annual Production ()	Foliar Cover (%)
-------	-------------	--------	-----------------	----------------------	------------------

Table 9. Community 2.3 plant community composition

Group	Common Name	Symbol	Scientific Name	Annual Production ()	Foliar Cover (%)
-------	-------------	--------	-----------------	----------------------	------------------

Table 10. Community 3.1 plant community composition

Group	Common Name	Symbol	Scientific Name	Annual Production ()	Foliar Cover (%)
-------	-------------	--------	-----------------	----------------------	------------------

Table 11. Community 4.1 plant community composition

Group	Common Name	Symbol	Scientific Name	Annual Production ()	Foliar Cover (%)
-------	-------------	--------	-----------------	----------------------	------------------

Table 12. Community 4.2 plant community composition

Group	Common Name	Symbol	Scientific Name	Annual Production ()	Foliar Cover (%)
-------	-------------	--------	-----------------	----------------------	------------------

Table 13. Community 5.1 plant community composition

Group	Common Name	Symbol	Scientific Name	Annual Production ()	Foliar Cover (%)
-------	-------------	--------	-----------------	----------------------	------------------

Table 14. Community 6.1 plant community composition

Group	Common Name	Symbol	Scientific Name	Annual Production ()	Foliar Cover (%)
-------	-------------	--------	-----------------	----------------------	------------------

Animal community

Livestock Interpretations: This site has limited value for livestock grazing, due to the low forage production. Grazing management should be keyed to dominant grasses and palatable shrubs production. Shadscale is a valuable browse species, providing a source of palatable, nutritious forage for a wide variety of livestock. Shadscale provides good browse for domestic sheep. Shadscale leaves and seeds are an

important component of domestic sheep and cattle winter diets. Bottlebrush squirreltail is very palatable winter forage for domestic sheep of Intermountain ranges. Domestic sheep relish the green foliage. Overall, bottlebrush squirreltail is considered moderately palatable to livestock. Stocking rates vary over time depending upon season of use, climate variations, site, and previous and current management goals. A safe starting stocking rate is an estimated stocking rate that is fine tuned by the client by adaptive management through the year and from year to year. Wildlife Interpretations: Shadscale is a valuable browse species, providing a source of palatable, nutritious forage for a wide variety of wildlife particularly during spring and summer before the hardening of spiny twigs. It supplies browse, seed, and cover for birds, small mammals, rabbits, deer, and pronghorn antelope. Bottlebrush squirreltail is a dietary component of several wildlife species. Bottlebrush squirreltail may provide forage for mule deer and pronghorn.

Hydrological functions

Runoff is low to medium. Permeability is moderately slow to moderate. Hydrologic soil group is B. Rills are none. Water flow patterns are rare to common depending on site location relative to major inflow areas. Pedestals are none. There are typically no gullies. Shrubs and deep-rooted perennial herbaceous bunchgrasses aid in infiltration. Sparse shrub canopy and associated litter help break raindrop impact.

Recreational uses

Aesthetic value is derived from the diverse floral and faunal composition and the colorful flowering of wild flowers and shrubs during the spring and early summer. This site offers rewarding opportunities to photographers and for nature study. This site has potential for upland and big game hunting.

Other products

Seeds of shadscale were used by Native Americans for bread and mush.

Other information

Bottlebrush squirreltail is tolerant of disturbance and is a suitable species for revegetation.

Inventory data references

NASIS soil component data.

Type locality

Location 1: Humboldt County, NV	
Township/Range/Section	T35N R34E S6
UTM zone	N
UTM northing	4532338
UTM easting	398683
Latitude	40° 56' 9"
Longitude	118° 12' 12"

General legal description	NW¼ About 6 miles southwest of Imlay, Humboldt Canyon area, Humboldt County, Nevada. This site also occurs in Pershing County, Nevada.
---------------------------	--

Other references

Fire Effects Information System (Online; <http://www.fs.fed.us/database/feis/plants/>).

USDA-NRCS Plants Database (Online; <http://www.plants.usda.gov>).

Banner, R.E. 1992. Vegetation types of Utah. *Journal of Range Management* 14(2):109-114.

Bich, B.S., J.L. Butler, and C.A. Schmidt. 1995. Effects of differential livestock use of key plant species and rodent populations within selected *Oryzopsis hymenoides*/*Hilaria jamesii* communities in Glen Canyon National Recreation Area. *The Southwestern Naturalist* 40(3):281-287.

Blaisdell, J.P. and R.C. Holmgren. 1984. Managing Intermountain rangelands – Salt-desert shrub ranges. USDA-FS General Technical Report INT-163. 52 p.

Booth, D. T., C. G. Howard, and C. E. Mowry. 2006. 'Nezpar' Indian ricegrass: description, justification for release, and recommendations for use. *Rangelands Archives* 2:53-54.

Britton, C.M., G.R. McPherson, and F.A. Sneva. 1990. Effects of burning and clipping on five bunchgrasses in eastern Oregon. *The Great Basin Naturalist* 50(2):115-120.

Carlson, J. 1984. *Atriplex* cultivar development. In: Proceedings--symposium on the biology of *Atriplex* and related chenopods; 1983 May 2-6; Provo, UT. Gen. Tech. Rep. INT-172. Ogden, UT: U.S. Department of Agriculture, Forest Service, Intermountain Forest and Range Experiment Station: Pgs 176-182.

Caudle, D., J. DiBenedetto, M. Karl, H. Sanchez, and C. Talbot. 2013. Interagency ecological site handbook for rangelands. Available at: <http://jornada.nmsu.edu/sites/jornada.nmsu.edu/files/InteragencyEcolSiteHandbook.pdf>. Accessed 4 October 2013.

Chambers, J.C. and B.E. Norton. 1993. Effects of grazing and drought on population dynamics of salt desert species on the Desert Experimental Range, Utah. *Journal of Arid Environments* 24:261-275.

Cibils, A.F., D.M. Swift and D.E. McArthur. 1998. Plant-herbivore interactions in *Atriplex*: current state of knowledge. Gen. Tech. Rep. RMRS-GTR-14. Ogden, UT: U.S. Department of Agriculture, Forest Service, Rocky Mountain Research Station. 31 p.

Cook, C.W. and R.D. Child. 1971. Recovery of desert plants in various states of vigor. *Journal of Range Management* 24(5):339-343.

Crofts, K. and G. Van Epps. 1975. Use of shadscale in revegetation of arid disturbed sites. In: Stutz, H.C. (ed.). *Wildland shrubs: symposium and workshop proceedings*; 1975 November 5-7; Provo, UT. Provo, UT: Brigham Young University: Pgs 151-152.

Eckert, R.E., Jr., F.F. Peterson, and F.L. Emmerich. 1987. A study of factors influencing secondary succession in the sagebrush [*Artemisia* spp. L.] type. In: Frasier, G.W. and R.A. Evans, (eds.). *Proceedings of the symposium: "Seed and seedbed ecology of rangeland plants"*; 1987 April 21-23; Tucson, AZ. Washington, DC: U.S. Department of Agriculture, Agricultural Research Service: Pgs 149-168.

Hironaka, M. and E.W. Tisdale. 1972. Growth and development of *Sitanion hystrix* and *Poa sandbergii*. Research Memorandum RM 72-24. U.S. International Biological Program, Desert Biome. 15 p.

Hutchings, S.S. and G. Stewart. 1953. Increasing forage yields and sheep production on Intermountain winter ranges. Circular No. 925. Washington, DC: U.S. Department of Agriculture. 63 p.

Johnson, K.L. 1978. Wyoming shrublands: Proceedings, 7th Wyoming shrub ecology workshop; 1978 May 31-June 1; Rock Springs, WY. Laramie, WY: University of Wyoming, Agricultural Extension Service. 58 p.

Knapp, P.A. 1998. Spatio-temporal patterns of large grassland fires in the Intermountain West, U.S.A. *Global Ecology and Biogeography Letters* 7(4):259-273.

Nelson, C.R., B.A. Haws and D.L. Nelson. 1990b. Mealybugs and related homoptera of shadscale: possible agents in the dieoff problem in the intermountain west. Pages 152–165 in Symposium on cheatgrass invasion, shrub die-off and other aspects of shrub biology and management. USDA Forest Service GTR-INT-276.

Nelson, D. L., D. J. Weber, and S. C. Garvin. 1990a. The possible role of plant disease in the recent wildland shrub dieoff in Utah. Pages 84–90 in Symposium on cheatgrass invasion, shrub die-off and other aspects of shrub biology and management. USDA Forest Service GTR-INT-276.

Pearson, L.C. 1964. Effect of harvest date on recovery of range grasses and shrubs. *Agronomy Journal* 56:80-82.

Pearson, L.C. 1965. Primary production in grazed and ungrazed desert communities of eastern Idaho. *Ecology* 46(3):278-285.

Peters, E. F. and S. C. Bunting. 1994. Fire conditions pre- and post-occurrence of annual grasses on the Snake River Plain. In: Monsen, S.B. and S.G. Kitchen (compilers). *Proceedings--ecology and management of annual rangelands*; 1992 May 18-22; Boise, ID. Gen. Tech. Rep. INT-GTR-313. Ogden, UT: U.S. Department of Agriculture, Forest Service, Intermountain Research Station: 31-36.

Quinones, F.A. 1981. Indian ricegrass evaluation and breeding. Bulletin 681. Las Cruces, NM: New Mexico State University, Agricultural Experiment Station. 19 p.

Sanderson, S. C., H.C. Stutz, and E.D. McArthur. 1990. Geographic differentiation in *Atriplex confertifolia*. *American Journal of Botany* 77(4):490-498.

Sharp, L.A., K. Sanders and N. Rimbey. 1990. Forty years of change in a shadscale stand in Idaho. *Rangelands* 12(6): 313-328.

Smith, S.D. and P.S. Nobel. 1986. Deserts. In: Baker, N.R. and S.P. Long (eds.). *Photosynthesis in contrasting environments*.

Amsterdam, The Netherlands: Elsevier Science Publishers: Pgs 13-62.

Stubbendieck, J., S.L. Hatch, and C.H. Butterfield. 1992. North American range plants. 4th ed. Lincoln, NE: University of Nebraska Press. 493 p.

USDA-SCS. 1993. Determining success of forage production seedings. TN- Plant Materials No. 32. Reno NV.

Vallentine, John F. 1989. Range Development and Improvements. Third Ed. Academic Press, Inc. San Diego, CA. p. 524.

Weber, D.J., D.L. Nelson, W.H. Hess and R.B. Bhat. 1990. Salinity and moisture stress in relation to dieoff of wildland shrubs. Pages 91-102 in Symposium on cheatgrass invasion, shrub die-off and other aspects of shrub biology and management. USDA Forest Service GTR-INT-276.

West, N.E. 1994. Effects of fire on salt-desert shrub rangelands. In: Monsen, S.B. and S.G. Kitchen (compilers). Proceedings--ecology and management of annual rangelands; 1992 May 18-22; Boise, ID. Gen. Tech. Rep. INT-GTR-313. Ogden, UT: U.S. Department of Agriculture, Forest Service, Intermountain Research Station: Pgs 71-74.

Williams, M.C. 1960. Effect of sodium and potassium salts on growth and oxalate content of halogeton. Plant Physiology 35:500-505.

Wood, B.W. and J.D. Brotherson. 1986. Ecological adaptation and grazing response of budsage (*Artemisia spinescens*). In: McArthur, E.D. and B.L. Welch (compilers). Proceedings--symposium on the biology of *Artemisia* and *Chrysothamnus*; 1984 July 9-13; Provo, UT. Gen. Tech. Rep. INT-200. Ogden, UT: U.S. Department of Agriculture, Forest Service, Intermountain Research Station: 75-92.

Wright, H. A. 1971. Why Squirreltail Is More Tolerant to Burning than Needle-and-Thread. Journal of Range Management 24:277-284.

Young, R.P. 1983. Fire as a vegetation management tool in rangelands of the Intermountain Region. In: Monsen, S.B. and N. Shaw (compilers). Managing Intermountain rangelands--improvement of range and wildlife habitats: Proceedings; 1981 September 15-17; Twin Falls, ID; 1982 June 22-24; Elko, NV. Gen. Tech. Rep. INT-157. Ogden, UT: U.S. Department of Agriculture, Forest Service, Intermountain Forest and Range Experiment Station: Pgs 18-31.

Young, J.A. and R.A. Evans 1977. Squirreltail seed germination. Journal of Range Management 30(1):33-36.

Young, J.A. and F. Tipton. 1990. Invasion of cheatgrass into arid environments of the Lahontan Basin. In: McArthur, E.D., E.M. Romney, S.D. Smith, and P.T. Tueller (compilers). Proceedings--symposium on cheatgrass invasion, shrub die-off, and other aspects of shrub biology and management; 1989 April 5-7; Las Vegas, NV. Gen. Tech. Rep. INT-276. Ogden, UT: U.S. Department of Agriculture, Forest Service, Intermountain Research Station: Pgs 37-40.

Zielinski, M.J. 1994. Controlling erosion on lands administered by the Bureau of Land Management, Winnemucca District, Nevada. In: Monsen, S.B. and S.G. Kitchen (compilers). Proceedings--ecology and management of annual rangelands; 1992 May 18-22; Boise, ID. Gen. Tech. Rep. INT-GTR-313. Ogden, UT: U.S. Department of Agriculture, Forest Service, Intermountain Research Station: Pgs 143-146.

Contributors

DK
TK Stringham

Approval

Kendra Moseley, 3/06/2025

Rangeland health reference sheet

Interpreting Indicators of Rangeland Health is a qualitative assessment protocol used to determine ecosystem condition based on benchmark characteristics described in the Reference Sheet. A suite of 17 (or more) indicators are typically considered in an assessment. The ecological site(s) representative of an assessment location must be known prior to applying the protocol and must be verified based on soils and climate. Current plant community cannot be used to identify the ecological site.

Author(s)/participant(s)	Patti Novak-Echenique
Contact for lead author	State Rangeland Management Specialist
Date	03/19/2010
Approved by	
Approval date	

Composition (Indicators 10 and 12) based on

Annual Production

Indicators

1. **Number and extent of rills:** Rills are none.

2. **Presence of water flow patterns:** Water flow patterns are rare to common depending on site location relative to major inflow areas.

3. **Number and height of erosional pedestals or terracettes:** Pedestals are none.

4. **Bare ground from Ecological Site Description or other studies (rock, litter, lichen, moss, plant canopy are not bare ground):**
Bare Ground \pm 80%.

5. **Number of gullies and erosion associated with gullies:** There are typically no gullies.

6. **Extent of wind scoured, blowouts and/or depositional areas:** None

7. **Amount of litter movement (describe size and distance expected to travel):** Fine litter (foliage of grasses and annual & perennial forbs) expected to move distance of slope length during periods of intense summer convection storms or run in of early spring snow melt flows. Persistent litter (large woody material) will remain in place except during unusual flooding (ponding) events.

8. **Soil surface (top few mm) resistance to erosion (stability values are averages - most sites will show a range of values):** Soil stability values will range from 1 to 4. (To be field tested.)

9. **Soil surface structure and SOM content (include type of structure and A-horizon color and thickness):** Structure of soil surface is thin to thick platy. Soil surface colors are light and soils are typified by an ochric epipedon. Organic matter of the surface 2 to 3 inches is typically less than 0.7 percent.

10. **Effect of community phase composition (relative proportion of different functional groups) and spatial distribution on infiltration and runoff:** Shrubs and deep-rooted perennial herbaceous bunchgrasses aid in infiltration. Sparse shrub canopy and associated litter help break raindrop impact.

11. **Presence and thickness of compaction layer (usually none; describe soil profile features which may be mistaken for compaction on this site):** Compacted layers are not typical. Platy, subangular blocky, or massive subsurface layers are normal for this

site and are not to be interpreted as compaction.

12. **Functional/Structural Groups (list in order of descending dominance by above-ground annual-production or live foliar cover using symbols: >>, >, = to indicate much greater than, greater than, and equal to):**

Dominant: Reference Plant Community: Low-stature shrubs or half-shrubs (shadscale)

Sub-dominant: Deep-rooted, cool season, perennial bunchgrasses > shallow-rooted cool season, perennial bunchgrasses > associated shrubs > deep-rooted, cool season, perennial forbs = fibrous, shallow-rooted, cool season, perennial and annual forbs

Other:

Additional:

13. **Amount of plant mortality and decadence (include which functional groups are expected to show mortality or decadence):**
Dead branches within individual shrubs common and standing dead shrub canopy material may be as much as 30% of total woody canopy.
-

14. **Average percent litter cover (%) and depth (in):** Between plant interspaces ($\pm 10\%$) and depth ($\pm \frac{1}{4}$ in.)
-

15. **Expected annual annual-production (this is TOTAL above-ground annual-production, not just forage annual-production):**
For normal or average growing season (March thru May) ± 200 lbs/ac.
-

16. **Potential invasive (including noxious) species (native and non-native). List species which BOTH characterize degraded states and have the potential to become a dominant or co-dominant species on the ecological site if their future establishment and growth is not actively controlled by management interventions. Species that become dominant for only one to several years (e.g., short-term response to drought or wildfire) are not invasive plants. Note that unlike other indicators, we are describing what is NOT expected in the reference state for the ecological site: Increasers include horsebrush, seepweed, and black greasewood. Invaders include annual mustards, annual kochia, Russian thistle, halogeton, knapweeds, and cheatgrass.**
-

17. **Perennial plant reproductive capability: All functional groups should reproduce in average (or normal) and above average growing season years.**
-