

Ecological site R030XB058NV GRANITIC FAN 5-7 P.Z.

Last updated: 2/24/2025
Accessed: 09/16/2026

General information

Provisional. A provisional ecological site description has undergone quality control and quality assurance review. It contains a working state and transition model and enough information to identify the ecological site.

Ecological site concept

This site occurs on fan remnants. Slope gradients of 2 to 8 percent are typical. Elevations are 2500 to 3600 feet. The soils of this site are derived from granite, schist or gneiss parent materials. These soils are shallow to moderately deep to an argillic horizon, a duripan, or a petrocalcic horizon. This is a group concept and provisional STM that also covers R030XB059NV.

Table 1. Dominant plant species

Tree	Not specified
Shrub	(1) <i>Larrea tridentata</i> (2) <i>Ambrosia dumosa</i>
Herbaceous	(1) <i>Muhlenbergia porteri</i> (2) <i>Achnatherum speciosum</i>

Physiographic features

This site occurs on fan remnants. Slope gradients of 2 to 8 percent are typical. Elevations are 2500 to 3600 feet.

Table 2. Representative physiographic features

Landforms	(1) Fan remnant
Flooding frequency	None
Ponding frequency	None
Elevation	760 – 1,100 m
Slope	0 – 10 %
Water table depth	0 cm

Aspect	Aspect is not a significant factor
--------	------------------------------------

Climatic features

The climate associated with this site is arid, characterized by cool, moist winters, and hot, dry summers. Average annual precipitation is 5 to 7 inches. Mean annual air temperature is 62 to 66 degrees F. The average growing season is about 210 to 250 days.

Table 3 Representative climatic features

Frost-free period (average)	250 days
Freeze-free period (average)	
Precipitation total (average)	180 mm

Influencing water features

There are no influencing water features associated with this site.

Soil features

The soils of this site are derived from granite, schist or gneiss parent materials. These soils are shallow to moderately deep to an argillic horizon, a duripan, or a petrocalcic horizon. The soils are well drained, have slow runoff and have moderate permeability. The soil series associated with this site includes; Arizo, Commiski, Crosgrain, Filaree, Lastchance, Orwash, Shamock, and Tenwell.

Table 4. Representative soil features

Surface texture	(1) Extremely gravelly loam (2) Very gravelly fine sandy loam (3) Extremely gravelly fine sandy loam
Family particle size	(1) Loamy
Drainage class	Well drained
Permeability class	Moderate to moderately slow
Soil depth	60 – 150 cm
Surface fragment cover <=3"	40 – 60 %
Surface fragment cover >3"	0 – 10 %

Available water capacity (0-101.6cm)	2.29 – 10.41 cm
Calcium carbonate equivalent (0-101.6cm)	20 – 50 %
Electrical conductivity (0-101.6cm)	Not specified
Sodium adsorption ratio (0-101.6cm)	0 – 10
Soil reaction (1:1 water) (0-101.6cm)	7.9 – 10
Subsurface fragment volume ≤3" (Depth not specified)	30 – 60 %
Subsurface fragment volume >3" (Depth not specified)	0 – 30 %

Ecological dynamics

This ecological site is dominated by creosotebush and white bursage. Ambrosia generally dominates more developed soils with the ability to hold moisture in the upper soil profile. The shallow roots of white bursage are able to effectively use moisture stored in the upper horizons when it is available and survive for extended periods of time when it is not (Hamerlynck et al. 2002). Ambrosia is not able to dominate the deep, weakly developed, coarse textured soils that store water deep in the profile, which is ideal for Larrea. The spatial distribution of soils allows Ambrosia and Larrea to share dominance on these ecological sites. Ambrosia and Larrea can share dominance throughout the successional process, although the relative abundance is likely to change (Marshall 1994). Creosotebush commonly uses white bursage as a nurse plant, young creosotebushes are frequently found rooted beneath mature white bursage plants (Marshall 1995). White bursage is well adapted to the desert environment but prolonged periods of drought will result a reduction of biomass and possibly kill the plant. Under natural conditions, low available fuel and low fire return interval allowed the establishment of long-lived desert perennial communities.

Fine, sandy alluvium on these sites provides material that the wind redistributes to mound-like coppice dunes beneath creosotebush canopies. Heights of coppice dunes increase as a function of creosotebush cover and are tallest on young alluvial surfaces. Moisture absorbed by and stored in the coppice dune enhances plant performance, in turn contributing to the plant's effectiveness as a windbreak allowing for further deposition of eolian sands (McAuliffe et al 2007). Nutrient concentrations in this shrub community are spatially variable. Nutrient resources are concentrated under shrub canopy relative to the interspaces, called islands of fertility (Kieft et al. 1998).

Most fires in the Mojave Desert are infrequent and of low severity because production of annual and perennial herbs seldom provides a fuel load capable of sustaining fire. Fires in creosotebush scrub were an infrequent event in pre-settlement desert habitats, because fine fuels from winter annual plants were probably sparse, only occurring in large amounts during exceptionally wet winters. Fire kills many creosotebush. Creosotebush is poorly adapted to fire because of its limited sprouting ability. Creosotebush survives some fires that burn patchily or are of low severity. Fire generally kills white bursage. Fire top-kills bush muhly and causes at least short-term decline of bush muhly. Recovery time is thought to vary considerably and is dependent on post fire weather and competition. Bush muhly establishes after fire by sprouting from the root crown and from soil-stored seed. Response to a single fire event may be favorable. Bush muhly's

loosely arranged culms burn cooler and transfer less heat to the root crown compared to culms of more compact bunchgrasses. However, bush muhly declines under repeated desert grassland fires enhanced by cheatgrass (*Bromus tectorum*) and red brome (*B. rubens*). Fire top-kills desert needlegrass. Most needlegrasses (*Achnatherum* spp.), especially young plants, are very susceptible to fire damage. Surviving tufts of desert needlegrass probably will sprout. Desert needlegrass has persistent dead leaf bases, which make it susceptible to burning.

State and transition model

Additional community tables

Table 5. Community 1.1 plant community composition

Group	Common Name	Symbol	Scientific Name	Annual Production ()	Foliar Cover (%)
Grass/Grasslike					
1	Primary Perennial Grasses			30-94	
	desert needlegrass	ACSP12	<i>Achnatherum speciosum</i>	17-50	–
	bush muhly	MUPO2	<i>Muhlenbergia porteri</i>	7-27	–
	big galleta	PLRI3	<i>Pleuraphis rigida</i>	7-17	–
2	Secondary Perennial Grasses			1-17	
	Indian ricegrass	ACHY	<i>Achnatherum hymenoides</i>	2-7	–
	threeawn	ARIST	<i>Aristida</i>	2-7	–
3	Annual Grasses			1-17	
Forb					
4	Perennial Forbs			7-27	
	desert globemallow	SPAM2	<i>Sphaeralcea ambigua</i>	2-7	–
	low woollygrass	DAPU7	<i>Dasyochloa pulchella</i>	0-6	–
5	Annual Forbs			1-27	
Shrub/Vine					
6	Primary Shrubs			101-202	
	burrobush	AMDU2	<i>Ambrosia dumosa</i>	50-101	–
	creosote bush	LATR2	<i>Larrea tridentata</i>	50-101	–
	desert globemallow	SPAM2	<i>Sphaeralcea ambigua</i>	0-6	–
7	Secondary Shrubs			17-67	
	brittlebush	ENFA	<i>Encelia farinosa</i>	3-10	–
	Virgin River brittlebush	ENVI	<i>Encelia virginensis</i>	3-10	–
	jointfir	EPHED	<i>Ephedra</i>	3-10	–
	goldenbush	ERICA2	<i>Ericameria</i>	3-10	–
	winterfat	KRLA2	<i>Krascheninnikovia lanata</i>	3-10	–
	water jacket	LYAN	<i>Lycium andersonii</i>	3-10	–
	Mojave yucca	YUSC2	<i>Yucca schidigera</i>	3-10	–

Table 6. Community 2.1 plant community composition

Group	Common Name	Symbol	Scientific Name	Annual Production ()	Foliar Cover (%)
-------	-------------	--------	-----------------	----------------------	------------------

Table 7. Community 2.2 plant community composition

Group	Common Name	Symbol	Scientific Name	Annual Production ()	Foliar Cover (%)
-------	-------------	--------	-----------------	----------------------	------------------

Table 8. Community 2.3 plant community composition

Group	Common Name	Symbol	Scientific Name	Annual Production ()	Foliar Cover (%)
-------	-------------	--------	-----------------	----------------------	------------------

Table 9. Community 2.4 plant community composition

Group	Common Name	Symbol	Scientific Name	Annual Production ()	Foliar Cover (%)
-------	-------------	--------	-----------------	----------------------	------------------

Table 10. Community 3.1 plant community composition

Group	Common Name	Symbol	Scientific Name	Annual Production ()	Foliar Cover (%)
-------	-------------	--------	-----------------	----------------------	------------------

Animal community

Livestock Interpretations: This site has limited value for livestock grazing, due to the low forage production. Grazing management should be keyed to perennial grass or palatable shrub production. White bursage is an important browse species. Browsing pressure on white bursage is particularly heavy during years of low precipitation, when production of winter annuals is low. White bursage is of intermediate forage value. It is fair to good forage for horses and fair to poor for cattle and sheep. However, because there is often little other forage where white bursage grows, it is often highly valuable to browsing animals. Many animals bed in or under creosotebush. Domestic sheep dig shallow beds under creosotebush because it provides the only shade in the desert scrub community. Creosotebush is unpalatable to livestock. Consumption of creosotebush may be fatal to sheep. Young desert needlegrass is palatable to all classes of livestock. Mature herbage is moderately grazed by horses and cattle, but rarely grazed by sheep. Bush muhly is readily eaten by livestock throughout the year when available; however, it is usually not abundant enough to provide much forage. It is grazed heavily in winter when other species become scarce. Because of its branching habit, it is extremely susceptible to heavy grazing. Bush muhly is damaged when continuously grazed to a stubble height of less than 4 inches (10 cm). Big galleta is considered a valuable forage plant for cattle and domestic sheep. Its coarse, rigid culms make it relatively resistant to heavy grazing and trampling. Stocking rates vary over time depending upon season of use, climate variations, site, and previous and current management goals. A safe starting stocking rate is an estimated stocking rate that is fine tuned by the client by adaptive management through the year and from year to year. Wildlife Interpretations: White bursage is an important browse species for wildlife. Many small mammals browse creosotebush or consume its seeds. Desert reptiles and amphibians use creosotebush as a food source and perch site and hibernate or estivate in burrows under creosotebush, avoiding predators and excessive daytime temperatures. Young desert needlegrass is palatable to many species of wildlife. Desert needlegrass produces considerable basal foliage and is good forage while young. Desert bighorn sheep graze desert needlegrass. The palatability of bush muhly for wildlife species is rated fair to poor. In southern Nevada, big galleta is heavily utilized by bighorn sheep and in some blackbrush communities it is referred to as preferred habitat. Mule deer utilize trace amounts of big galleta.

Hydrological functions

Runoff is medium to high. Permeability is moderate to moderately slow.

Recreational uses

Aesthetic value is derived from the diverse floral and faunal composition and the colorful flowering of wild flowers and shrubs during the spring and early summer. This site offers rewarding opportunities to photographers and for nature study. This site is used for camping and hiking and has potential for upland and big game hunting.

Other products

White bursage is a host for sandfood, a parasitic plant with a sweet, succulent, subterranean flowerstalk. Sandfood was a valuable food supply for Native Americans. Creosotebush has been highly valued for its medicinal properties by desert peoples. It has been used to treat at least 14 illnesses. Twigs and leaves may be boiled as tea, steamed, pounded into a powder, pressed into a poultice, or heated into an infusion.

Other information

White bursage may be used to revegetate disturbed sites in southwestern deserts. Creosotebush may be used to rehabilitate disturbed environments in southwestern deserts. Once established, creosotebush may improve sites for annuals that grow under its canopy by trapping fine soil, organic matter, and symbiont propagules. It may also increase water infiltration and storage. Big galleta's clumped growth form stabilizes blowing sand.

Type locality

Location 1: Clark County, NV

Other references

Fire Effects Information System (Online; <http://www.fs.fed.us/database/feis/>).

Hamerlynck, E.P., J.R. McAuliffe, E.V. McDonald and S.D. Smith. 2002. Ecological Responses of Two Mojave Desert Shrubs to Soils Horizon Development and Soil Water Dynamics. *Ecology*. 83.3: 768-779.

McAuliffe, J.R., E.P. Hamerlynck, and M.C. Eppes. 2007. Landscape dynamics fostering the development and persistence of long-lived creosotebush (*Larrea tridentata*) clones in the Mojave Desert. *J. of Arid Environments*. 69:96-126.

USDA-NRCS Plants Database (Online; <http://plants.usda.gov/>).

Contributors

RWA

Approval

Sarah Quistberg, 2/24/2025

Rangeland health reference sheet

Interpreting Indicators of Rangeland Health is a qualitative assessment protocol used to determine ecosystem condition based on benchmark characteristics described in the Reference Sheet. A suite of 17 (or more) indicators are typically considered in an assessment. The ecological site(s) representative of an assessment location must be known prior to applying the protocol and must be verified based on soils and climate. Current plant community cannot be used to identify the ecological site.

Author(s)/participant(s)	GK BRACKLEY
Contact for lead author	State Rangeland Management Specialist
Date	04/01/2010
Approved by	
Approval date	
Composition (Indicators 10 and 12) based on	Annual Production

Indicators

1. **Number and extent of rills:** Rills are none to rare, but may be evident in areas recently subjected to summer convection storms.

2. **Presence of water flow patterns:** Water flow patterns are none to rare. A few water flow patterns may be evident in areas recently subjected to summer convection storms. Where flow patterns are observed, they are short in length and stable.

3. **Number and height of erosional pedestals or terracettes:** Pedestals are none to rare with occurrence typically limited to areas within water flow patterns.

4. Bare ground from Ecological Site Description or other studies (rock, litter, lichen, moss, plant canopy are not bare ground): Bare Ground $\pm 50\%$; surface rock fragments to $\pm 35\%$; shrub canopy to 10% ; basal area for perennial herbaceous plants $\pm 1\%$.
-

5. Number of gullies and erosion associated with gullies: None
-

6. Extent of wind scoured, blowouts and/or depositional areas: None
-

7. Amount of litter movement (describe size and distance expected to travel): Fine litter (foliage from grasses and annual & perennial forbs) expected to move distance of slope length during intense summer convection storms or rapid snowmelt events. Persistent litter (large woody material) will remain in place except during catastrophic events.
-

8. Soil surface (top few mm) resistance to erosion (stability values are averages - most sites will show a range of values): Soil stability values should be 1 to 3 on the coarse soil textures found on this site. (To be field tested.)
-

9. Soil surface structure and SOM content (include type of structure and A-horizon color and thickness): Surface soil structure is typically single grained to medium platy. Soil surface colors are light and typified by an ochric epipedon. Organic matter of the surface 2 to 3 inches is less than 1 percent.
-

10. Effect of community phase composition (relative proportion of different functional groups) and spatial distribution on infiltration and runoff: Shrub canopy and associated litter break raindrop impact.
-

11. Presence and thickness of compaction layer (usually none; describe soil profile features which may be mistaken for compaction on this site): None. Platy or massive sub-surface horizons, subsoil argillic horizons or duripan are not to be interpreted as compacted layers.
-

12. Functional/Structural Groups (list in order of descending dominance by above-ground annual-production or live foliar cover using symbols: >>, >, = to indicate much greater than, greater than, and equal to):

Dominant: Reference Plant Community: Mojave Desert shrubs

Sub-dominant: cool-season, perennial bunchgrasses = warm- season, perennial grasses = deep-rooted, perennial, forbs = fibrous, shallow-rooted, perennial forbs = annual forbs

Other:

Additional:

13. Amount of plant mortality and decadence (include which functional groups are expected to show mortality or decadence):
Dead branches within individual shrubs common and standing dead shrub canopy material may be as much as 25% of total woody canopy; mature bunchgrasses commonly ($\pm 15\%$) have dead centers.
-
14. Average percent litter cover (%) and depth (in): Between plant interspaces (Trace).
-
15. Expected annual annual-production (this is TOTAL above-ground annual-production, not just forage annual-production):
For normal or average growing season ± 300 lbs/ac.
-
16. Potential invasive (including noxious) species (native and non-native). List species which BOTH characterize degraded states and have the potential to become a dominant or co-dominant species on the ecological site if their future establishment and growth is not actively controlled by management interventions. Species that become dominant for only one to several years (e.g., short-term response to drought or wildfire) are not invasive plants. Note that unlike other indicators, we are describing what is NOT expected in the reference state for the ecological site: Invaders on this site include Red brome, filaree, and Mediterranean grass.
-
17. Perennial plant reproductive capability: All functional groups should reproduce in above average growing season years.
-