

Ecological site R030XB141NV GRAVELLY FAN APRON 5-7 PZ

Last updated: 3/10/2025
Accessed: 08/19/2026

General information

Provisional. A provisional ecological site description has undergone quality control and quality assurance review. It contains a working state and transition model and enough information to identify the ecological site.

Ecological site concept

This site occurs on fan aprons. Slopes range from 2 to 8 percent. Elevations range from 1800 to 3800 feet. The soils of this site are very deep, well drained, and formed in alluvium derived from limestone. This site is part of group concept R030XD015CA.

Associated sites

R030XB137NV	GRAVELLY WASH 5-7 P.Z.
R030XB139NV	COBBLY FAN 5-7 P.Z.

Similar sites

R030XB039NV	LIMY FAN 5-7 P.Z. Shrubs minor component, much more productive.
R030XB043NV	CLAYPAN 5-7 P.Z. More productive site. MUPO2 and KRLA2 important species.

Table 1. Dominant plant species

Tree	Not specified
Shrub	(1) <i>Ambrosia dumosa</i>
Herbaceous	(1) <i>Pleuraphis rigida</i>

Physiographic features

This site occurs on fan aprons. Slopes range from 2 to 8 percent. Elevations range from 1800 to 3800 feet.

Table 2. Representative physiographic features

Landforms	(1) Fan apron
Flooding duration	Very brief (4 to 48 hours)
Flooding frequency	Very rare
Elevation	550 – 1,160 m
Slope	0 – 10 %
Aspect	Aspect is not a significant factor

Climatic features

The climate is typical of the Mojave Desert transitional to Sonoran Desert with hot, dry summers and cool, moist winters. Majority of precipitation comes in the winter with a lesser secondary peak in the summer. The average annual precipitation ranges from 5 to 7 inches. Mean annual air temperature is about 64 degrees F. The average frost-free period is 210 to 250 days.

Table 3 Representative climatic features

Frost-free period (average)	250 days
Freeze-free period (average)	
Precipitation total (average)	180 mm

Influencing water features

This site is very rarely flooded during intense summer convective storms.

Soil features

The soils of this site are very deep, well drained, and formed in alluvium derived from limestone. The soil profile is modified with 40 to 70 percent gravels. The soils are rarely flooded, have moderately high saturated hydraulic conductivity, and medium runoff. The soil series correlated to this ecological site includes Elbowcanyon, classified as a loamy-skeletal, carbonatic, thermic Typic Torriorthent.

Table 4. Representative soil features

Parent material	(1) Alluvium – limestone
Surface texture	(1) Very gravelly fine sandy loam
Family particle size	(1) Loamy

Drainage class	Well drained
Permeability class	Moderately rapid
Soil depth	180 – 210 cm
Surface fragment cover <=3"	80 – 100 %
Surface fragment cover >3"	0 – 10 %
Available water capacity (0-101.6cm)	3.18 – 7.8 cm
Calcium carbonate equivalent (0-101.6cm)	30 – 50 %
Electrical conductivity (0-101.6cm)	Not specified
Sodium adsorption ratio (0-101.6cm)	Not specified
Soil reaction (1:1 water) (0-101.6cm)	10 – 8.2
Subsurface fragment volume <=3" (Depth not specified)	60 – 90 %
Subsurface fragment volume >3" (Depth not specified)	0 – 20 %

Ecological dynamics

The topographic position of this site results in increased run-in moisture from the surrounding landscape, increasing the abundance of warm season perennial bunchgrasses. Warm season grasses have higher light and temperature requirements to begin photosynthesis and grow most actively during the summer, when precipitation is lowest. Primary production in this system is largely constrained by drought. High temperatures and small rainfall events allow nutrients to accumulate during extended dry periods when plant and microbial growth is restricted. When precipitation events do occur, they trigger a pulse of biological activities including plant growth and nutrient uptake (Collins et al 2008). The resource pulses influence plant community structure through annual growth and seedling recruitment. Grasses are intensive exploiters; they extract a large portion of their moisture from shallow soil horizons through their dense network of

shallow roots (Burgess 1995). This trait makes grasses very efficient competitors for limited shallow soil moisture, especially during summer precipitation events. The shrubs of this site are extensive exploiters; they have roots systems that penetrate large volumes of soil both shallow and deep layers (Burgess 1995). This allows shrubs to extract moisture from layers that are too deep or distributed too erratically for intensive exploiters.

Vegetation plays an important role in reducing the erodibility of the soil surface. Incorrect management actions may result in reduced vegetative cover and increased soil erosion. Long-term surface disturbance or reoccurring wildfire will reduce native plant cover, plant density, and species diversity of this site. As ecological condition deteriorates big galleta and perennial forbs decrease, short-lived perennials initially increase. Non-native annual grasses and forbs may increase as native perennial vegetation is lost.

Fire Ecology:

Historically this plant community was influenced by infrequent, low severity fires due to low herbaceous production and widely spaced shrubs. Big galleta's clumped rhizomatous growth form and rigid clumps make it relatively resistant to trampling and other disturbance. Big galleta sprouts from the root crown following fire. Litter and old shoots from galleta provide valuable organic matter when incorporated into the soil. White bursage is tolerant of disturbance and is a common pioneer species on disturbed areas in the Mojave Desert. White bursage is generally killed by wildfire. It reproduces both vegetatively and sexually and will quickly recover. Creosotebush reproduces vegetatively and is able to sprout from the root crown following disturbance. Fire generally kills creosotebush. It has limited sprouting ability following low severity fire, damage can vary depending on season of burning. White ratany is generally completely top killed by fire, but readily sprouts from the root crown and quickly recovers.

State and transition model

Additional community tables

Table 5. Community 1.1 plant community composition

Group	Common Name	Symbol	Scientific Name	Annual Production ()	Foliar Cover (%)
Grass/Grasslike					
1	Primary Perennial Grasses			78-98	
	big galleta	PLRI3	<i>Pleuraphis rigida</i>	78-98	–
2	Secondary Perennial Grasses			8-39	
	Indian ricegrass	ACHY	<i>Achnatherum hymenoides</i>	2-12	–
	low woollygrass	DAPU7	<i>Dasyochloa pulchella</i>	2-12	–
	bush muhly	MUPO2	<i>Muhlenbergia porteri</i>	2-12	–
	redroot cryptantha	CRMIM	<i>Cryptantha micrantha</i> var. <i>micrantha</i>	1-4	–
	Fremont's phacelia	PHFR2	<i>Phacelia fremontii</i>	1-4	–
Forb					
2	Annual Forbs			1-8	
	smallflowered milkvetch	ASNU4	<i>Astragalus nuttallianus</i>	2-8	–
	desert marigold	BAMU	<i>Baileya multiradiata</i>	2-8	–
	desert trumpet	ERIN4	<i>Eriogonum inflatum</i>	2-8	–
	desert globemallow	SPAM2	<i>Sphaeralcea ambigua</i>	2-8	–
3	Perennial Forbs			8-39	
Shrub/Vine					
5	Primary Shrubs			122-188	
	burrobush	AMDU2	<i>Ambrosia dumosa</i>	78-118	–
	creosote bush	LATR2	<i>Larrea tridentata</i>	39-59	–
	white ratany	KRGR	<i>Krameria grayi</i>	4-11	–
6	Secondary Shrubs			19-59	
	rayless goldenhead	ACSP	<i>Acamptopappus sphaerocephalus</i>	4-12	–
	Nevada jointfir	EPNE	<i>Ephedra nevadensis</i>	4-12	–
	Eastern Mojave buckwheat	ERFAP	<i>Eriogonum fasciculatum</i> var. <i>polifolium</i>	4-12	–
	spiny hopsage	GRSP	<i>Grayia spinosa</i>	4-12	–
	winterfat	KRLA2	<i>Krascheninnikovia lanata</i>	4-12	–

	water jacket	LYAN	<i>Lycium andersonii</i>	4-12	–
	Joshua tree	YUBR	<i>Yucca brevifolia</i>	4-12	–
	Mojave yucca	YUSC2	<i>Yucca schidigera</i>	4-12	–
	beavertail pricklypear	OPBA2	<i>Opuntia basilaris</i>	1-4	–
	grizzlybear pricklypear	OPPOE	<i>Opuntia polyacantha</i> var. <i>erinacea</i>	1-4	–
	California barrel cactus	FECYC	<i>Ferocactus cylindraceus</i> var. <i>cylindraceus</i>	1-4	–
	Wiggins' cholla	CYEC3	<i>Cylindropuntia echinocarpa</i>	1-4	–

Table 6. Community 1.2 plant community composition

Group	Common Name	Symbol	Scientific Name	Annual Production ()	Foliar Cover (%)
-------	-------------	--------	-----------------	----------------------	------------------

Table 7. Community 2.1 plant community composition

Group	Common Name	Symbol	Scientific Name	Annual Production ()	Foliar Cover (%)
-------	-------------	--------	-----------------	----------------------	------------------

Table 8. Community 2.2 plant community composition

Group	Common Name	Symbol	Scientific Name	Annual Production ()	Foliar Cover (%)
-------	-------------	--------	-----------------	----------------------	------------------

Table 9. Community 2.3 plant community composition

Group	Common Name	Symbol	Scientific Name	Annual Production ()	Foliar Cover (%)
-------	-------------	--------	-----------------	----------------------	------------------

Animal community

Livestock Interpretations: This site is suitable for livestock grazing. Big galleta dominates the site and is considered fair forage for domestic livestock. Other grasses comprise less than ten percent of the total annual production and provide a relatively unimportant grazing resource. White bursage is of intermediate forage value. In areas where little else is available is can be extremely important. White ratany is an important forage for all classes of livestock. White ratany will decrease in response to heavy grazing pressure. Creosotebush is unpalatable to livestock, however, it is used as bedding cover. Stocking rates vary over time depending upon season of use, climatic variations, site and previous and current management goals. A safe starting stocking rate is an estimated stocking rate that is fine tuned by the client by adaptive management through the year and from year to year. Wildlife Interpretations: Many wildlife species find valuable foraging and habitat resources on this ecological site. Big galleta is extensively used by bighorn sheep and is preferred habitat in some areas. White bursage is an important browse species, especially on ranges where little else is available. White ratany is heavily used by mule deer and desert bighorn sheep. Black-tailed jackrabbits rely almost exclusively on white ratany during the winter. The seeds of white ratany comprise up to 5 percent of the diet of quail in some areas. Many species bed under creosotebush. Desert reptiles and amphibians use creosotebush as a food source, a perch site and also burrow under the shrub to avoid high daytime temperatures. Black-tailed jackrabbits heavily depend on creosotebush for forage throughout the year.

Hydrological functions

This ecological site is characterized by medium runoff and moderately rapid permeability. Rills and waterflow patterns are none to rare. Sparse shrub canopy and associated litter provide some protection from raindrop impact. Perennial herbaceous plants slow runoff and increase infiltration.

Recreational uses

Aesthetic value is derived from the diverse floral and faunal composition and the colorful flowering of wild flowers and shrubs during the spring and early summer. This site offers rewarding opportunities for photography and nature study. This site is used for hiking and has potential for upland and big game hunting.

Other products

Native Americans have used white ratany for many medicinal puposes. An infusion of the roots was used to treat skin irritations and sores. Roots were also used to produce a reddish-yellow dye. The flowers were used to treat upset stomach and diarrhea. The stems were dried and ground and applied to sores. Creosotebush is used by Native Americans to treat many illnesses. Twigs and leaves may be boiled as tea, steamed, pounded into a powder, pressed into a poultice, or heated into an infusion.

Type locality

Location 1: Clark County, NV	
Township/Range/Section	T15S R63E S7
UTM zone	N
UTM northing	4057383
UTM easting	681167
Latitude	36° 38'41"
Longitude	114° 58'24"
General legal description	Approximately 5 miles from Wamp Spring in Desert National Wildlife Refuge, west U.S. Highway 93.USGS Arrow Canyon NW 7.5 minute topographic Quadrangle.

Other references

Burgess, T.L. 1995. Desert Grassland, Mixed Shrub Savanna, Shrub Steppe, or Semidesert Scrub? Pp. 31-67 in M.P. McClaran and T.R. Van Devender (eds.), the Desert Grassland. University of Arizona Press, Tucson Arizona.

Collins, S.L., R.L. Sinsabaugh, C. Crenshaw, L. Green, A. Porras-Alfaro, M. Stursova and L.H. Zeglin. 2008. Pulse dynamics and microbial processes in aridland ecosystems. J. of Ecology. 96: 413-420.

Fire Effects Information System (Online; <http://www.fs.fed.us/database/feis/plants/>)
 USDA-NRCS Plants Database (Online; <http://plants.usda.gov>)

Contributors

E. Hourihan
 PNOVAK-ECHENIQUE

Approval

Kendra Moseley, 3/10/2025

Rangeland health reference sheet

Interpreting Indicators of Rangeland Health is a qualitative assessment protocol used to determine ecosystem condition based on benchmark characteristics described in the Reference Sheet. A suite of 17 (or more) indicators are typically considered in an assessment. The ecological site(s) representative of an assessment location must be known prior to applying the protocol and must be verified based on soils and climate. Current plant community cannot be used to identify the ecological site.

Author(s)/participant(s)	P. Novak-Echenique
--------------------------	--------------------

Contact for lead author	State Rangeland Management Specialist
Date	06/29/2011
Approved by	
Approval date	
Composition (Indicators 10 and 12) based on	Annual Production

Indicators

1. **Number and extent of rills:** Rills are none to rare. Surface rock fragments armor the surface.

2. **Presence of water flow patterns:** Water flow patterns none to rare and may be evident in areas recently subject to intense summer rainfall.

3. **Number and height of erosional pedestals or terracettes:** Pedestals are none to rare with occurrence typically limited to areas within water flow patterns.

4. **Bare ground from Ecological Site Description or other studies (rock, litter, lichen, moss, plant canopy are not bare ground):** Bare Ground up to 10%, depending on amount of surface rock fragments.

5. **Number of gullies and erosion associated with gullies:** None.

6. **Extent of wind scoured, blowouts and/or depositional areas:** None.

7. **Amount of litter movement (describe size and distance expected to travel):** Fine litter (foliage from grasses and annual & perennial forbs) expected to move distance of slope length (<10 ft) during intense summer convection storms or rapid snowmelt events. Persistent litter (large woody material) will remain in place except during large rainfall events.

8. **Soil surface (top few mm) resistance to erosion (stability values are averages - most sites will show a range of values):** Soil stability values should be 1 to 3 in interspaces and 3 to 6 under canopy. (To be field tested.)

9. Soil surface structure and SOM content (include type of structure and A-horizon color and thickness): Surface structure is typically moderate, very thick platy. Soil surface colors are yellowish browns and soils are typified by an ochric epipedon. Organic matter of the surface 2 to 3 inches is less than 1 percent.

10. Effect of community phase composition (relative proportion of different functional groups) and spatial distribution on infiltration and runoff: Sparse shrub canopy and associated litter break raindrop impact. Perennial herbaceous plants slow runoff and increase infiltration.

11. Presence and thickness of compaction layer (usually none; describe soil profile features which may be mistaken for compaction on this site): None. Platy or subangular blocky structure is not to be interpreted as compacted layers.

12. Functional/Structural Groups (list in order of descending dominance by above-ground annual-production or live foliar cover using symbols: >>, >, = to indicate much greater than, greater than, and equal to):

Dominant: Reference Plant Community: Mojave Desert shrubs > >

Sub-dominant: deep-rooted, warm-season, bunchgrasses > perennial forbs > annual forbs > deep-rooted, cool-season, bunchgrasses > annual grasses

Other: succulents

Additional:

13. Amount of plant mortality and decadence (include which functional groups are expected to show mortality or decadence): Dead branches within individual shrubs common and standing dead shrub canopy material may be as much as 25% of total woody canopy; mature bunchgrasses commonly (<20%) have dead centers.

14. Average percent litter cover (%) and depth (in): Between and under canopy 20-25% and depth (<¼-inch)

15. Expected annual annual-production (this is TOTAL above-ground annual-production, not just forage annual-production): For normal or average growing season ±350 lbs/ac. Favorable years 550 lbs/ac and unfavorable years 200 lbs/ac.

16. Potential invasive (including noxious) species (native and non-native). List species which BOTH characterize degraded states and have the potential to become a dominant or co-dominant species on the ecological site if their future establishment and growth is not actively controlled by management interventions. Species that become dominant for only one to several years (e.g., short-term response to drought or wildfire) are not invasive plants. Note that unlike other indicators, we are describing what is NOT expected in the reference state for the ecological site: Potential invaders on this site include red brome, redstem filaree, and Mediterranean grass.

17. Perennial plant reproductive capability: All functional groups should reproduce in above average and normal growing season years. Less reproduction will occur in below average precipitation years. Some functional groups may not reproduce during below average

years.
