

# Ecological site R030XD021CA

## Occasionally Flooded, Hyperthermic, Desert Pavement Ephemeral Stream

Last updated: 10/21/2024

Accessed: 08/17/2026

### General information

**Provisional.** A provisional ecological site description has undergone quality control and quality assurance review. It contains a working state and transition model and enough information to identify the ecological site.

### MLRA notes

Major Land Resource Area (MLRA): 030X–Mojave Basin and Range

MLRA statement: Major Land Resource Area (MLRA) 30, Mojave Desert, is found in southern California, southern Nevada, the extreme southwest corner of Utah and northwestern Arizona within the Basin and Range Province of the Intermontane Plateaus. The climate of the area is hot (primarily hyperthermic and thermic; however at higher elevations, generally above 5000 feet, mesic, cryic and frigid) and dry (aridic). Elevations range from below sea level to over 12,000 feet in the higher mountain areas found within the MLRA. Due to the extreme elevational range found within this MLRA, land resource units (LRUs) were designated to group the MLRA into similar land units.

### LRU notes

"XY" Land Resource Unit (LRU): This LRU is found throughout the Mojave Desert MLRA. This LRU designation is set aside for ecological sites that are ubiquitous throughout the MLRA. These sites are driven by environmental or chemical features that override the climatic designations of the other LRU's or are atypical compared to the surrounding landscape. Common overriding XY characteristics within this MLRA include: ephemeral streams subject to flash flood events, riparian areas or other water features, and soils with strong chemical influence (Na, Ca, etc).

### Ecological site concept

This small ephemeral stream is on inset fans, drainageways, and stream terraces on fan remnants which typically have desert pavement surfaces. Soils are hyperthermic, very deep, and have sandy and gravelly textures. Small, narrow drainageways collect runoff from adjacent fan remnants, and are not associated with larger stream systems. Occasional floods have low volume and intensity, and are contained within the drainageway. Creosote bush (*Larrea tridentata*), brittlebush (*Encelia farinosa*), burrobush (*Ambrosia dumosa*), and white ratany (*Krameria grayi*) are important species. Data ranges in the physiographic data, climate data, water features, and soil data sections of this Ecological Site Description are based on major and minor components, since it is often only associated with minor components. This is a group concept and provisional STM that also covers R030XY021CA.

### Associated sites

|             |                                                                                                                                              |
|-------------|----------------------------------------------------------------------------------------------------------------------------------------------|
| R030XD002CA | <b>Desert Pavement</b><br><br>This is a desert pavement ecological site which is adjacent to this site. There is low cover of creosote bush. |
| R030XD006CA | <b>Abandoned Fan</b><br><br>This ecological site is on adjacent alluvial fans with creosote bush.                                            |
| R030XD039CA | <b>Coarse Gravelly Fans</b><br><br>This ecological site is on alluvial fans with creosote bush and brittlebush.                              |

|                    |                                                                                                                                                                                                     |
|--------------------|-----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| <b>R030XD042CA</b> | <p><b>Hyperthermic Shallow To Moderately Deep Fan Remnants</b></p> <p>This ecological site is on fan remnants with shallow to moderately deep soils over a duripan. Sparse creosote is present.</p> |
|--------------------|-----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|

### Similar sites

|                    |                                                                                                                                                                                                                                                                    |
|--------------------|--------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| <b>R040XD021CA</b> | <p><b>Very Gravelly Wash</b></p> <p>This ecological site is on similar drainageways among fan remnants with desert pavement surfaces, but it is in the Sonoran Desert, generally at lower elevations, with higher temperatures, and more summer precipitation.</p> |
|--------------------|--------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|

**Table 1. Dominant plant species**

|            |                                                                        |
|------------|------------------------------------------------------------------------|
| Tree       | Not specified                                                          |
| Shrub      | <p>(1) <i>Encelia farinosa</i></p> <p>(2) <i>Larrea tridentata</i></p> |
| Herbaceous | <p>(1) <i>Cryptantha</i></p> <p>(2) <i>Chamaesyce abramsiana</i></p>   |

### Physiographic features

This site occurs on inset fans, drainageways and stream terraces (rarely on fan aprons) that drain stable fan remnants covered with desert pavement. Elevations range from 570 to 2800 feet, and slopes are typically 2 to 8 percent, but may range from 0 to 15 percent.

**Table 2. Representative physiographic features**

|                    |                                                                       |
|--------------------|-----------------------------------------------------------------------|
| Landforms          | <p>(1) Inset fan</p> <p>(2) Drainageway</p> <p>(3) Stream terrace</p> |
| Flooding duration  | Very brief (4 to 48 hours)                                            |
| Flooding frequency | None to occasional                                                    |
| Elevation          | 170 – 850 m                                                           |
| Slope              | 0 – 20 %                                                              |
| Aspect             | Aspect is not a significant factor                                    |

### Climatic features

The climate of this ecological site is characterized by hot temperatures, aridity, and a bimodal precipitation pattern. Precipitation falls as rain, with 30 percent falling in summer between July and October, and 65 percent falling in winter between November and March. The mean annual air temperature is 68 to 73 degrees, and mean annual precipitation is 3 to 5 inches. The frost free period is 300 to 340 days, and freeze free period is estimated to range from 320 to 360 days.

Maximum and minimum monthly climate data for this ESD were generated by the Climate Summarizer ([http://www.nm.nrcs.usda.gov/technical/handbooks/nrph/Climate\\_Summarizer.xls](http://www.nm.nrcs.usda.gov/technical/handbooks/nrph/Climate_Summarizer.xls)) using data from the following climate stations (results are unweighted averages).

42598, Eagle Mountain, CA (Period of record = 1933 to 2011) [1]

43855, Hayfield Reservoir, CA (Period of record = 1933 to 2011) [1]

049099, Twentynine Palms, California (Period of record = 1935 to 2011) [1]

The data from multiple weather were combined to most accurately reflect the climatic conditions of this ecological site.

Maximum and minimum monthly climate data for this ESD were generated by the Climate Summarizer ([http://www.nm.nrcs.usda.gov/technical/handbooks/nrph/Climate\\_Summarizer.xls](http://www.nm.nrcs.usda.gov/technical/handbooks/nrph/Climate_Summarizer.xls)) using data from the following climate stations (results are weighted averages; numbers in square brackets represent relative weights):

42598, Eagle Mountain, CA (Period of record = 1933 to 2011) [1]

43855, Hayfield Reservoir, CA (Period of record = 1933 to 2011) [1]

049099, Twentynine Palms, California (Period of record = 1935 to 2011) [1]

The data from multiple weather were combined to most accurately reflect the climatic conditions of this ecological site.

**Table 3 Representative climatic features**

|                               |          |
|-------------------------------|----------|
| Frost-free period (average)   | 340 days |
| Freeze-free period (average)  | 360 days |
| Precipitation total (average) | 130 mm   |

## Influencing water features

### Soil features

The soils associated with this ecological site are very deep, excessively drained, and formed in alluvium from granitoid, igneous or mixed parent material. Surface textures are very gravelly loamy sand, gravelly loamy sand, and gravelly sand. Subsurface horizons (1 to 59 inches) are composed of stratified layers of sandy textures with gravelly to extremely gravelly modifiers. Surface rock fragments less than 3 inches range from 34 to 79 percent cover, and fragments greater than 3 inches range from 1 to 20 percent cover. Subsurface percent by volume of rock fragments less than 3 inches ranges from 15 to 60, and greater than 3 inch fragments range from 1 to 10 percent.

The Carrizo soils are Sandy-skeletal, mixed, hyperthermic Typic Torriorthents. This site is associated with a minor component of Cambidic Haplodurids, but data has not been entered in NASIS at time of this writing.

This ecological site has been correlated with the following map units and soil components within the Joshua Tree National Park Soil Survey Area (CA794):

Map unit ID; Map unit name; Component; phase; percent

1540; Carrizo-Russiroks complex, 2 to 8 percent slopes; Carrizo; occasionally flooded, rocky surface; 20

1541; Carrizo-Cambidic Haplodurids association, 4 to 15 percent slopes; Cambidic Haplodurids; rarely flooded; 2

2068; Aquapeak-Carpetflat-Pintobasin complex, 0 to 4 percent slopes; Carrizo; occasionally flooded, rocky surface; 1

2070; Missionsweet-Carpetflat association, 2 to 30 percent slopes; occasionally flooded, rocky surface; 6  
 2076; Oldale-Carrizo complex, 2 to 8 percent slopes; occasionally flooded, rocky surface; 5  
 2077; Oldale-Carrizo association, 2 to 8 percent slopes; Carrizo; occasionally flooded, rocky surface; 10  
 2110; Descent association, 4 to 50 percent slopes; Carrizo; occasionally flooded, rocky surface; 5

**Table 4. Representative soil features**

|                                             |                                                                              |
|---------------------------------------------|------------------------------------------------------------------------------|
| Parent material                             | (1) Alluvium – granite                                                       |
| Surface texture                             | (1) Very gravelly loamy sand<br>(2) Gravelly loamy sand<br>(3) Gravelly sand |
| Family particle size                        | (1) Sandy                                                                    |
| Drainage class                              | Well drained to excessively drained                                          |
| Permeability class                          | Moderately rapid to rapid                                                    |
| Soil depth                                  | 150 cm                                                                       |
| Surface fragment cover <=3"                 | 30 – 80 %                                                                    |
| Surface fragment cover >3"                  | 0 – 20 %                                                                     |
| Available water capacity<br>(0-101.6cm)     | 1.27 – 5.59 cm                                                               |
| Calcium carbonate equivalent<br>(0-101.6cm) | 0 – 10 %                                                                     |
| Electrical conductivity<br>(0-101.6cm)      | Not specified                                                                |
| Sodium adsorption ratio<br>(0-101.6cm)      | Not specified                                                                |
| Soil reaction (1:1 water)<br>(0-101.6cm)    | 6.4 – 8.8                                                                    |

|                                                          |           |
|----------------------------------------------------------|-----------|
| Subsurface fragment volume <=3"<br>(Depth not specified) | 20 – 60 % |
| Subsurface fragment volume >3"<br>(Depth not specified)  | 0 – 10 %  |

## Ecological dynamics

These small ephemeral streams are on inset fans, drainageways, and stream terraces that drain stable fan remnants that typically have desert pavement surfaces. These ephemeral streams form dendritic (branched) drainage patterns as they collect runoff from adjacent fan remnants. Desert pavement is distinguished by almost barren, level surfaces covered with tightly interlocked surface gravels, which are often darkly colored by desert varnish. Ephemeral streams among desert pavement surfaces appear as wavy lines of vegetation among the dark barren surfaces. Desert pavement has near surface soil features that reduce the rate of infiltration, thus excess flow drains across the flat surfaces to the multiple small ephemeral streams that have developed on the fan remnants. Drainageways among desert pavement surfaces receive continuous soil disturbance from runoff events that prevents formation of desert pavement features, such as an interlocking gravelly surface, or a vesicular horizon. Instead, they are composed of very deep sands with very gravelly or extremely gravelly modifiers. Regular flooding of the drainageways flushes the fine eolian dust deposits from the surface and inhibits the development of the vesicular horizon and the associated uplift of gravels. The deep sands allow for quick and deep infiltration of water. Vegetation is dense and diverse in contrast to the barren desert pavement surfaces, because they can access the deeper water. Creosote bush, brittlebush, burrobush, and white ratany are dominant. A diversity of other shrubs and forbs may be present.

The stable gravel surface and the flat topography of the desert pavement are resistant to erosion, and provide little sediment to the streams during flood events. Occasional floods have low volume and intensity, and are contained within the drainageway. These drainageways are in stable positions, and are not inclined to braid or migrate, unless disturbed.

If the desert pavement surface is disturbed, and the protective gravels are removed, surface erosion and gullyng may increase sediment and run off to the adjacent ephemeral streams. Road development and associated ditches alter drainage patterns, which may cause drainageways to become wider, deeper, and braided.

This ecological site is very unlikely to burn, or to burn extensively due to the large expanses of barren desert pavement between the vegetated channels. Non-native species were not recorded in the plot data, but are present in low amounts in the area. However, they will not likely form a continuous fuel load to carry fire.

## State and transition model

Figure 3. R030XY021CA Model

## Additional community tables

Table 5. Community 2.1 plant community composition

| Group             | Common Name            | Symbol | Scientific Name                            | Annual Production (t) | Foliar Cover (%) |
|-------------------|------------------------|--------|--------------------------------------------|-----------------------|------------------|
| <b>Shrub/Vine</b> |                        |        |                                            |                       |                  |
| 1                 | <b>Shrubs</b>          |        |                                            | 67-129                |                  |
|                   | brittlebush            | ENFA   | <a href="#">Encelia farinosa</a>           | 45-95                 | 1-20             |
|                   | creosote bush          | LATR2  | <a href="#">Larrea tridentata</a>          | 22-56                 | 2-20             |
|                   | white ratany           | KRGR   | <a href="#">Krameria grayi</a>             | 2-22                  | 1-6              |
|                   | desertsenna            | SEAR8  | <a href="#">Senna armata</a>               | 0-22                  | 0-2              |
|                   | desert lavender        | HYEM   | <a href="#">Hyptis emoryi</a>              | 0-22                  | 0-2              |
|                   | sweetbush              | BEJU   | <a href="#">Bebbia juncea</a>              | 0-17                  | 0-4              |
|                   | jojoba                 | SICH   | <a href="#">Simmondsia chinensis</a>       | 0-11                  | 0-1              |
|                   | burrobush              | AMDU2  | <a href="#">Ambrosia dumosa</a>            | 0-1                   | 0-3              |
|                   | Wiggins' cholla        | CYEC3  | <a href="#">Cylindropuntia echinocarpa</a> | 0-1                   | 0-1              |
|                   | branched pencil cholla | CYRA9  | <a href="#">Cylindropuntia ramosissima</a> | 0-1                   | 0-1              |

| Grass/Grasslike |                           |       |                               |        |     |
|-----------------|---------------------------|-------|-------------------------------|--------|-----|
| 2               | Perennial Grass           |       |                               | 0-22   |     |
|                 | big galleta               | PLRI3 | <i>Pleuraphis rigida</i>      | 0-22   | 0-2 |
| Forb            |                           |       |                               |        |     |
| 3               | Forbs                     |       |                               | 45-101 |     |
|                 | cryptantha                | CRYPT | <i>Cryptantha</i>             | 28-39  | 0-5 |
|                 | Abrams' sandmat           | CHAB2 | <i>Chamaesyce abramsiana</i>  | 17-28  | 0-1 |
|                 | smooth desertdandelion    | MAGL3 | <i>Malacothrix glabrata</i>   | 11-22  | 0-3 |
|                 | desert Indianwheat        | PLOV  | <i>Plantago ovata</i>         | 4-9    | 0-2 |
|                 | chia                      | SACO6 | <i>Salvia columbariae</i>     | 0-1    | 0-1 |
|                 | Mojave desertstar         | MOBE2 | <i>Monoptilon bellioides</i>  | 0-1    | 0-1 |
|                 | ghost flower              | MOCO  | <i>Mohavea confertiflora</i>  | 0-1    | 0-1 |
|                 | birdcage evening primrose | OEDE2 | <i>Oenothera deltoides</i>    | 0-1    | 0-1 |
|                 | brittle spineflower       | CHBR  | <i>Chorizanthe brevicornu</i> | 0-1    | 0-1 |
|                 | pincushion flower         | CHFR  | <i>Chaenactis fremontii</i>   | 0-1    | 0-1 |
|                 | buckwheat                 | ERIOG | <i>Eriogonum</i>              | 0-1    | 0-1 |

Table 6. Community 2.2 plant community composition

| Group | Common Name | Symbol | Scientific Name | Annual Production () | Foliar Cover (%) |
|-------|-------------|--------|-----------------|----------------------|------------------|
|-------|-------------|--------|-----------------|----------------------|------------------|

Table 7. Community 3.1 plant community composition

| Group | Common Name | Symbol | Scientific Name | Annual Production () | Foliar Cover (%) |
|-------|-------------|--------|-----------------|----------------------|------------------|
|-------|-------------|--------|-----------------|----------------------|------------------|

## Animal community

Creosote bush shrublands provides a home for an abundance of specialist insect species, for example, creosote bush flowers provide nutrition for over twenty species of bees, and the creosote bush grasshopper (*Boottettix argentatus*) feeds solely on creosote leaves (Pavlik 2008). The sparse vegetation of this ecological site does not provide good cover or food for animals. Brittlebush is used as forage by desert bighorn sheep and mule deer.

## Hydrological functions

Ephemeral drainages maintain water quality by allowing energy dissipation from flow events. These systems transport nutrients and sediments, and store sediments and nutrients in deposition zones. Ephemeral drainages provide temporary storage of surface water, and longer duration storage of subsurface water (Levick et al. 2008).

## Recreational uses

This site may be used for hiking and aesthetic enjoyment.

## Other products

Brittlebush has medicinal uses for Native Americans, including as a poultice for pain and for toothaches. Brittlebush resin is used as chewing gum, to fasten arrow points to twigs, to waterproof water bottles, and is melted to make a varnish. Brittlebush twigs were used as kindling for quick fires. <http://herb.umd.umich.edu/herb/search.pl?searchstring=Encelia+farinosa>. Brittlebush resin is burned as incense in churches in Mexico (Tesky 1993). Creosote bush is an important medicinal plant for Native Americans. It has a very wide range of uses from treatment for consumption, bowl complaints, and menstrual cramps, to induce vomiting, relief for arthritis, rheumatism, aching bones and sprains, congestion and cold, as an antiseptic and disinfectant, dandruff, antispasmodic, to induce urination, gonorrhea, and to cancer treatment. (This list is not exhaustive). <http://herb.umd.umich.edu/herb/search.pl?searchstring=Larrea+tridentata>. Creosote bush stems are used to make weapons, digging tools, and basket handles, and creosote gum is used for knife and awl handles. Creosote bush branches are used as thatch in dwelling construction. <http://herb.umd.umich.edu/herb/search.pl?searchstring=Larrea+tridentata>.

## Inventory data references

The following NRCS plots were used to describe this ecological site: J5-M 701-break PIMO-04B

### Type locality

|                                  |                                                                                                              |
|----------------------------------|--------------------------------------------------------------------------------------------------------------|
| Location 1: Riverside County, CA |                                                                                                              |
| UTM zone                         | N                                                                                                            |
| UTM northing                     | 3749208                                                                                                      |
| UTM easting                      | 659021                                                                                                       |
| General legal description        | The type location is west of HW177, in Joshua Tree National Park, about 15 miles north of Desert Center, CA. |

### Other references

Beatley, J. C. 1969. Dependence of desert rodents on winter annuals and precipitation. *Ecology* 50:721-724.

Beatley, J. C. 1974. Effects of rainfall and temperature on the distribution and behavior of *Larrea tridentata* (Creosote-bush) in the Mojave Desert of Nevada. *Ecology* 55:245-261.

Bowers, J. E. 1994. Natural conditions for seedling emergence of three woody species in the northern Sonoran Desert. *Madroño* 41:73-84.

Bowers, J. E. 2005. Effects of drought on shrub survival and longevity in the northern Sonoran Desert. *Journal of the Torrey Botanical Society* 132:421-431.

Ehleringer, J. R. and C. S. Cook. 1990. Characteristics of *Encelia* species differing in leaf reflectance and transpiration rate under common garden conditions. *Oecologia* 82:484-489.

Hereford, R., R. H. Webb, and C. I. Longpre. 2006. Precipitation history and ecosystem response to multidecadal precipitation variability in the Mojave Desert region, 1893-2001. *Journal of Arid Environments* 67:13-34.

Housman, D. C., M. V. Price, and R. A. Redak. 2002. Architecture of coastal and desert *Encelia farinosa* (Asteraceae): consequences of plastic and heritable variation in leaf characteristics. *American Journal of Botany* 89:1303-1310.

Levick, L., J. Fonseca, D. Goodrich, M. Hernandez, D. Semmens, J. Stromberg, R. Leidy, M. Scianni, D. P. Guertin, M. Tluczek, and W. Kepner. 2008. The ecological and hydrological significance of ephemeral and intermittent streams in the arid and semi-arid American Southwest.

Miriti, M. N., S. Rodriguez-Buritica, S. J. Wright, and H. F. Howe. 2007. Episodic death across species of desert shrubs. *Ecology* 88:32-36.

Pavlik, B. M. 2008. *The California Deserts: an ecological rediscovery*. University of California Press, Ltd., Berkeley and Los Angeles, California.

Sandquist, D. R. and J. R. Ehleringer. 1997. Intraspecific variation in leaf pubescence and drought response in *Encelia farinosa* associated with contrasting desert environments. *New Phytologist* 135:635-644.

Sandquist, D. R. and J. R. Ehleringer. 2003. Population- and family-level variation of brittlebush (*Encelia farinosa*, Asteraceae) pubescence: its relation to drought and implications for selection in variable environments. *American Journal of Botany* 90:1481-1486.

Sandquist, J. R. and J. R. Ehleringer. 1996. Potential adaptability and constraints of response to changing climates for *Encelia farinosa* var. *phenicodonta* from southern Baja California, Mexico. *Madroño* 43:465-478.

Sawyer, J. O., T. Keeler-Woolf, and J. M. Evans. 2009. *A manual of California vegetation*. 2nd edition. California Native Plant Society, Sacramento, California.

Tesky, J. L. 1993. *Hymenoclea salsola*. In: Fire Effects Information System. U.S. Department of Agriculture, Forest Service, Rocky Mountain Research Station, Fire Sciences Laboratory.

Webb, R. H., M. B. Muroy, T. C. Esque, D. E. Boyer, L. A. DeFalco, D. F. Haines, D. Oldershaw, S. J. Scoles, K. A. Thomas, J. B. Blainey, and P. A. Medica. 2003. Perennial vegetation data from permanent plots on the Nevada Test Site, Nye County, Nevada. U.S. Geological Society, Tucson, AZ.

Approval

Kendra Moseley, 10/21/2024

Rangeland health reference sheet

**Interpreting Indicators of Rangeland Health** is a qualitative assessment protocol used to determine ecosystem condition based on benchmark characteristics described in the Reference Sheet. A suite of 17 (or more) indicators are typically considered in an assessment. The ecological site(s) representative of an assessment location must be known prior to applying the protocol and must be verified based on soils and climate. Current plant community cannot be used to identify the ecological site.

|                                             |                   |
|---------------------------------------------|-------------------|
| Author(s)/participant(s)                    |                   |
| Contact for lead author                     |                   |
| Date                                        | 08/17/2026        |
| Approved by                                 |                   |
| Approval date                               |                   |
| Composition (Indicators 10 and 12) based on | Annual Production |

Indicators

1. Number and extent of rills:  

---
2. Presence of water flow patterns:  

---
3. Number and height of erosional pedestals or terracettes:  

---
4. Bare ground from Ecological Site Description or other studies (rock, litter, lichen, moss, plant canopy are not bare ground):  

---
5. Number of gullies and erosion associated with gullies:  

---

6. Extent of wind scoured, blowouts and/or depositional areas:

---

7. Amount of litter movement (describe size and distance expected to travel):

---

8. Soil surface (top few mm) resistance to erosion (stability values are averages - most sites will show a range of values):

---

9. Soil surface structure and SOM content (include type of structure and A-horizon color and thickness):

---

10. Effect of community phase composition (relative proportion of different functional groups) and spatial distribution on infiltration and runoff:

---

11. Presence and thickness of compaction layer (usually none; describe soil profile features which may be mistaken for compaction on this site):

---

12. Functional/Structural Groups (list in order of descending dominance by above-ground annual-production or live foliar cover using symbols: >>, >, = to indicate much greater than, greater than, and equal to):

Dominant:

Sub-dominant:

Other:

Additional:

---

13. Amount of plant mortality and decadence (include which functional groups are expected to show mortality or decadence):

---

14. Average percent litter cover (%) and depth ( in):

---

15. Expected annual annual-production (this is TOTAL above-ground annual-production, not just forage annual-production):

---

16. Potential invasive (including noxious) species (native and non-native). List species which BOTH characterize degraded states and have the potential to become a dominant or co-dominant species on the ecological site if their future establishment and growth is not actively controlled by management interventions. Species that become dominant for only one to several years (e.g., short-term response to drought or wildfire) are not invasive plants. Note that unlike other indicators, we are describing what is NOT expected in the reference state for the ecological site:

---

17. Perennial plant reproductive capability:

---