

Ecological site R030XY023NV

SALINE MEADOW

Last updated: 2/24/2025

Accessed: 09/06/2026

General information

Provisional. A provisional ecological site description has undergone quality control and quality assurance review. It contains a working state and transition model and enough information to identify the ecological site.

Ecological site concept

This site occurs on alluvial flats, lake plains, or axial stream floodplains on all exposures. Slopes range from 0 to 4 percent, but slope gradients of 0 to 2 percent are most typical. Elevations are 1000 to 5400 feet. This site falls under group concept R030XB023CA.

Associated sites

R030XY024NV	SALINE BOTTOM
-------------	---------------

Similar sites

R030XY022NV	WET MEADOW more productive site; JUNCU-CAREX codominant species; SPAI minor species
R030XY024NV	SALINE BOTTOM less productive site; ATLE dominant shrub

Table 1. Dominant plant species

Tree	Not specified
Shrub	Not specified
Herbaceous	(1) <i>Sporobolus airoides</i> (2) <i>Juncus</i>

Physiographic features

This site occurs on alluvial flats, lake plains, or axial stream floodplains on all exposures. Slopes range from 0 to 4 percent, but slope gradients of 0 to 2 percent are most typical. Elevations are 1000 to 5400 feet.

Table 2. Representative physiographic features

Landforms	(1) Alluvial flat (2) Flood plain (3) Lake plain
Flooding duration	Very brief (4 to 48 hours) to brief (2 to 7 days)
Flooding frequency	Rare to occasional
Ponding frequency	None
Elevation	310 – 1,650 m
Slope	0 %
Water table depth	100 – 150 cm
Aspect	Aspect is not a significant factor

Climatic features

The climate is hot and arid, with mild winters and very hot summers. Precipitation is greatest in the winter with a lesser secondary peak in summer, typical of the Mojave Desert. Average annual precipitation is 3 to 7 inches. Mean annual air temperature is 55 to 76 degrees F. The average growing season is about 140 to 360 days.

Table 3 Representative climatic features

Frost-free period (average)	360 days
Freeze-free period (average)	
Precipitation total (average)	130 mm

Influencing water features

This site may experience brief ponding in depressional areas and flooding adjacent to perennial streams.

Soil features

The soils are deep to very deep alluvium from mixed sources. These soils are moderately to strongly alkaline, calcareous throughout and poorly drained. Water table depths range from 27 to 48 inches during the summer months. Permeability is slow to moderate, runoff is very slow and there may be some brief ponding in depression areas. Soil series associated with this ecological site include Land, a fine-silty, mixed superactive, thermic Typic Aquisolid and Paradise, a coarse-loamy, thermic Typic Calciaquoll.

Table 4. Representative soil features

Surface texture	(1) Silty clay loam
Family particle size	(1) Loamy
Drainage class	Somewhat poorly drained
Permeability class	Slow
Soil depth	130 – 150 cm
Surface fragment cover <=3"	Not specified
Surface fragment cover >3"	Not specified
Available water capacity (0-101.6cm)	5.61 – 10.11 cm
Calcium carbonate equivalent (0-101.6cm)	10 – 40 %
Electrical conductivity (0-101.6cm)	0 – 10 mmhos/cm
Sodium adsorption ratio (0-101.6cm)	0 – 10
Soil reaction (1:1 water) (0-101.6cm)	7.9 – 10
Subsurface fragment volume <=3" (Depth not specified)	0 – 20 %
Subsurface fragment volume >3" (Depth not specified)	Not specified

Ecological dynamics

The reference plant community is dominated by alkali sacaton, rush and inland saltgrass. Drought and infrequent wildfire are natural disturbances determining long-term plant community dynamics. Plant community composition is also determined by soil salinity. The soil associated with this site is strongly affected by salt in the upper profile limiting the native vegetation to those able to grow well in salt affected soils.

Inappropriate grazing management will allow shrubs, inland saltgrass and unpalatable rushes and sedges increase while, alkali sacaton and other palatable grasses, sedges and rushes decrease. Forbs such as dock and thistle will increase under continuous disturbance. Willow, saltbush, mesquite, Russian olive and saltcedar readily increase and/or invade this site under drought conditions.

Where severe stream entrenchment occurs, the potential for this site is lost due to the change in soil moisture balance. Stream entrenchment lowers the groundwater level, which is important for maintaining the desirable vegetation composition dominated by graminoids. Deep-rooted shrubs and trees tolerate lower groundwater levels and drought conditions better than relatively shallow rooted forbs and grasses.

Fire Ecology:

Historic fire return intervals of adjacent plant communities (creosotebush scrub) ranged from 35 to >100 years. Alkali sacaton is classified as tolerant of, but not resistant to, fire. Top-killing by fire is probably frequent, and the plants can be killed by severe fire. Rush is fire tolerant when dormant and top-killed by fire during the growing season. It establishes after fire through seed and/or lateral spread by rhizomes. Saltgrass rhizomes occur deep in the soil where they are insulated from the heat of most fires. Saltgrass survives fire by sending up new growth from rhizomes. Sedge is top-killed by fire, with rhizomes protected by insulating soil. The rhizomes of sedge species may be killed by high-severity fires that remove most of the soil organic layer. Reestablishment after fire occurs by seed establishment and/or rhizomatous spread.

Shrubs are uncommon on this site, but provide important habitat and browse for wildlife. Mesquite is top-killed by fire but commonly re-sprouts. Willow also sprouts from the root crown following fire and is among the first species to appear. Saltbush can survive low severity fires. Post-fire regeneration commonly occurs through seed, big saltbush is capable of producing an abundant amount of seed.

State and transition model

Additional community tables

Table 5. Community 1.1 plant community composition

Group	Common Name	Symbol	Scientific Name	Annual Production ()	Foliar Cover (%)
Grass/Grasslike					
1	Primary Perennial Grasses/Grasslikes			874-1513	
	alkali sacaton	SPAI	<i>Sporobolus airoides</i>	504-841	
	rush	JUNCU	<i>Juncus</i>	168-336	—
	saltgrass	DISP	<i>Distichlis spicata</i>	168-252	—
	sedge	CAREX	<i>Carex</i>	34-84	—
2	Secondary Perennial Grasses			84-252	
	common reed	PHAU7	<i>Phragmites australis</i>	9-84	—
	alkali cordgrass	SPGR	<i>Spartina gracilis</i>	9-84	—
Forb					
3	Perennial			84-252	
	Mojave thistle	CIMO	<i>Cirsium mohavense</i>	9-50	—
	marsh elder	IVA	<i>Iva</i>	9-50	—
	dock	RUMEX	<i>Rumex</i>	9-50	—
4	Annual			1-50	
Shrub/Vine					
5	Primary Shrubs			1-50	
	big saltbush	ATLE	<i>Atriplex lentiformis</i>	1-34	—
	mesquite	PROSO	<i>Prosopis</i>	1-34	—
	willow	SALIX	<i>Salix</i>	1-34	

Table 6. Community 1.2 plant community composition

Group	Common Name	Symbol	Scientific Name	Annual Production ()	Foliar Cover (%)
-------	-------------	--------	-----------------	----------------------	------------------

Table 7. Community 1.3 plant community composition

Group	Common Name	Symbol	Scientific Name	Annual Production ()	Foliar Cover (%)
-------	-------------	--------	-----------------	----------------------	------------------

Table 8. Community 2.1 plant community composition

Group	Common Name	Symbol	Scientific Name	Annual Production ()	Foliar Cover (%)
-------	-------------	--------	-----------------	----------------------	------------------

Table 9. Community 2.2 plant community composition

Group	Common Name	Symbol	Scientific Name	Annual Production ()	Foliar Cover (%)
-------	-------------	--------	-----------------	----------------------	------------------

Table 10. Community 2.3 plant community composition

Group	Common Name	Symbol	Scientific Name	Annual Production ()	Foliar Cover (%)
-------	-------------	--------	-----------------	----------------------	------------------

Table 11. Community 3.1 plant community composition

Group	Common Name	Symbol	Scientific Name	Annual Production ()	Foliar Cover (%)
-------	-------------	--------	-----------------	----------------------	------------------

Table 12. Community 3.2 plant community composition

Group	Common Name	Symbol	Scientific Name	Annual Production ()	Foliar Cover (%)
-------	-------------	--------	-----------------	----------------------	------------------

Animal community

Livestock Interpretations: This site is suitable for livestock grazing. Grazing management should be keyed to perennial grass production. Alkali sacaton is a valuable forage species in arid and semiarid regions. Plants are tolerant to moderate grazing and can produce abundant herbage utilized by livestock. Rush is described as a fair to good forage species for cattle. On average, rush's palatability is considered medium to moderately low. Rush is considered palatable early in the growing season when plants are young and tender, but as stems mature and toughen palatability declines. Saltgrass's value as forage depends primarily on the relative availability of other grasses of higher nutritional value and palatability. It can be an especially important late summer grass in arid environments after other forage grasses have deceased. Saltgrass is rated as a fair to good forage species only because it stays green after most other grasses dry. Livestock generally avoid saltgrass due to its coarse foliage. Saltgrass is described as an increaser under grazing pressure. Sedge provides good to fair forage for domestic grazing. Stocking rates vary over time depending upon season of use, climate variations, site, and previous and current management goals. A safe starting stocking rate is an estimated stocking rate that is fine tuned by the client by adaptive management through the year and from year to year. Wildlife Interpretations: The western salt desert shrub and grassland communities where alkali sacaton is common support an abundance of mule deer, pronghorn, carnivores, small mammals, birds, amphibians, and reptiles. Rush provides food for several wildlife species and waterfowl. Rush is an important cover species for a variety of small birds, upland game birds, birds of prey, and waterfowl. Saltgrass provides cover for a variety of bird species, small mammals, and arthropods and is on occasion used as forage for several big game wildlife species. Sedges have a high to moderate resource value for elk and a medium value for mule deer. Elk consume beaked sedge later in the growing season.

Hydrological functions

Runoff is low to very low. Permeability is somewhat poorly drained. Rills and waterflow patterns are none. Deep-rooted perennial herbaceous bunchgrasses slow runoff and increase infiltration. Relatively dense foliar cover of perennial grasses and associated litter break raindrop impact and slow overland flow.

Recreational uses

This site offers rewarding opportunities to photographers and for nature study. This site has potential for upland and big game hunting.

Other information

Alkali sacaton is one of the most commonly used species for seeding and stabilizing disturbed lands. Due to alkali sacaton's salt tolerance, is recommended for native grass seeding on subirrigated saline sites. Given its extensive system of rhizomes and roots which form a dense sod, saltgrass is considered a suitable species for controlling wind and water erosion.

Type locality

Location 1: Clark County, NV	
Township/Range/Section	T16S R68E S30
UTM zone	N
UTM northing	730311
UTM easting	4044005
Latitude	36° 30′49″
Longitude	114° 25′41″
General legal description	Along Virgin River near Overton, Clark County, Nevada.
Location 2: Nye County, NV	
Township/Range/Section	T18S R50E S3
UTM zone	N
UTM northing	560182
UTM easting	4030309
Latitude	36° 24′57″
Longitude	116° 19′43″
General legal description	Ash Meadows area, Nye County, Nevada.

Other references

Fire Effects Information System (Online; <http://www.fs.fed.us/database/feis/plants/>).

USDA-NRCS Plants Database (Online; <http://www.plants.usda.gov>).

Contributors

HA

Approval

Sarah Quistberg, 2/24/2025

Rangeland health reference sheet

Interpreting Indicators of Rangeland Health is a qualitative assessment protocol used to determine ecosystem condition based on benchmark characteristics described in the Reference Sheet. A suite of 17 (or more) indicators are typically considered in an assessment. The ecological site(s) representative of an assessment location must be known prior to applying the protocol and must be verified based on soils and climate. Current plant community cannot be used to identify the ecological site.

Author(s)/participant(s)	P.Novak-Echenique/E.Hourihan
Contact for lead author	State Rangeland Management Specialist
Date	10/18/2011
Approved by	
Approval date	
Composition (Indicators 10 and 12) based on	Annual Production

Indicators

1. Number and extent of rills: None

2. Presence of water flow patterns: None

3. Number and height of erosional pedestals or terracettes: None

4. Bare ground from Ecological Site Description or other studies (rock, litter, lichen, moss, plant canopy are not bare ground):
Bare Ground 10-20%

5. Number of gullies and erosion associated with gullies: None

6. Extent of wind scoured, blowouts and/or depositional areas: None

7. Amount of litter movement (describe size and distance expected to travel): Fine litter (foliage of grasses and annual & perennial forbs) only expected to move during periods of flooding by adjacent streams. Persistent litter (large woody material) will remain in place except during peak flooding periods.
-
8. Soil surface (top few mm) resistance to erosion (stability values are averages - most sites will show a range of values): Soil stability values will range from 3 to 6. (To be field tested.)
-
9. Soil surface structure and SOM content (include type of structure and A-horizon color and thickness): Soil surface structure is platy or massive. Soil surface colors are light and soils are characterized by an ochric epipedon. Organic matter can range from 1 to 3 percent in the upper 10 inches.
-
10. Effect of community phase composition (relative proportion of different functional groups) and spatial distribution on infiltration and runoff: Deep-rooted perennial herbaceous bunchgrasses slow runoff and increase infiltration. Relatively dense foliar cover of perennial grasses and associated litter break raindrop impact and slow overland flow.
-
11. Presence and thickness of compaction layer (usually none; describe soil profile features which may be mistaken for compaction on this site): None - Platy or massive subsurface layers are not to be interpreted as compaction.
-
12. Functional/Structural Groups (list in order of descending dominance by above-ground annual-production or live foliar cover using symbols: >>, >, = to indicate much greater than, greater than, and equal to):
- Dominant: Reference Plant Community: Deep-rooted, warm season, perennial grasses >>
- Sub-dominant: rhizomatous, cool season, perennial grasses and grass-like plants > deep-rooted, cool season, perennial forbs > shallow-rooted, cool season, perennial bunchgrasses > fibrous, shallow-rooted, cool season, perennial forbs > shrubs.
- Other: Grass-like plants
- Additional:
-
13. Amount of plant mortality and decadence (include which functional groups are expected to show mortality or decadence): Herbaceous plant mortality or decadence uncommon.
-
14. Average percent litter cover (%) and depth (in): Within plant interspaces ($\pm$ 80%) and depth (1 to 3 in).
-
15. Expected annual annual-production (this is TOTAL above-ground annual-production, not just forage annual-production): For normal or average growing season (September thru July) $\pm$ 1500 lbs/ac; favorable 3000+ lbs/ac; unfavorable 1000 lbs/ac
-

16. **Potential invasive (including noxious) species (native and non-native).** List species which BOTH characterize degraded states and have the potential to become a dominant or co-dominant species on the ecological site if their future establishment and growth is not actively controlled by management interventions. Species that become dominant for only one to several years (e.g., short-term response to drought or wildfire) are not invasive plants. Note that unlike other indicators, we are describing what is NOT expected in the reference state for the ecological site: Potential invaders include saltcedar and Russian olive.
-

17. **Perennial plant reproductive capability:** All functional groups should reproduce in most years.
-