

Ecological site R030XY025NV

SODIC FLAT

Last updated: 2/24/2025
Accessed: 08/27/2026

General information

Provisional. A provisional ecological site description has undergone quality control and quality assurance review. It contains a working state and transition model and enough information to identify the ecological site.

Ecological site concept

This site occurs on the margins of playas, on alluvial flats, and on lake plains. Slopes range from 0 to 4 percent, but slope gradients of 0 to 2 percent are most typical. Elevations are 800 to about 5000 feet. The soils associated with this site are deep alluvium from mixed sources. Surface is very high in salts. Textures vary from loams to clays. Please refer to group concept R030XB025CA to view the provisional STM.

Associated sites

R030XB028NV	VALLEY WASH
R030XY040NV	SODIC TERRACE
R030XY046NV	OUTWASH PLAIN

Similar sites

R030XY046NV	OUTWASH PLAIN ATPO dominant shrub
R030XY040NV	SODIC TERRACE more productive site; LYCIU codominant shrub

Table 1. Dominant plant species

Tree	Not specified
Shrub	(1) <i>Atriplex confertifolia</i> (2) <i>Atriplex hymenelytra</i>
Herbaceous	Not specified

Physiographic features

This site occurs on the margins of playas, on alluvial flats, and on lake plains. Slopes range from 0 to 4 percent, but slope gradients of 0 to 2 percent are most typical. Elevations are 800 to about 5000 feet.

Table 2. Representative physiographic features

Landforms	(1) Alluvial flat
-----------	-------------------

Climatic features

The climate is hot and arid, with mild winters and very hot summers. Precipitation is greatest in the winter with a lesser secondary peak in summer, typical of the Mojave Desert. Average annual precipitation is 3 to about 7 inches. Mean annual air temperature is 55 to 76 degrees F. The average growing season is about 140 to 360 days.

Table 3 Representative climatic features

Frost-free period (average)	360 days
Freeze-free period (average)	
Precipitation total (average)	180 mm

Influencing water features

There are no influencing water features associated with this site.

Soil features

The soils associated with this site are deep alluvium from mixed sources. Surface is very high in salts. Textures vary from loams to clays. A high water table is present. Drainage is very poor. The surface will crust. The soils have low water holding capacity. Potential for sheet and rill erosion is slight to moderate.

Ecological dynamics

Please refer to group concept R030XB025CA to view the provisional STM.

As ecological condition deteriorates, shadscale, desertholly, and cresotebush increase. Species likely to invade this site are mesquite and annuals.

Fire Ecology:

The mean fire return interval for shadscale-greasewood communities ranges from 35 to 100 years. Shadscale communities are usually unaffected by fire because of low fuel loads, although a year of exceptionally heavy winter rains can generate fuels by producing a heavy stand of annual forbs and grasses. Increased presence of non-native annual grasses, such as cheatgrass, can alter fire regimes in shadscale communities by increasing fire frequency under wet to near-normal summer moisture conditions. When fire does occur, the effect on the ecosystem may be extreme. Shadscale is fire intolerant and it does not readily recover from fire, except for establishment through seed. Fall prescribed burning killed 100% of shadscale on study plots in a basin big sagebrush community. Spring burning left a few surviving shadscale plants but greatly reduced shadscale density and frequency.

State and transition model

Additional community tables

Table 4. Community 1.1 plant community composition

Group	Common Name	Symbol	Scientific Name	Annual Production ()	Foliar Cover (%)
Grass/Grasslike					

1	Primary Perennial Grasses			1-9	
	Indian ricegrass	ACHY	<i>Achnatherum hymenoides</i>	1-6	—
	saltgrass	DISP	<i>Distichlis spicata</i>	1-6	—
	alkali sacaton	SPAI	<i>Sporobolus airoides</i>	1-6	—
Forb					
2	Perennial			1-9	
	marsh elder	IVA	<i>Iva</i>	1-3	—
	niterwort	NITRO	<i>Nitrophila</i>	1-3	—
3	Annual			1-3	
Shrub/Vine					
4	Primary Shrubs			64-115	
	shadscale saltbush	ATCO	<i>Atriplex confertifolia</i>	50-67	—
	desertholly	ATHY	<i>Atriplex hymenelytra</i>	11-39	—
	seepweed	SUAED	<i>Suaeda</i>	2-9	—
5	Secondary Shrubs			11-28	
	iodinebush	ALOC2	<i>Allenrolfea occidentalis</i>	1-6	—
	burrobush	AMDU2	<i>Ambrosia dumosa</i>	1-6	—
	fourwing saltbush	ATCA2	<i>Atriplex canescens</i>	1-6	—
	cattle saltbush	ATPO	<i>Atriplex polycarpa</i>	1-6	—
	Torrey's saltbush	ATTO	<i>Atriplex torreyi</i>	1-6	—
	burrobrush	HYSA	<i>Hymenoclea salsola</i>	1-6	—
	alkali goldenbush	ISAC2	<i>Isocoma acradenia</i>	1-6	—
	creosote bush	LATR2	<i>Larrea tridentata</i>	1-6	—
	desert-thorn	LYCIU	<i>Lycium</i>	1-6	—

Animal community

Livestock Interpretations: This site has limited value for livestock grazing due to low forage. Shadscale is a valuable browse species, providing a source of palatable, nutritious forage for a wide variety of livestock. Shadscale provides good browse for domestic sheep. Shadscale leaves and seeds are an important component of domestic sheep and cattle winter diets. Stocking rates vary over time depending upon season of use, climate variations, site, and previous and current management goals. A safe starting stocking rate is an estimated stocking rate that is fine tuned by the client by adaptive management through the year and from year to year. Wildlife Interpretations: Shadscale is a valuable browse species, providing a source of palatable, nutritious forage for a wide variety of wildlife particularly during spring and summer before the hardening of spiny twigs. It supplies browse, seed, and cover for birds, small mammals, rabbits, deer, and pronghorn antelope.

Recreational uses

Aesthetic value is derived from the diverse floral and faunal composition and the colorful flowering of wild flowers and shrubs during the spring and early summer. This site offers rewarding opportunities to photographers and for nature study. This site is used for camping and hiking and has potential for upland and big game hunting.

Other products

Seeds of shadscale were used by Native Americans for bread and mush.

Type locality

Location 1: Clark County, NV	
Township/Range/Section	T25S R63E S10

UTM zone	N
UTM northing	687237
UTM easting	3962341
Latitude	35° 47' 14"
Longitude	114° 55' 41"
General legal description	Eldorado Valley area, Clark County, Nevada. This site also occurs in southern Nye County, Nevada.

Other references

Fire Effects Information System (Online; <http://www.fs.fed.us/database/feis/plants/>).

USDA-NRCS Plants Database (Online; <http://www.plants.usda.gov>).

Contributors

HA

Approval

Sarah Quistberg, 2/24/2025

Rangeland health reference sheet

Interpreting Indicators of Rangeland Health is a qualitative assessment protocol used to determine ecosystem condition based on benchmark characteristics described in the Reference Sheet. A suite of 17 (or more) indicators are typically considered in an assessment. The ecological site(s) representative of an assessment location must be known prior to applying the protocol and must be verified based on soils and climate. Current plant community cannot be used to identify the ecological site.

Author(s)/participant(s)	P NOVAK-ECHENIQUE
Contact for lead author	State Rangeland Management Specialist
Date	07/20/2012
Approved by	
Approval date	

Composition (Indicators 10 and 12) based on	Annual Production
---	-------------------

Indicators

1. **Number and extent of rills: None**

2. **Presence of water flow patterns: Waterflow patterns are rare to common depending on site location relative to major inflow areas. Waterflow patterns are typically short, ending in depressional areas where water ponds.**

3. **Number and height of erosional pedestals or terracettes: None**

4. **Bare ground from Ecological Site Description or other studies (rock, litter, lichen, moss, plant canopy are not bare ground): Bare Ground up to 80%**

5. **Number of gullies and erosion associated with gullies: None**

6. **Extent of wind scoured, blowouts and/or depositional areas: None**

7. **Amount of litter movement (describe size and distance expected to travel): Fine litter (foliage of grasses and annual & perennial forbs) expected to move distance of slope length during periods of intense summer convection storms. Persistent litter (large woody material) will remain in place except during large rainfall events.**

8. **Soil surface (top few mm) resistance to erosion (stability values are averages - most sites will show a range of values): Soil stability values will range from 1 to 4. (To be field tested.)**

9. **Soil surface structure and SOM content (include type of structure and A-horizon color and thickness): Structure of soil surface will be moderate to strong medium platy. The soil surface will crust. Soil surface colors are white or pale browns and soils are typified by a natric horizon near the surface. Organic matter is less than 1.0 percent.**

10. **Effect of community phase composition (relative proportion of different functional groups) and spatial distribution on infiltration and runoff: The soils have slow permeability and brief ponding may occur during the winter months. The surface layer will also crust resulting in poor infiltration. Sparse shrub canopy and associated litter provide little protection from raindrop impact. Runoff is slow to moderately rapid.**

11. Presence and thickness of compaction layer (usually none; describe soil profile features which may be mistaken for compaction on this site): Compacted layers are none. Subsoil natric or argillic horizons are normal for this site and are not to be interpreted as compaction.
-

12. Functional/Structural Groups (list in order of descending dominance by above-ground annual-production or live foliar cover using symbols: >>, >, = to indicate much greater than, greater than, and equal to):

Dominant: salt-desert shrubs

Sub-dominant: deep-rooted, warm season, perennial bunchgrasses > deep-rooted perennial forbs > cool season perennial bunchgrasses > fibrous, shallow-rooted, cool season, perennial and annual forbs

Other: rhizomatous grasses

Additional:

13. Amount of plant mortality and decadence (include which functional groups are expected to show mortality or decadence):

Dead branches within individual shrubs common and standing dead shrub canopy material may be as much as 30% of total woody canopy

14. Average percent litter cover (%) and depth (in): Between plant interspaces and under shrubs up to 5% and depth < ¼ in.
-

15. Expected annual annual-production (this is TOTAL above-ground annual-production, not just forage annual-production):

For normal or average growing season (March thru May) ± 100 lbs/ac; Favorable years 300 lbs/ac and unfavorable years <50 lbs/ac.

16. Potential invasive (including noxious) species (native and non-native). List species which BOTH characterize degraded states and have the potential to become a dominant or co-dominant species on the ecological site if their future establishment and growth is not actively controlled by management interventions. Species that become dominant for only one to several years (e.g., short-term response to drought or wildfire) are not invasive plants. Note that unlike other indicators, we are describing what is NOT expected in the reference state for the ecological site: Potential invaders include: red brome, Mediterranean grass, redstem filaree, annual mustards, Russian thistle.
-

17. Perennial plant reproductive capability: All functional groups should reproduce in average (or normal) and above average growing season years. Little growth or reproduction occurs in drought years.
-