

Ecological site DX032X01B162

Shallow Loamy (SwLy)

Big Horn Basin Rim

Last updated: 3/26/2025

Accessed: 07/11/2026

General information

Provisional. A provisional ecological site description has undergone quality control and quality assurance review. It contains a working state and transition model and enough information to identify the ecological site.

MLRA notes

Major Land Resource Area (MLRA): 032X–Northern Intermountain Desertic Basins

032X – Northern Intermountain Desertic Basins – This MLRA is comprised of two major Basins, the Big Horn and Wind River. These two basins are distinctly different and are split by LRU's to allow individual ESD descriptions. These warm basins are surrounded by uplifts and rimmed by mountains, creating a unique set of plant responses and communities. Unique characteristics of the geology and geomorphology single these two basins out. Further information regarding MLRAs, refer to: United States Department of Agriculture, Natural Resources Conservation Service. 2006. Land Resource Regions and Major Land Resource Areas of the United States, the Caribbean, and the Pacific Basin. U.S. Department of Agriculture Handbook 296. Available electronically at: http://www.nrcs.usda.gov/wps/portal/nrcs/detail/soils/ref/?cid=nrcs142p2_053624#handbook.

LRU notes

Land Resource Unit (LRU): 32X01B (WY): This LRU is the Big Horn Basin within MLRA 32. This LRU is lower in elevation, slightly warmer and receives slightly less overall precipitation than the Wind River Basin (LRU 02). This LRU was originally divided into two LRU's - LRU A which was the core and LRU B which was the rim. With the most current standards, this LRU is divided into two subsets. This subset is Subset B, referred to as the Rim, is a transitional band between the basin floor and the lower foothills. The subset encircles Subset A which was originally LRU A. As the LRU shifts towards the south and tracks east, changes in geology and relation to the mountain position, creates a minor shift in soil chemistry influencing the variety of ecological sites and plant interactions. The extent of soils currently correlated to this ecological site does not fit within the digitized boundary. Many of the noted soils are provisional and will be reviewed and corrected in mapping update projects. Other map units are correlated as small inclusions within other MLRA's/LRU's based on elevation, landform, and biological references. Moisture Regime: Ustic Aridic – Prior to 2012, many of the soils within this group were correlated as Frigid Ustic Aridic or as Mesic Typic Aridic, with few mapped within this cross over zone. As progressive soil survey mapping continues, these “crossover” or transitional areas are being identified and corrected. Temperature Regime: Mesic Dominant Cover: Rangeland, with Saltbush flats the dominant vegetative cover for this LRU/ESD. Representative Value (RV) Effective Precipitation: 10-14 inches (254 – 355 mm) RV Frost-Free Days: 105-125 days

Classification relationships

Relationship to Other Established Classification Systems: National Vegetation Classification System (NVC): 3 Xeromorphic Woodland, Scrub & Herb Vegetation Class 3.B Cool Semi-Desert Scrub & Grassland Subclass 3.B.1 Cool Semi-Desert Scrub & Grassland formation 3.B.1.NE Western North American Cool Semi-Desert Scrub & Grassland Division M169 Great Basin & Intermountain Tall Sagebrush Shrubland & Steppe Macrogroup G302 Artemisia Tridentata - Artemisia tripartita - Purshia tridentata Big Sagebrush Steppe Group CEG001535 - Artemisia tridentata ssp. wyomingensis/Pseudoroegneria spicata Herbaceous Vegetation or CEG001009 - Artemisia tridentata ssp. wyomingensis/Pseudoroegneria spicata Shrubland Ecoregions (EPA): Level I: 10 North American Deserts Level II: 10.1 Cold Deserts Level III: 10.1.18 Wyoming Basin Level IV: 10.1.18.g Big Horn Salt Desert Shrub Basin

Ecological site concept

- Site receives no additional water.
- Slope is 45%
- Soils Characteristics:
 - o Textures range from very fine sandy loam to clay loam in top 4" (10 cm) of mineral soil surface
 - o All subsurface horizons have a weighted average of >18% clay but 35% clay.
 - o Shallow (10-20 in. (25-50 cm))
 - o 10% stone and boulder cover and 10% cobble and gravel cover
 - o Not skeletal (35% rock fragments) within 20" (50 cm) of mineral soil surface
 - o Non-saline, sodic, or saline-sodic

Associated sites

DX032X01B166	<p>Shallow Sandy (SwSy) Big Horn Basin Rim</p> <p>Shallow Sandy sites will occur in association with Shallow Loamy sites along the outcropping of inter-bedded sedimentary bedrock and sandstone, they will also occur on the dipslopes and ridges of escarpments formed by these same parent materials.</p>
DX032X01B112	<p>Gravelly (Gr) Big Horn Basin Rim</p> <p>Gravelly sites will occur along the exposed shoulder and down slope or inward on the landform will find Shallow Loamy Sites.</p>
DX032X01B122	<p>Loamy (Ly) Big Horn Basin Rim</p> <p>Loamy sites will be found down slope or on lower landscape positions along the same inter-bedded sedimentary bedrock or sandstone, they will also occur on the dipslopes and ridges of escarpments formed by these same parent materials.</p>

Similar sites

R032XY262WY	<p>Shallow Loamy (SwLy) 5-9" Wind River Basin Precipitation Zone</p> <p>Shallow Loamy (SwLy) 5-9" Mesic Wind River Basin Ecological Sites have the same ecological site concept as DX032X01A162 but are found in LRU 02 within the Major Land Resource Area 032X. These sites have lower annual production being in a lower precipitation zone.</p>
DX032X01A162	<p>Shallow Loamy (SwLy) Big Horn Basin Core</p> <p>Shallow Loamy (SwLy) Big Horn Basin Core has the same ecological site concept as DX032X01B162 but falls within a lower precipitation zone (5-9</p>

Table 1. Dominant plant species

Tree	Not specified
Shrub	(1) <i>Artemisia tridentata ssp. wyomingensis</i>
Herbaceous	(1) <i>Pseudoroegneria spicata</i> (2) <i>Hesperostipa comata</i>

Legacy ID

R032XB162WY

Physiographic features

Shallow Loamy Ecological Sites occur on steep slopes ranging up to 45 percent. Ridge tops, escarpments and hillsides are the major landforms where this site is found. This site generally comprises a small component of the soils mapped on these landforms. Soil deposition is minimal on steep slopes and soils are less developed and frequently shallow or very shallow to bedrock. The geology and inherent soil chemistry is complex in the Northern Intermountain Desertic Basins. The Shallow Loamy Ecological Site lacks influence from saline/sodic soils and is a key factor to categorize it separately from other shallow sites.

Table 2. Representative physiographic features

Landforms	(1) Intermontane basin > Hill (2) Intermontane basin > Ridge (3) Intermontane basin > Escarpment
Runoff class	Negligible to high
Elevation	1,650 – 2,290 m
Slope	0 – 50 %
Aspect	Aspect is not a significant factor

Climatic features

Annual Precipitation and modeled relative effective annual precipitation ranges from 10 to 14 inches (254 to 355 mm). The normal precipitation pattern shows peaks in May and June and a secondary peak in September. This amounts to about 50 percent of the mean annual precipitation. Much of the moisture that falls in the latter part of the summer is lost by evaporation and much of the moisture that falls during the winter is lost by sublimation. Average snowfall is about 20 inches annually. Wide fluctuations may occur in yearly precipitation and result in more dry years than those with more than normal precipitation.

Temperatures show a wide range between summer and winter and between daily maximums and minimums, due to the high elevation and dry air, which permits rapid incoming and outgoing radiation. Cold air outbreaks from Canada in winter move rapidly from northwest to southeast and account for extreme minimum temperatures. Chinook winds may occur in winter and bring rapid rises in temperature. Extreme storms may occur during the winter, but most severely affect ranch operations during late winter and spring. High winds are generally blocked from the basin by high mountains but can occur in conjunction with an occasional thunderstorm. Growth of native cool-season plants begins about April 1st and continues to about July 1st. Cool weather and moisture in September may produce some green up of cool season plants that will continue to late October.

For detailed information visit the Natural Resources Conservation Service National Water and Climate Center at <http://www.wcc.nrcs.usda.gov/>. The Clark 3NE, Cody, Cody 12SE, Heart Mtn, Powell Fld Stn, Shell 1NE, and Thermopolis 9NE are the representative weather stations. The following graphs and charts are a collective sample representing the averaged normals and 30-year annual rainfall data for the selected weather stations from 1981 to 2010.

Table 3 Representative climatic features

Frost-free period (characteristic range)	90-90 days
Freeze-free period (characteristic range)	110-120 days
Precipitation total (characteristic range)	180-280 mm
Frost-free period (actual range)	80-110 days
Freeze-free period (actual range)	110-130 days
Precipitation total (actual range)	180-310 mm

Frost-free period (average)	90 days
Freeze-free period (average)	120 days
Precipitation total (average)	230 mm

- (1) HEART MTN [USC00484411], Powell, WY
- (2) POWELL FLD STN [USC00487388], Powell, WY
- (3) CLARK 3NE [USC00481775], Powell, WY
- (4) THERMOPOLIS 9NE [USC00488884], Thermopolis, WY
- (5) SHELL 1NE [USC00488124], Shell, WY
- (6) CODY 12SE [USC00481850], Meeteetse, WY
- (7) CODY [USC00481840], Cody, WY

Influencing water features

The characteristics of these upland soils have no influence from groundwater (water table below 60 inches (150 cm)) and have minimal influence from surface water or overland flow. There may be isolated features that are affected by snow pack that persists longer than surrounding areas due to position on the landform (shaded/protected pockets); but overflow is not a suitable fit. No streams are classified within this ecological site.

Wetland description

No associated wetlands.

Soil features

The soils of this site are shallow (less than 20 inches to bedrock) well-drained soils formed in alluvium over residuum or in residuum. These soils have moderately slow to moderate permeability and may occur on all slopes. The bedrock may be any kind which is virtually impenetrable to plant roots, except igneous. The surface soil will have one or more of the following textures: very fine sandy loam, loam, silt loam, sandy clay loam, silty clay loam, and clay loam. Thin ineffectual layers of other textures are disregarded. The soil characteristics having the most influence on the plant community are the shallow depth, and potential for elevated quantities of soluble salts.

Major Soil Series correlated to this site include: Blazon, Stutzman, Dalhquist, Brownsto, Thermopolis, Blazon, Shingle, Pensore, Cragosen, Rootel, Pilotpeak, Asholler, Worf

Table 4. Representative soil features

Parent material	(1) Residuum – sandstone and shale (2) Alluvium – igneous, metamorphic and sedimentary rock
Surface texture	(1) Loam (2) Silt loam (3) Clay loam (4) Sandy clay loam (5) Sandy loam (6) Fine sandy loam

Family particle size	(1) Loamy
Drainage class	Well drained
Permeability class	Moderately slow to moderate
Depth to restrictive layer	30 – 50 cm
Soil depth	30 – 50 cm
Surface fragment cover <=3"	0 – 10 %
Surface fragment cover >3"	0 – 10 %
Available water capacity (0-101.6cm)	1.27 – 8.89 cm
Calcium carbonate equivalent (0-101.6cm)	0 – 10 %
Electrical conductivity (0-101.6cm)	0 – 10 mmhos/cm
Sodium adsorption ratio (0-101.6cm)	Not specified
Soil reaction (1:1 water) (0-101.6cm)	10 – 8.4
Subsurface fragment volume <=3" (Depth not specified)	0 – 20 %

Ecological dynamics

Potential vegetation on this site is dominated by mid cool-season perennial grasses. Other significant vegetation includes green rabbitbrush, big sagebrush, and a variety of forbs. The expected potential composition for this site is about 80 percent grasses, 10 percent forbs and 10 percent woody plants. The composition and production will vary naturally due to historical use, fluctuating precipitation and fire frequency.

As this site deteriorates, species such as blue grama and Wyoming big sagebrush will increase. Plains prickly pear and weedy annuals will invade. Cool-season grasses such as bluebunch wheatgrass, Indian ricegrass, and rhizomatous wheatgrasses will decrease in frequency and production.

The Reference State (description follows the plant community diagram) has been determined by study of rangeland relic areas, or areas protected from excessive disturbance. Trends in plant communities going from heavily grazed areas to lightly grazed areas, seasonal use pastures, and historical accounts have also been used.

The following is a State and Transition Model Diagram that illustrates the common plant communities (states) that can occur on the site and the transitions between these communities. The ecological processes will be discussed in more detail in the plant community narratives following the diagram.

State and transition model

Additional community tables

Table 5. Community 1.1 plant community composition

Group	Common Name	Symbol	Scientific Name	Annual Production ()	Foliar Cover (%)
Grass/Grasslike					
1	bluebunch wheatgrass			84-118	
	Montana wheatgrass	ELAL7	<i>Elymus albicans</i>	140-280	–
	bluebunch wheatgrass	PSSP6	<i>Pseudoroegneria spicata</i>	140-280	–
2	western wheatgrass			50-84	
	needle and thread	HECO26	<i>Hesperostipa comata</i>	28-84	–
3	thickspike			50-84	
	Indian ricegrass	ACHY	<i>Achnatherum hymenoides</i>	28-71	–
4	Indian ricegrass			34-84	
	prairie Junegrass	KOMA	<i>Koeleria macrantha</i>	28-56	–
5	needle and thread			17-34	
	western wheatgrass	PASM	<i>Pascopyrum smithii</i>	28-56	–
Forb					
7				0-34	
	Forb, perennial	2FP	<i>Forb, perennial</i>	0-28	–
	textile onion	ALTE	<i>Allium textile</i>	0-28	–
	rosy pussytoes	ANRO2	<i>Antennaria rosea</i>	0-28	–
	prairie sagewort	ARFR4	<i>Artemisia frigida</i>	0-28	–
	Missouri milkvetch	ASMI10	<i>Astragalus missouriensis</i>	0-28	–
	Indian paintbrush	CASTI2	<i>Castilleja</i>	0-28	–
	little larkspur	DEBI	<i>Delphinium bicolor</i>	0-28	–
	fleabane	ERIGE2	<i>Erigeron</i>	0-28	–
	sulphur-flower buckwheat	ERUM	<i>Eriogonum umbellatum</i>	0-28	–
	beardtongue	PENST	<i>Penstemon</i>	0-28	–
	spiny phlox	PHHO	<i>Phlox hoodii</i>	0-28	–
	scarlet globemallow	SPCO	<i>Sphaeralcea coccinea</i>	0-28	–
	thrift mock goldenweed	STAR10	<i>Stenotus armerioides</i>	0-28	–
Shrub/Vine					
8				0-17	
	Shrub (>.5m)	2SHRUB	<i>Shrub (>.5m)</i>	0-28	–
	black sagebrush	ARNO4	<i>Artemisia nova</i>	0-28	–

	Wyoming big sagebrush	ARTRW8	<i>Artemisia tridentata ssp. wyomingensis</i>	0-28	-
	yellow rabbitbrush	CHVI8	<i>Chrysothamnus viscidiflorus</i>	0-28	-
	winterfat	KRLA2	<i>Krascheninnikovia lanata</i>	0-28	-

Table 6. Community 1.2 plant community composition

Group	Common Name	Symbol	Scientific Name	Annual Production ()	Foliar Cover (%)
-------	-------------	--------	-----------------	----------------------	------------------

Table 7. Community 2.1 plant community composition

Group	Common Name	Symbol	Scientific Name	Annual Production ()	Foliar Cover (%)
-------	-------------	--------	-----------------	----------------------	------------------

Table 8. Community 3.1 plant community composition

Group	Common Name	Symbol	Scientific Name	Annual Production ()	Foliar Cover (%)
-------	-------------	--------	-----------------	----------------------	------------------

Table 9. Community 4.1 plant community composition

Group	Common Name	Symbol	Scientific Name	Annual Production ()	Foliar Cover (%)
-------	-------------	--------	-----------------	----------------------	------------------

Animal community

Animal Community – Wildlife Interpretations 1.1 - Reference Plant Community: Bluebunch Wheatgrass/Rhizomatous Wheatgrasses: The predominance of grasses in this plant community favors grazers and mixed-feeders, such as bison, elk, and antelope. Suitable thermal and escape cover for deer may be limited due to the low quantities of woody plants. However, topographical variations could provide some escape cover. When found adjacent to sagebrush dominated states, this plant community may provide brood rearing/foraging areas for sage grouse, as well as lek sites. Other birds that would frequent this plant community include western meadowlarks, horned larks, and golden eagles. Many grassland obligate small mammals would occur here. 1.2 - Perennial Grass/Mixed Shrub Plant Community: The combination of an overstory of sagebrush and an understory of grasses and forbs provide a very diverse plant community for wildlife. The crowns of sagebrush tend to break up hard crusted snow on winter ranges, so mule deer and antelope may use this state for foraging and cover year-round, as would cottontail and jack rabbits. It provides important winter, nesting, brood-rearing, and foraging habitat for sage grouse. Brewer's sparrows' nest in big sagebrush plants, and hosts of other nesting birds utilize stands in the 20 to 30 percent cover range. 2.1 - Mixed Shrub/Bare Ground Plant Community: This plant community can provide important winter foraging for elk, mule deer and antelope, as sagebrush can approach 15 percent protein and 40 to 60 percent digestibility during that time. This community provides excellent escape and thermal cover for large ungulates, as well as nesting habitat for sage grouse. 3.1 - Blue Grama Sod Plant Community: These communities provide limited foraging for antelope and other grazers. They may be used as a foraging site by sage grouse if proximal to woody cover and if the Reference Plant Community or the Perennial Grass/Mixed Shrub Plant Community is limiting. Generally, these are not target plant communities for wildlife habitat management. 4.1 - Invasive Plant Community: Early spring and fall green-up of cheatgrass provides foraging opportunities for many of our grazers and mixed feeders. However, as the invasive species increase, decreasing the desirable species, the wildlife species benefits are decreased as well. Animal Community – Grazing Interpretations The following table lists suggested stocking rates for cattle under continuous season-long grazing under normal growing conditions. These are conservative estimates that should be used only as guidelines in the initial stages of the conservation planning process. Often, the current plant composition does not entirely match any particular plant community (as described in this ecological site description). Because of this, a field visit is recommended, in all cases, to document plant composition and production. More precise carrying capacity estimates should eventually be calculated using this information along with animal preference data, particularly when grazers other than cattle are involved. Under more intensive grazing management, improved harvest efficiencies can result in an increased carrying capacity. If distribution problems occur, stocking rates must be reduced to maintain plant health and vigor. Plant Community Production Carrying Capacity* (lb./ac) (AUM/ac) Reference 150-375 0.08 Perennial Grass/Mixed Shrub 100-300 0.05 Mixed Shrub/Bare Ground 75-225 0.03 Blue Grama Sod 25-125 *** Invasive ***** * - The Carrying capacity is calculated as the production for a normal year X .25 efficiency factor / 912.5 #/AUM to calculate the AUM's/Acre. Grazing by domestic livestock is one of the major income-producing industries in the area. Rangeland in this area may provide yearlong forage for cattle, sheep, or horses. During the dormant period, the forage for livestock use needs to be supplemented with protein because the quality does not meet minimum livestock requirements.

Hydrological functions

Water is the principal factor limiting forage production on this site. This site is dominated by soils in hydrologic group B and C, with localized areas in hydrologic group D. Infiltration ranges from moderate to moderately rapid. Runoff potential for this site varies from moderate to high depending on soil hydrologic group and ground cover. In many cases, areas with greater than 75 percent ground cover have the greatest potential for high infiltration and lower runoff. An example of an exception would be where short-grasses form a strong sod and dominate the site. Areas where ground cover is less than 50 percent have the greatest potential to have reduced infiltration and higher runoff (refer to Part 630, NRCS National Engineering Handbook for detailed hydrology information). Rills and gullies should not typically be present. Water flow patterns should be barely distinguishable if at all present. Pedestals are only slightly present in association with bunchgrasses such as bluebunch wheatgrass. Litter typically falls in place, and signs of movement are not common.

Chemical and physical crusts are rare to non-existent. Cryptogamic crusts are present, but only cover one to two percent of the soil surface.

Recreational uses

This site provides hunting opportunities for upland game species. The wide varieties of plants which bloom from spring until fall have an aesthetic value that appeals to visitors. Outside of plants, the extent offers a variety of cultural resources to view on the landscape based on the location of many of these sites on higher ground on the benches and fans which also provides a rich source of geology for exploration.

Wood products

No appreciable wood products are present on the site.

Other products

Herbs: The forb species of the Shallow Loamy Ecological Site have medicinal characteristics and have been used by the Native Americans in this area and more recently by the naturopathic profession. Ornamental Species: The forbs commonly found as well as the shrub component of these communities have been used in landscaping and xeriscaping.

Inventory data references

Information presented in the original site description was derived from NRCS inventory data. Field observations from range-trained personnel also were used. Those involved in developing the original site include Chris Krassin, Range Management Specialist, NRCS and Everet Bainter, Range Management Specialist. Other sources used as references include USDA NRCS Water and Climate Center, USDA NRCS National Range and Pasture Handbook, USDI and USDA Interpreting Indicators of Rangeland Health Version 3, and USDA NRCS Soil Surveys from various counties. Those involved in the development of the new concept for the Sandy ecological site include Tricia Hatle, Range Management Specialist, US Department of the Interior-Bureau of Land Management (USDI-BLM); Karen Hepp, Range Management Specialist, USDI-BLM; and Marji Patz, Ecological Site Specialist, NRCS. Quality control and quality assurance completed by NRCS: Dan Mattke, Area Resource Soil Scientist; Daniel Wood, MLRA Soil Survey Leader; John Hartung, Wyoming State Rangeland Management Specialist; James Bauchert, Wyoming State Soil Scientist; and Scott Woodall, Regional Quality Assurance Ecological Site Specialist. For specific data inquiries, contact the Powell, Wyoming Soil Survey Office (USDA-NRCS). Inventory Data References: Ocular field estimations observed by trained personnel were completed at each site. Then sites were selected where a 100-foot tape was stretched, and the following sample procedures were completed by inventory staff. For full sampling protocol and guidelines with forms please refer to the Wyoming ESI Operating Procedures, compiled in 2012 for the Powell and Rock Springs Soil Survey Office, USDA-NRCS. • Double Sampling Production Data (9.6 hoop used to estimate 10 points, clipped a minimum of three of these estimated points, with two 21-foot X 21-foot square extended shrub plots). • Line Point Intercept (overstory and understory captured with soil cover). Height of herbaceous and woody cover is collected every three feet along established transect.) • Continuous Line Intercept (Woody Canopy Cover, with minimum gap of 0.2 foot for all woody species and succulents. Intercept height collected at each measurement.), • Gap Intercept (Basal Gap measured with a minimum gap requirement of 0.7 foot.), • Sample Point (Ten 1-meter square point photographs taken at set distances on transect. Read using the sample point computer program established by the High Plains Agricultural Research Center, WY). • Soil Stability (Slake Test – surface and subsurface samples collected and processed according to the soil stability guidelines provided by the Jornada Research Center, NM.)

Other references

Baker, William L. 2006. Fire and Restoration of Sagebrush Ecosystems. *Wildlife Society Bulletin* 34(1): 177-185.

Bestelmeyer, B., and J.R. Brown. 2005. State-and-transition models 101: a fresh look at vegetation change. *The Quivira Coalition Newsletter*, Vol. 7, No. 3.

Bestelmeyer, B., J.R. Brown, K.M. Havstad, B. Alexander, G. Chavez, and J.E. Herrick. 2003. Development and use of state and transition models for rangelands. *Journal of Range Management* 56(2):114-126.

Bestelmeyer, B., J.E. Herrick, J.R. Brown, D.A. Trujillo, and K.M. Havstad. 2004. Land management in the American Southwest: a state-and-transition approach to ecosystem complexity. *Environmental Management* 34(1):38-51.

Herrick, J.E., J.W. Van Zee, K.M. Havstad, L.M. Burkett, and W.G. Whitford. 2005. Monitoring manual for grassland, shrubland and savanna ecosystems. Volume I Quick Start. USDA - ARS Jornada Experimental Range, Las Cruces, New Mexico.

Herrick, J.E., J.W. Van Zee, K.M. Havstad, L.M. Burkett, and W.G. Whitford. 2005. Monitoring manual for grassland,

shrubland and savanna ecosystems. Volume II: Design, supplementary methods and interpretation. USDA - ARS Jornada Experimental

Range, Las Cruces, New Mexico.

United States Department of Agriculture, Natural Resources Conservation Service. (electronic) National Water and Climate Center. Available online at <http://www.wcc.nrcs.usda.gov/>. Accessed November 2014.

United States Department of Agriculture, Natural Resources Conservation Service. 2009. Plant Guide: Cheatgrass. Prepared by Skinner et al., National Plant Data Center.

Pellant, M., P. Shaver, D. A. Pyke, and J. E. Herrick. 2005. Interpreting indicators of rangeland health. Version 4. Technical Reference 1734-6. USDI-BLM.

Ricketts, M. J., R. S. Noggles, and B. Landgraf-Gibbons. 2004. Pryor Mountain Wild Horse Range Survey and Assessment. USDA-Natural Resources Conservation Service.

Schoeneberger, P.J., D.A. Wysocki, E.C. Benham, and Soil Survey Staff. 2012. Field book for describing and sampling soils, Version 3.0. Natural Resources Conservation Service, National Soil Survey Center, Lincoln, NE .

Stringham, T.K. and W.C. Krueger. 2001. States, transitions, and thresholds: further refinement for rangeland applications. Agricultural Experiment Station, Oregon State University. Special Report 1024.

Stringham, T.K., W.C. Kreuger, and P.L Shaver. 2003. State and transition modeling: an ecological process approach. Journal of Range Management 56(2):106-113.

United States Department of Agriculture. Soil Survey Division Staff. 1993. Soil Survey Manual, United States Department of Agriculture Handbook No. 18, Chapter 3: Examination and Description of Soils. p.192- 196.

United States Department of Agriculture, Natural Resources Conservation Service. 1997. National Range and Pasture Handbook. (<http://www.glti.nrcs.usda.gov/technical/publications/nrph.html>). Accessed October 2014.

Trlica, M.J. 1999. Grass growth and response to grazing. Range . Colorado State University Cooperative Extension, Natural Resource Series. No. 6.108.

U.S. Department of Agriculture, Natural Resources Conservation Service (USDA/NRCS). 2007. The PLANTS Database (<http://plants.usda.gov>). National Plant Data Center.

U.S. Department of Agriculture, Natural Resources Conservation Service (USDA/NRCS), Soil Survey Staff. 2010. Keys to Soil Taxonomy, 11th Edition.

USDA/NRCS Soil survey manuals for various counties within MLRA 32X. Web soil survey is available online at: <https://websoilsurvey.sc.egov.usda.gov/App/HomePage.htm>.

Western Regional Climate Center. 2014. Electronic station metadata. Available online at: <http://www.wrcc.dri.edu/summary/climsmwy.html>.

Contributors

Marji Patz
Daniel Holter

Approval

Kirt Walstad, 3/26/2025

Acknowledgments

Everet Bainter, retired USDA-NRCS State Rangeland Management Specialist

Rangeland health reference sheet

Interpreting Indicators of Rangeland Health is a qualitative assessment protocol used to determine ecosystem condition based on benchmark characteristics described in the Reference Sheet. A suite of 17 (or more) indicators are typically considered in an assessment. The ecological site(s) representative of an assessment location must be known prior to applying the protocol and must be verified based on soils and climate. Current plant community cannot be used to identify the ecological site.

Author(s)/participant(s)	Ray Gullion, E. Bainter
Contact for lead author	ray.gullion@wy.usda.gov or 307-347-2456
Date	05/02/2008
Approved by	
Approval date	
Composition (Indicators 10 and 12) based on	Annual Production

Indicators

1. **Number and extent of rills:** Rare to nonexistent. Where present, short and widely spaced.

2. **Presence of water flow patterns:** Barely observable.

3. **Number and height of erosional pedestals or terracettes:** Rare to nonexistent.

4. **Bare ground from Ecological Site Description or other studies (rock, litter, lichen, moss, plant canopy are not bare ground):**
Bare ground can range from 15-45%.

5. **Number of gullies and erosion associated with gullies:** Active gullies should not be present.

6. **Extent of wind scoured, blowouts and/or depositional areas:** Rare to nonexistent.

7. **Amount of litter movement (describe size and distance expected to travel):** Herbaceous litter expected to move only in small amounts (to leeward side of shrubs) due to wind. Large woody debris from sagebrush will show no movement.

8. **Soil surface (top few mm) resistance to erosion (stability values are averages - most sites will show a range of values):** Soil Stability Index ratings range from 1 (interspaces) to 5 (under plant canopy), but average values should be 3.0 or greater.

9. **Soil surface structure and SOM content (include type of structure and A-horizon color and thickness):** Soil data is limited for this site. Described A-horizons vary from 1-15 inches (3-38 cm) with OM of .5 to 1.5%.

10. **Effect of community phase composition (relative proportion of different functional groups) and spatial distribution on infiltration and runoff:** Plant community consists of 70-85% grasses, 10% forbs, and 5-20% shrubs. Evenly distributed plant canopy (30-60%) and litter plus slow to moderate infiltration rates result in minimal runoff. Basal cover is typically less than 5% for this site and does very little to effect runoff on this site. Surface rock fragments of 10-20% provide stability to the site, but reduce infiltration.

11. **Presence and thickness of compaction layer (usually none; describe soil profile features which may be mistaken for compaction on this site):** None.

12. **Functional/Structural Groups (list in order of descending dominance by above-ground annual-production or live foliar cover using symbols: >>, >, = to indicate much greater than, greater than, and equal to):**

Dominant: Mid-size, cool season bunchgrasses

Sub-dominant: cool season rhizomatous grasses perennial shrubs

Other: perennial forbs short, cool season bunchgrasses

Additional:

13. **Amount of plant mortality and decadence (include which functional groups are expected to show mortality or decadence):** Minimal decadence, typically associated with shrub component.

14. **Average percent litter cover (%) and depth (in):** Litter ranges from 10-25% of total canopy measurement with total litter (including beneath the plant canopy) from 25-65% expected. Herbaceous litter depth typically ranges from 3-10mm. Woody litter can be up to a couple inches (4-6 cm).

15. **Expected annual annual-production (this is TOTAL above-ground annual-production, not just forage annual-production):** English: 350-700 lb/ac (525 lb/ac average); Metric 392-784 kg/ha (588 kg/ha average).

16. **Potential invasive (including noxious) species (native and non-native). List species which BOTH characterize degraded states and have the potential to become a dominant or co-dominant species on the ecological site if their future establishment and growth is not actively controlled by management interventions. Species that become dominant for only one to several years (e.g., short-term response to drought or wildfire) are not invasive plants. Note that unlike other indicators, we are describing what is NOT expected in the reference state for the ecological site:** Bare ground greater than 75% is the most common indicator of a threshold being crossed. Rabbitbrush, big sagebrush, blue grama, Sandberg bluegrass, buckwheat, and phlox are common increasers. Annual weeds such as kochia, mustards, lambsquarter, and Russian thistle are common invasive species in disturbed sites.

17. Perennial plant reproductive capability: All species are capable of reproducing, except in drought years.
