

Ecological site R032XY218WY
Impervious Clay (IC)
5-9" Wind River Basin Precipitation Zone

Accessed: 08/16/2026

General information

Provisional. A provisional ecological site description has undergone quality control and quality assurance review. It contains a working state and transition model and enough information to identify the ecological site.

Associated sites

R032XY204WY	Clayey (Cy) 5-9" Wind River Basin Precipitation Zone
R032XY238WY	Saline Lowland (SL) 5-9" Wind River Basin Precipitation Zone
R032XY240WY	Saline Lowland Drained (SLDr) 5-9" Wind River Basin Precipitation Zone
R032XY244WY	Saline Upland (SU) 5-9" Wind River Basin Precipitation Zone

Table 1. Dominant plant species

Tree	Not specified
Shrub	Not specified
Herbaceous	Not specified

Physiographic features

This site occurs on nearly level up to 30% slopes.

Table 2. Representative physiographic features

Landforms	(1) Hill (2) Ridge (3) Escarpment
Flooding frequency	None

Ponding frequency	None
Elevation	1,370 – 2,010 m
Slope	0 – 30 %
Ponding depth	0 cm
Aspect	Aspect is not a significant factor

Climatic features

Annual precipitation ranges from 5-9 inches per year. The normal precipitation pattern shows peaks in May and June and a secondary peak in September. This amounts to about 50% of the mean annual precipitation. Much of the moisture that falls in the latter part of the summer is lost by evaporation and much of the moisture that falls during the winter is lost by sublimation. Average snowfall is about 20 inches annually. Wide fluctuations may occur in yearly precipitation and result in more dry years than those with more than normal precipitation.

Temperatures show a wide range between summer and winter and between daily maximums and minimums, due to the high elevation and dry air, which permits rapid incoming and outgoing radiation. Cold air outbreaks from Canada in winter move rapidly from northwest to southeast and account for extreme minimum temperatures. Chinook winds may occur in winter and bring rapid rises in temperature. Extreme storms may occur during the winter, but most severely affect ranch operations during late winter and spring.

High winds are generally blocked from the basin by high mountains, but can occur in conjunction with an occasional thunderstorm.

Growth of native cool-season plants begins about April 1 and continues to about July 1. Cool weather and moisture in September may produce some green up of cool season plants that will continue to late October.

The following information is from the "Pavillion" climate station:

Minimum Maximum 5 yrs. out of 10 between

Frost-free period (days): 95 175 May 19 – September 19

Freeze-free period (days): 98 185 May 6 – October 3

Mean Annual Precipitation (inches): 2.50 12.54

Mean annual precipitation: 7.85 inches

Mean annual air temperature: 44.53 F (30.5 F Avg. Min. to 58.5 F Avg. Max.)

For detailed information visit the Natural Resources Conservation Service National Water and Climate Center at <http://www.wcc.nrcs.usda.gov/> website. Other climate station(s) representative of this precipitation zone include "Riverton", "Arminto", and "Lost Cabin".

Table 3 Representative climatic features

Frost-free period (average)	180 days
Freeze-free period (average)	190 days
Precipitation total (average)	230 mm

Influencing water features

Stream Type: None

Soil features

The soils of this site are very shallow (less than 10") to very deep, well to poorly drained soils formed in alluvium or alluvium over residuum. Layers of the soil most influential to the plant community varies from 3 to 6 inches thick. These soils have slow to very slow permeability. The topsoil, except for thin ineffectual layers, will be heavy clays and/or soils that develop large cracks when dry and are very sticky when wet. These sites typically have moderate saline and /or alkaline soils, but high amounts of soluble salt can occur. The soil characteristics having the most influence on plants are the very slow infiltration rate, which reduces the available moisture, and the amount of soluble salts.

Table 4. Representative soil features

Surface texture	(1) Clay (2) Silty clay (3) Clay loam
Family particle size	(1) Clayey
Drainage class	Poorly drained to well drained
Permeability class	Slow to very slow
Soil depth	10 – 150 cm
Surface fragment cover <=3"	Not specified
Surface fragment cover >3"	Not specified
Available water capacity (0-101.6cm)	1.42 – 20 cm
Calcium carbonate equivalent (0-101.6cm)	0 – 20 %
Electrical conductivity (0-101.6cm)	0 – 20 mmhos/cm
Sodium adsorption ratio (0-101.6cm)	0 – 30
Soil reaction (1:1 water) (0-101.6cm)	7.9 – 10

Subsurface fragment volume <=3" (Depth not specified)	Not specified
Subsurface fragment volume >3" (Depth not specified)	Not specified

Ecological dynamics

Ecological Dynamics of the Site:

Potential vegetation on this site is dominated by salt tolerant, drought resistant, mid cool-season perennial grasses and shrubs. The expected potential composition for this site is about 50% grasses, 10% forbs and 40% woody plants. The composition and production will vary naturally due to historical use, fluctuating precipitation and fire frequency.

As this site deteriorates, species such as Sandberg bluegrass and birdfoot sage will increase. Weedy annuals will invade. Cool season grasses such as rhizomatous wheatgrass, bottlebrush squirreltail, and Indian ricegrass will decrease in frequency and production.

The Historic Climax Plant Community (description follows the plant community diagram) has been determined by study of rangeland relic areas, or areas protected from excessive disturbance. Trends in plant communities going from heavily grazed areas to lightly grazed areas, seasonal use pastures, and historical accounts have also been used.

The following is a State and Transition Model Diagram that illustrates the common plant communities (states) that can occur on the site and the transitions between these communities. The ecological processes will be discussed in more detail in the plant community narratives following the diagram.

State and transition model

Additional community tables

Table 5. Community 1.1 plant community composition

Group	Common Name	Symbol	Scientific Name	Annual Production ()	Foliar Cover (%)
Grass/Grasslike					
1				45-78	
	western wheatgrass	PASM	<i>Pascopyrum smithii</i>	45-78	–
2				22-45	
	squirreltail	ELELE	<i>Elymus elymoides ssp. elymoides</i>	22-45	–
3				11-34	
	Indian ricegrass	ACHY	<i>Achnatherum hymenoides</i>	11-34	–
4				11-22	
	Grass, perennial	2GP	<i>Grass, perennial</i>	0-11	–
	blue grama	BOGR2	<i>Bouteloua gracilis</i>	0-11	–
	Sandberg bluegrass	POSE	<i>Poa secunda</i>	0-11	–
Forb					
5				11-22	
	Forb, perennial	2FP	<i>Forb, perennial</i>	0-11	–
	textile onion	ALTE	<i>Allium textile</i>	0-11	–
	small-leaf pussytoes	ANPA4	<i>Antennaria parvifolia</i>	0-11	–
	sulphur-flower buckwheat	ERUM	<i>Eriogonum umbellatum</i>	0-11	–
	cous biscuitroot	LOCO4	<i>Lomatium cous</i>	0-11	–

	leafy wildparsley	MUDI	<i>Musineon divaricatum</i>	0-11	–
	plains pricklypear	OPPO	<i>Opuntia polyacantha</i>	0-11	–
	spiny phlox	PHHO	<i>Phlox hoodii</i>	0-11	–
	scarlet globemallow	SPCO	<i>Sphaeralcea coccinea</i>	0-11	–
	western aster	SYAS3	<i>Symphyotrichum ascendens</i>	0-11	–
	smooth woodyaster	XYGL	<i>Xylorhiza glabriuscula</i>	0-11	–
Shrub/Vine					
6				45-90	
	birdfoot sagebrush	ARPE6	<i>Artemisia pedatifida</i>	45-90	–
7				11-22	
	Gardner's saltbush	ATGA	<i>Atriplex gardneri</i>	11-22	–
8				11-22	
	Shrub (>.5m)	2SHRUB	<i>Shrub (>.5m)</i>	0-11	–
	Wyoming big sagebrush	ARTRW8	<i>Artemisia tridentata ssp. wyomingensis</i>	0-11	–
	yellow rabbitbrush	CHVI8	<i>Chrysothamnus viscidiflorus</i>	0-11	–
	winterfat	KRASC	<i>Krascheninnikovia</i>	0-11	–

Table 6. Community 2.1 plant community composition

Group	Common Name	Symbol	Scientific Name	Annual Production ()	Foliar Cover (%)
-------	-------------	--------	-----------------	----------------------	------------------

Table 7. Community 3.1 plant community composition

Group	Common Name	Symbol	Scientific Name	Annual Production ()	Foliar Cover (%)
-------	-------------	--------	-----------------	----------------------	------------------

Animal community

Animal Community – Wildlife Interpretations Historic Climax Plant Community: The composition of grasses and shrubs in this plant community favors mixed-feeders such as antelope. Because of the low growing shrub component this is not suitable for thermal and escape cover for deer, but may be preferred by antelope or other wildlife. When found adjacent to sagebrush dominated states, this plant community may provide brood rearing/foraging areas for sage grouse, as well as lek sites. Other birds that would frequent this plant community include western meadowlarks, horned larks, and golden eagles. Prairie dogs and other small mammals prefer these areas as taller denser stands of vegetation impedes recognition and escape from predators. Mixed Perennial Grass/Birdfoot Sagebrush: This plant community exhibits a low level of plant species diversity but is an important winter range for antelope. When found adjacent to sagebrush dominated states, this plant community may provide brood rearing/foraging areas for sage grouse, as well as lek sites. Other birds that would frequent this plant community include western meadowlarks, horned larks, and golden eagles. Prairie dogs and other small mammals prefer these areas as taller denser stands of vegetation impedes recognition and escape from predators. Birdfoot Sagebrush/Bare Ground: This plant community has a low level of diversity. Due to the dominance of annual weeds and grasses feed for large mammals is limited. Areas of bare ground may provide leks for birds such as sage grouse or habitat for prairie dogs and other small mammals. Animal Community – Grazing Interpretations The following table lists suggested stocking rates for cattle under continuous season-long grazing under normal growing conditions. These are conservative estimates that should be used only as guidelines in the initial stages of the conservation planning process. Often, the current plant composition does not entirely match any particular plant community (as described in this ecological site description). Because of this, a field visit is recommended, in all cases, to document plant composition and production. More precise carrying capacity estimates should eventually be calculated using this information along with animal preference data, particularly when grazers other than cattle are involved. Under more intensive grazing management, improved harvest efficiencies can result in an increased carrying capacity. If distribution problems occur, stocking rates must be reduced to maintain plant health and vigor. Plant Community Production Carrying Capacity* (lb./ac) (AUM/ac) Historic Climax Plant Community 100-300 .08 Mixed Perennial Grass/Birdfoot Sagebrush 50-225 .06 Birdfoot Sagebrush/Bare Ground 25-100 .02 * - Continuous, season-long grazing by cattle under average growing conditions. Grazing by domestic livestock is one of the major income-producing industries in the area. Rangeland in this area may provide yearlong forage for cattle, sheep, or horses. During the dormant period, the forage for livestock use needs to be supplemented with protein because the quality does not meet minimum livestock requirements.

Hydrological functions

Water is the principal factor limiting forage production on this site. This site is dominated by soils in hydrologic group C and D. Infiltration ranges from slow to very slow. Runoff potential for this site varies from high to very high depending on soil hydrologic group and ground cover. In many cases, areas with greater than 75% ground cover have the greatest potential for high infiltration and lower runoff. An example of an exception would be where short-grasses form a strong sod and dominate the site. Areas where ground cover is less than

50% have the greatest potential to have reduced infiltration and higher runoff (refer to Part 630, NRCS National Engineering Handbook for detailed hydrology information). Rills and gullies should not typically be present. Water flow patterns should be barely distinguishable if at all present. Pedestals are only slightly present in association with bunchgrasses. Litter typically falls in place, and signs of movement are not common. Chemical and physical crusts are rare to non-existent. Cryptogamic crusts are present, but only cover 1-2% of the soil surface.

Recreational uses

This site provides hunting opportunities for upland game species. The wide variety of plants which bloom from spring until fall have an esthetic value that appeals to visitors.

Wood products

No appreciable wood products are present on the site.

Other products

None noted.

Inventory data references

Information presented here has been derived from NRCS inventory data. Field observations from range trained personnel were also used. Those involved in developing this site include: Chris Krassin, Range Management Specialist, NRCS and Everet Bainter, Range Management Specialist, NRCS. Other sources used as references include USDA NRCS Water and Climate Center, USDA NRCS National Range and Pasture Handbook, USDI and USDA Interpreting Indicators of Rangeland Health Version 3, and USDA NRCS Soil Surveys from various counties.

Contributors

C. Krassin

Rangeland health reference sheet

Interpreting Indicators of Rangeland Health is a qualitative assessment protocol used to determine ecosystem condition based on benchmark characteristics described in the Reference Sheet. A suite of 17 (or more) indicators are typically considered in an assessment. The ecological site(s) representative of an assessment location must be known prior to applying the protocol and must be verified based on soils and climate. Current plant community cannot be used to identify the ecological site.

Author(s)/participant(s)	
Contact for lead author	
Date	07/01/2005
Approved by	
Approval date	
Composition (Indicators 10 and 12) based on	Annual Production

Indicators

1. **Number and extent of rills:** Due to the wide slope range associated with this site, the number and extent of rills will vary from none on slope < 9% to common on slopes > 25%

2. **Presence of water flow patterns:** Barely observable Due to the wide slope range associated with this site, water flow patterns vary from barely observable on slopes of < 9% and from broken and irregular in appearance to continuous on slopes > 25%

3. **Number and height of erosional pedestals or terracettes:** Not evident on slopes < 9%. Erosional pedestals will be present with terracettes present at debris dams on slopes >9%.

4. **Bare ground from Ecological Site Description or other studies (rock, litter, lichen, moss, plant canopy are not bare ground):** Bare ground is 25-40% 50-70%, occurring in small openings throughout the site.

5. **Number of gullies and erosion associated with gullies:** Active gullies should not be present. Active gullies restricted to concentrated water flow patterns.

6. **Extent of wind scoured, blowouts and/or depositional areas:** None

7. **Amount of litter movement (describe size and distance expected to travel):** Little to no plant litter movement. Plant litter remains in place and is not moved by erosional forces. Little to no plant litter movement occurs on slopes < 9%. Litter movement does occur on slopes > 25%.

8. **Soil surface (top few mm) resistance to erosion (stability values are averages - most sites will show a range of values):** Plant cover and litter is at 50% 30% or greater of soil surface and maintains soil surface integrity. Soil Stability class is anticipated to be 5 or greater.

9. **Soil surface structure and SOM content (include type of structure and A-horizon color and thickness):** Use Soil Series description for depth and color of A-horizon

10. **Effect of community phase composition (relative proportion of different functional groups) and spatial distribution on infiltration and runoff:** Grass canopy and basal cover should reduce raindrop impact and slow overland flow providing increased time for infiltration to occur Sparse plant canopy, slow infiltration rates, and the high amount of bare ground contribute . Infiltration varies with soil texture from moderately rapid to rapid. very slow to slow infiltration rates, the amount of bare ground, and steepness of slopes results in a naturally high runoff rate on slopes > 25%, even in HCPC.

11. **Presence and thickness of compaction layer (usually none; describe soil profile features which may be mistaken for compaction on this site):** No compaction layer or soil surface crusting should be present.

12. Functional/Structural Groups (list in order of descending dominance by above-ground annual-production or live foliar cover using symbols: >>, >, = to indicate much greater than, greater than, and equal to):

Dominant:

Sub-dominant:

Other:

Additional: Shrubs > Mid stature Grasses > Forbs = short stature grasses

13. Amount of plant mortality and decadence (include which functional groups are expected to show mortality or decadence):
Some plant mortality and decadence is expected
-

14. Average percent litter cover (%) and depth (in): Average litter cover is 10-20% 5-15% with depths of 0.10 to 0.25 inches
-

15. Expected annual annual-production (this is TOTAL above-ground annual-production, not just forage annual-production):
200 lbs/ac
-

16. Potential invasive (including noxious) species (native and non-native). List species which BOTH characterize degraded states and have the potential to become a dominant or co-dominant species on the ecological site if their future establishment and growth is not actively controlled by management interventions. Species that become dominant for only one to several years (e.g., short-term response to drought or wildfire) are not invasive plants. Note that unlike other indicators, we are describing what is NOT expected in the reference state for the ecological site: Birdfoot sagebrush, Sandberg bluegrass, Woody aster, Annuals, Exotics, and Species found on Noxious Weed List
-

17. Perennial plant reproductive capability: May be Limited due to effective moisture and seed to soil contact
-