

Ecological site R032XY246WY
Sands (Sa)
5-9" Wind River Basin Precipitation Zone

Accessed: 08/16/2026

General information

Provisional. A provisional ecological site description has undergone quality control and quality assurance review. It contains a working state and transition model and enough information to identify the ecological site.

Associated sites

R032XY266WY	Shallow Sandy (SwSy) 5-9" Wind River Basin Precipitation Zone
-------------	---

Table 1. Dominant plant species

Tree	Not specified
Shrub	Not specified
Herbaceous	Not specified

Physiographic features

This site occurs on nearly level to 30% slopes.

Table 2. Representative physiographic features

Landforms	(1) Alluvial fan (2) Hill (3) Stream terrace
Flooding frequency	None
Ponding frequency	None
Elevation	1,370 – 2,010 m
Slope	0 – 30 %
Ponding depth	0 cm

Aspect	Aspect is not a significant factor
--------	------------------------------------

Climatic features

Annual precipitation ranges from 5-9 inches per year. The normal precipitation pattern shows peaks in May and June and a secondary peak in September. This amounts to about 50% of the mean annual precipitation. Much of the moisture that falls in the latter part of the summer is lost by evaporation and much of the moisture that falls during the winter is lost by sublimation. Average snowfall is about 20 inches annually. Wide fluctuations may occur in yearly precipitation and result in more dry years than those with more than normal precipitation.

Temperatures show a wide range between summer and winter and between daily maximums and minimums, due to the high elevation and dry air, which permits rapid incoming and outgoing radiation. Cold air outbreaks from Canada in winter move rapidly from northwest to southeast and account for extreme minimum temperatures. Chinook winds may occur in winter and bring rapid rises in temperature. Extreme storms may occur during the winter, but most severely affect ranch operations during late winter and spring.

High winds are generally blocked from the basin by high mountains, but can occur in conjunction with an occasional thunderstorm.

Growth of native cool-season plants begins about April 1 and continues to about July 1. Cool weather and moisture in September may produce some green up of cool season plants that will continue to late October.

The following information is from the "Pavillion" climate station:

Minimum Maximum 5 yrs. out of 10 between

Frost-free period (days): 95 175 May 19 – September 19

Freeze-free period (days): 98 185 May 6 – October 3

Mean Annual Precipitation (inches): 2.50 12.54

Mean annual precipitation: 7.85 inches

Mean annual air temperature: 44.53 F (30.5 F Avg. Min. to 58.5 F Avg. Max.)

For detailed information visit the Natural Resources Conservation Service National Water and Climate Center at <http://www.wcc.nrcs.usda.gov/> website. Other climate station(s) representative of this precipitation zone include "Riverton", "Arminto", and "Lost Cabin".

Table 3 Representative climatic features

Frost-free period (average)	180 days
Freeze-free period (average)	190 days
Precipitation total (average)	230 mm

Influencing water features

Stream Type: None

Soil features

The soils of this site are moderately deep (greater than 20" to bedrock) to very deep, well-drained soils that formed in alluvium, alluvium over residuum, or eolian deposits. These soils have moderate, moderately rapid, or rapid permeability. The surface soil will vary from 3 to 6 inches deep. The coarser topsoils may be included if underlain by finer textured subsoil. The soil characteristic having the most influence on plant community is the light texture which can affect the available moisture.

Major Soil Series correlated to this site include: Tipperary and Tipper

Table 4. Representative soil features

Surface texture	(1) Fine sandy loam (2) Sandy loam (3) Loamy very fine sand
Family particle size	(1) Sandy
Drainage class	Well drained to excessively drained
Permeability class	Moderately rapid to rapid
Soil depth	50 – 150 cm
Surface fragment cover <=3"	Not specified
Surface fragment cover >3"	Not specified
Available water capacity (0-101.6cm)	4.57 – 10.67 cm
Calcium carbonate equivalent (0-101.6cm)	0 – 10 %
Electrical conductivity (0-101.6cm)	Not specified
Sodium adsorption ratio (0-101.6cm)	0 – 10
Soil reaction (1:1 water) (0-101.6cm)	6.6 – 8.4
Subsurface fragment volume <=3" (Depth not specified)	Not specified

Subsurface fragment volume >3" (Depth not specified)	Not specified
---	---------------

Ecological dynamics

Potential vegetation on this site is dominated by tall and mid perennial grasses. Other significant vegetation includes winterfat, fourwing saltbush, and silver and big sagebrush, and a variety of forbs. The expected potential composition for this site is about 75% grasses, 10% forbs and 15% woody plants. The composition and production will vary naturally due to historical use, fluctuating precipitation and fire frequency.

As this site deteriorates, species such as threadleaf sedge, yucca, skunkbush sumac, and big sagebrush will increase. Weedy annuals will invade. Mid-grasses such as needleandthread, prairie sandreed and Indian ricegrass will decrease in frequency and production.

The Historic Climax Plant Community (description follows the plant community diagram) has been determined by study of rangeland relic areas, or areas protected from excessive disturbance. Trends in plant communities going from heavily grazed areas to lightly grazed areas, seasonal use pastures, and historical accounts have also been used.

The following is a State and Transition Model Diagram that illustrates the common plant communities (states) that can occur on the site and the transitions between these communities. The ecological processes will be discussed in more detail in the plant community narratives following the diagram.

Plant Community Narratives

Following are the narratives for each of the described plant communities. These plant communities may not represent every possibility, but they probably are the most prevalent and repeatable plant communities. The plant composition tables shown above have been developed from the best available knowledge at the time of this revision. As more data is collected, some of these plant communities may be revised or removed, and new ones may be added. None of these plant communities should necessarily be thought of as "Desired Plant Communities". According to the USDA NRCS National Range and Pasture Handbook, Desired Plant Communities (DPC's) will be determined by the decision-makers and will meet minimum quality criteria established by the NRCS. The main purpose for including any description of a plant community here is to capture the current knowledge and experience at the time of this revision.

State and transition model

Additional community tables

Table 5. Community 1.1 plant community composition

Group	Common Name	Symbol	Scientific Name	Annual Production ()	Foliar Cover (%)
-------	-------------	--------	-----------------	----------------------	------------------

Table 6. Community 2.1 plant community composition

Group	Common Name	Symbol	Scientific Name	Annual Production ()	Foliar Cover (%)
-------	-------------	--------	-----------------	----------------------	------------------

Table 7. Community 3.1 plant community composition

Group	Common Name	Symbol	Scientific Name	Annual Production ()	Foliar Cover (%)
-------	-------------	--------	-----------------	----------------------	------------------

Table 8. Community 4.1 plant community composition

Group	Common Name	Symbol	Scientific Name	Annual Production ()	Foliar Cover (%)
Grass/Grasslike					
1				168-224	
	needle and thread	HECO26	<i>Hesperostipa comata</i>	168-224	–
2				56-112	
	Indian ricegrass	ACHY	<i>Achnatherum hymenoides</i>	56-112	–
3				28-56	
	western wheatgrass	PASM	<i>Pascopyrum smithii</i>	28-56	–

4				112-168	
	prairie sandreed	CALO	<i>Calamovilfa longifolia</i>	112-168	–
5				28-56	
	sand dropseed	SPCR	<i>Sporobolus cryptandrus</i>	28-56	–
6				28-84	
	Grass, perennial	2GP	<i>Grass, perennial</i>	0-28	–
	threeawn	ARIST	<i>Aristida</i>	0-28	–
	blue grama	BOGR2	<i>Bouteloua gracilis</i>	0-28	–
	threadleaf sedge	CAFI	<i>Carex filifolia</i>	0-28	–
	prairie Junegrass	KOMA	<i>Koeleria macrantha</i>	0-28	–
	Sandberg bluegrass	POSE	<i>Poa secunda</i>	0-28	–
Forb					
7				28-56	
	miner's candle	CRVI4	<i>Cryptantha virgata</i>	0-28	–
	little larkspur	DEBI	<i>Delphinium bicolor</i>	0-28	–
	bigseed biscuitroot	LOMA3	<i>Lomatium macrocarpum</i>	0-28	–
	spiny phlox	PHHO	<i>Phlox hoodii</i>	0-28	–
	Forb, perennial	2FP	<i>Forb, perennial</i>	0-28	–
	textile onion	ALTE	<i>Allium textile</i>	0-28	–
	Franklin's sandwort	ARFR	<i>Arenaria franklinii</i>	0-28	–
	Missouri milkvetch	ASMI10	<i>Astragalus missouriensis</i>	0-28	–
	bastard toadflax	COUM	<i>Comandra umbellata</i>	0-28	–
	miner's candle	CRVI4	<i>Cryptantha virgata</i>	0-28	–
	little larkspur	DEBI	<i>Delphinium bicolor</i>	0-28	–
	parsnipflower buckwheat	ERHE2	<i>Eriogonum heracleoides</i>	0-28	–
	bigseed biscuitroot	LOMA3	<i>Lomatium macrocarpum</i>	0-28	–
	leafy wildparsley	MUDI	<i>Musineon divaricatum</i>	0-28	–
	plains pricklypear	OPPO	<i>Opuntia polyacantha</i>	0-28	–
	white locoweed	OXSES2	<i>Oxytropis sericea var. speciosa</i>	0-28	–
	spiny phlox	PHHO	<i>Phlox hoodii</i>	0-28	–
	lemon scurfpea	PSLA3	<i>Psoralegium lanceolatum</i>	0-28	–
	scarlet globemallow	SPCO	<i>Sphaeralcea coccinea</i>	0-28	–
	smooth woodyaster	XYGL	<i>Xylorhiza glabriuscula</i>	0-28	–
Shrub/Vine					
8				6-28	
	fourwing saltbush	ATCA2	<i>Atriplex canescens</i>	6-28	–
9				6-28	
	winterfat	KRASC	<i>Krascheninnikovia</i>	6-28	–
10				28-84	
	Shrub (>.5m)	2SHRUB	<i>Shrub (>.5m)</i>	0-28	–
	silver sagebrush	ARCA13	<i>Artemisia cana</i>	0-28	–
	Wyoming big sagebrush	ARTRW8	<i>Artemisia tridentata ssp. wyomingensis</i>	0-28	–
	shadscale saltbush	ATCO	<i>Atriplex confertifolia</i>	0-28	–
	yellow rabbitbrush	CHVI8	<i>Chrysothamnus viscidiflorus</i>	0-28	–
	rubber rabbitbrush	ERNA10	<i>Ericameria nauseosa</i>	0-28	–
	skunkbush sumac	RHTR	<i>Rhus trilobata</i>	0-28	–

	soapweed yucca	YUGL	<i>Yucca glauca</i>	0-28	–
--	----------------	------	---------------------	------	---

Table 9. Community 5.1 plant community composition

Group	Common Name	Symbol	Scientific Name	Annual Production ()	Foliar Cover (%)
-------	-------------	--------	-----------------	----------------------	------------------

Animal community

Historic Climax Plant Community: The predominance of grasses in this plant community favors grazers and mixed-feeders, such as bison, elk, and antelope. Suitable thermal and escape cover for deer may be limited due to the low quantities of woody plants. However, topographical variations could provide some escape cover. When found adjacent to sagebrush dominated states, this plant community may provide brood rearing/foraging areas for sage grouse. Other birds that would frequent this plant community include western meadowlarks, horned larks, and golden eagles. Many grassland obligate small mammals would occur here. Perennial Grass/Mixed Shrub: These communities provide foraging for antelope and other grazers. They may be used as a foraging site by sage grouse if proximal to woody cover. Indian Ricegrass/Yucca: These communities provide foraging for antelope and other grazers. They may be used as a foraging site by sage grouse if proximal to woody cover. Blowout Dune/Mixed Shrub Plant Community: This plant community can provide winter foraging for mule deer and antelope, as shrubs can approach 15% protein and 40-60% digestibility during that time. This community provides escape cover for large ungulates. Needleandthread/Threadleaf Sedge/Yucca Plant Community: These communities provide foraging for antelope and other grazers. They may be used as a foraging site by sage grouse if proximal to woody cover and if the Historic Climax Plant Community or the Perennial Grass/ Mixed Shrub Plant Community is limiting. Generally, these are not target plant communities for wildlife habitat management. Animal Community – Grazing Interpretations The following table lists suggested stocking rates for cattle under continuous season-long grazing under normal growing conditions. These are conservative estimates that should be used only as guidelines in the initial stages of the conservation planning process. Often, the current plant composition does not entirely match any particular plant community (as described in this ecological site description). Because of this, a field visit is recommended, in all cases, to document plant composition and production. More precise carrying capacity estimates should eventually be calculated using this information along with animal preference data, particularly when grazers other than cattle are involved. Under more intensive grazing management, improved harvest efficiencies can result in an increased carrying capacity. If distribution problems occur, stocking rates must be reduced to maintain plant health and vigor. Plant Community Production Carrying Capacity* (lb./ac) (AUM/ac) Historic Climax Plant Community 350-700 .20 Indian Ricegrass/Yucca 350-700 .20 Perennial Grass/Mixed Shrub 200-525 .16 Blowout Dune/Mixed Shrub 75-300 .05 Needleandthread/Threadleaf Sedge/Yucca 100-325 .10 * - Continuous, season-long grazing by cattle under average growing conditions. Grazing by domestic livestock is one of the major income-producing industries in the area. Rangeland in this area may provide yearlong forage for cattle, sheep, or horses. During the dormant period, the forage for livestock use needs to be supplemented with protein because the quality does not meet minimum livestock requirements.

Hydrological functions

Water is the principal factor limiting forage production on this site. This site is dominated by soils in hydrologic group B, with localized areas in hydrologic group C. Infiltration potential for this site varies from moderately rapid to rapid depending on soil hydrologic group and ground cover. Runoff varies from low to moderate. In many cases, areas with greater than 75% ground cover have the greatest potential for high infiltration and lower runoff. Areas where ground cover is less than 50% have the greatest potential to have reduced infiltration and higher runoff (refer to Part 630, NRCS National Engineering Handbook for detailed hydrology information). Rills and gullies should not typically be present. Water flow patterns should be barely distinguishable if at all present. Pedestals are only slightly present in association with bunchgrasses. Litter typically falls in place, and signs of movement are not common. Chemical and physical crusts are rare to non-existent. Cryptogamic crusts are present, but only cover 1-2% of the soil surface.

Recreational uses

Recreational Uses This site provides hunting opportunities for upland game species. The wide variety of plants which bloom from spring until fall have an esthetic value that appeals to visitors.

Wood products

No appreciable wood products are present on the site.

Other products

None noted.

Inventory data references

Information presented here has been derived from NRCS inventory data. Field observations from range trained personnel were also used. Those involved in developing this site include: Chris Krassin, Range Management Specialist, NRCS and Everet Bainter, Range Management Specialist, NRCS. Other sources used as references include USDA NRCS Water and Climate Center, USDA NRCS

Contributors

C. Krassin

Rangeland health reference sheet

Interpreting Indicators of Rangeland Health is a qualitative assessment protocol used to determine ecosystem condition based on benchmark characteristics described in the Reference Sheet. A suite of 17 (or more) indicators are typically considered in an assessment. The ecological site(s) representative of an assessment location must be known prior to applying the protocol and must be verified based on soils and climate. Current plant community cannot be used to identify the ecological site.

Author(s)/participant(s)	
Contact for lead author	
Date	07/01/2005
Approved by	
Approval date	
Composition (Indicators 10 and 12) based on	Annual Production

Indicators

1. Number and extent of rills: Rills should not be present

2. Presence of water flow patterns: Barely observable

3. Number and height of erosional pedestals or terracettes: Essentially non-existent

4. Bare ground from Ecological Site Description or other studies (rock, litter, lichen, moss, plant canopy are not bare ground):
Bare ground is 25-35% occurring in small areas throughout site

5. Number of gullies and erosion associated with gullies: Active gullies should not be present

6. Extent of wind scoured, blowouts and/or depositional areas: Active blowouts should not be present

7. Amount of litter movement (describe size and distance expected to travel): Little to no plant litter movement. Plant litter remains in place and is not moved by erosional forces.

8. Soil surface (top few mm) resistance to erosion (stability values are averages - most sites will show a range of values): Plant cover and litter is at 60% or greater of soil surface and maintains soil surface integrity. Soil Stability class is anticipated to be 4 or greater.

9. Soil surface structure and SOM content (include type of structure and A-horizon color and thickness): Use Soil Series description for depth and color of A-horizon

10. Effect of community phase composition (relative proportion of different functional groups) and spatial distribution on infiltration and runoff: Grass canopy and basal cover should reduce raindrop impact and slow overland flow providing increased time for infiltration to occur. Healthy deep rooted native grasses enhance infiltration and reduce runoff. Infiltration is Moderately rapid to Very Rapid.

11. Presence and thickness of compaction layer (usually none; describe soil profile features which may be mistaken for compaction on this site): No compaction layer or soil surface crusting should be present.

12. Functional/Structural Groups (list in order of descending dominance by above-ground annual-production or live foliar cover using symbols: >>, >, = to indicate much greater than, greater than, and equal to):

Dominant:

Sub-dominant:

Other:

Additional: Tall and Mid stature Grasses > Shrubs > Short stature grasses/grasslikes > Forbs

13. Amount of plant mortality and decadence (include which functional groups are expected to show mortality or decadence): Very Low

14. Average percent litter cover (%) and depth (in): Average litter cover is 20-30% with depths of 0.1 to 0.25 inches

15. Expected annual annual-production (this is TOTAL above-ground annual-production, not just forage annual-production): 500 lbs/ac

16. Potential invasive (including noxious) species (native and non-native). List species which BOTH characterize degraded states and have the potential to become a dominant or co-dominant species on the ecological site if their future establishment and growth is not actively controlled by management interventions. Species that become dominant for only one to several years (e.g., short-term response to drought or wildfire) are not invasive plants. Note that unlike other indicators, we are describing what is NOT expected in the reference state for the ecological site: Threadleaf sedge, Threeawns, Yucca, Skunkbush sumac, Big sagebrush, Annuals, Exotics, Unpalatable forbs, and Species found on Noxious Weed List
-

17. Perennial plant reproductive capability: All species are capable of reproducing
-