

Ecological site R032XY250WY
Sandy (Sy)
5-9" Wind River Basin Precipitation Zone

Accessed: 08/20/2026

General information

Provisional. A provisional ecological site description has undergone quality control and quality assurance review. It contains a working state and transition model and enough information to identify the ecological site.

Associated sites

R032XY204WY	Clayey (Cy) 5-9" Wind River Basin Precipitation Zone
R032XY212WY	Gravelly (Gr) 5-9" Wind River Basin Precipitation Zone
R032XY222WY	Loamy (Ly) 5-9" Wind River Basin Precipitation Zone
R032XY228WY	Lowland (LL) 5-9" Wind River Basin Precipitation Zone
R032XY266WY	Shallow Sandy (SwSy) 5-9" Wind River Basin Precipitation Zone

Similar sites

R032XY350WY	Sandy (Sy) 10-14" East Precipitation Zone Sandy 10-14" East P.Z. has higher production.
-------------	---

Table 1. Dominant plant species

Tree	Not specified
Shrub	Not specified
Herbaceous	Not specified

Physiographic features

This site occurs on nearly level to 35% slopes.

Table 2. Representative physiographic features

Landforms	(1) Alluvial fan (2) Hill (3) Plateau
Flooding duration	Brief (2 to 7 days)
Flooding frequency	None to rare
Ponding duration	Very brief (4 to 48 hours)
Ponding frequency	None to rare
Elevation	1,370 – 2,010 m
Slope	0 – 40 %
Ponding depth	0 cm
Aspect	Aspect is not a significant factor

Climatic features

Annual precipitation ranges from 5-9 inches per year. The normal precipitation pattern shows peaks in May and June and a secondary peak in September. This amounts to about 50% of the mean annual precipitation. Much of the moisture that falls in the latter part of the summer is lost by evaporation and much of the moisture that falls during the winter is lost by sublimation. Average snowfall is about 20 inches annually. Wide fluctuations may occur in yearly precipitation and result in more dry years than those with more than normal precipitation.

Temperatures show a wide range between summer and winter and between daily maximums and minimums, due to the high elevation and dry air, which permits rapid incoming and outgoing radiation. Cold air outbreaks from Canada in winter move rapidly from northwest to southeast and account for extreme minimum temperatures. Chinook winds may occur in winter and bring rapid rises in temperature. Extreme storms may occur during the winter, but most severely affect ranch operations during late winter and spring.

High winds are generally blocked from the basin by high mountains, but can occur in conjunction with an occasional thunderstorm.

Growth of native cool-season plants begins about April 1 and continues to about July 1. Cool weather and moisture in September may produce some green up of cool season plants that will continue to late October.

The following information is from the "Pavillion" climate station:

Minimum Maximum 5 yrs. out of 10 between

Frost-free period (days): 95 175 May 19 – September 19

Freeze-free period (days): 98 185 May 6 – October 3

Mean Annual Precipitation (inches): 2.50 12.54

Mean annual precipitation: 7.85 inches

Mean annual air temperature: 44.53 F (30.5 F Avg. Min. to 58.5 F Avg. Max.)

For detailed information visit the Natural Resources Conservation Service National Water and Climate Center at <http://www.wcc.nrcs.usda.gov/> website. Other climate station(s) representative of this precipitation zone include "Riverton", "Arminto", and

“Lost Cabin”.

Table 3 Representative climatic features

Frost-free period (average)	180 days
Freeze-free period (average)	190 days
Precipitation total (average)	230 mm

Influencing water features

Stream Type: None

Soil features

The soils of this site are moderately deep (greater than 20" to bedrock) to very deep, well-drained soils that formed in alluvium or alluvium over residuum. These soils have moderately slow, moderate, moderately rapid, or rapid permeability. The surface soil will vary from 3 to 6 inches deep. Coarser topsoils may be included if underlain by finer textured subsoil. The soil characteristics having the most influence on plant community are light texture, which can affect the available moisture and amount of calcium carbonates in the profile.

Major Soil Series correlated to this site include: Apron, Griffy, Wallson, Worland, Saddle, Trook, Kinnear, Pavillion, Ethete

Table 4. Representative soil features

Surface texture	(1) Fine sandy loam (2) Sandy loam (3) Loam
Family particle size	(1) Sandy
Drainage class	Well drained to excessively drained
Permeability class	Moderately slow to rapid
Soil depth	50 – 150 cm
Surface fragment cover <=3"	Not specified
Surface fragment cover >3"	Not specified
Available water capacity (0-101.6cm)	3.05 – 20 cm

Calcium carbonate equivalent (0-101.6cm)	0 – 40 %
Electrical conductivity (0-101.6cm)	0 – 10 mmhos/cm
Sodium adsorption ratio (0-101.6cm)	Not specified
Soil reaction (1:1 water) (0-101.6cm)	7.4 – 8.4
Subsurface fragment volume <=3" (Depth not specified)	Not specified
Subsurface fragment volume >3" (Depth not specified)	Not specified

Ecological dynamics

Potential vegetation on this site is dominated by mid cool-season perennial grasses. Other significant vegetation includes winterfat, big sagebrush, and a variety of forbs. The expected potential composition for this site is about 70% grasses, 15% forbs and 15% woody plants. The composition and production will vary naturally due to historical use, fluctuating precipitation and fire frequency.

As this site deteriorates, species such as threadleaf sedge, blue grama, and big sagebrush will increase. Weedy annuals will invade. Cool season grasses such as needleandthread, Indian ricegrass, and rhizomatous wheatgrasses will decrease in frequency and production.

Big sagebrush may become dominant on areas with an absence of fire and sufficient amount of precipitation. Wildfires are actively controlled in recent times and as a result old decadent stands of big sagebrush persist. Chemical control using herbicides has replaced the historic role of fire on this site. Recently, prescribed burning has regained some popularity.

Due to the amount and pattern of the precipitation, the big sagebrush component may not be resilient once it has been removed or severely reduced if a vigorous stand of grass exists and is maintained. On these areas, threadleaf sedge and blue grama may become dominant if the area is subjected to a combination of frequent and severe grazing especially yearlong grazing. As a result, a dense sod cover of threadleaf sedge and blue grama will become established.

The Historic Climax Plant Community (description follows the plant community diagram) has been determined by study of rangeland relic areas, or areas protected from excessive disturbance. Trends in plant communities going from heavily grazed areas to lightly grazed areas, seasonal use pastures, and historical accounts have also been used.

The following is a State and Transition Model Diagram that illustrates the common plant communities (states) that can occur on the site and the transitions between these communities. The ecological processes will be discussed in more detail in the plant community narratives following the diagram.

Plant Community Narratives

Following are the narratives for each of the described plant communities. These plant communities may not represent every possibility, but they probably are the most prevalent and repeatable plant communities. The plant composition tables shown above have been developed from the best available knowledge at the time of this revision. As more data is collected, some of these plant communities may

be revised or removed, and new ones may be added. None of these plant communities should necessarily be thought of as “Desired Plant Communities”. According to the USDA NRCS National Range and Pasture Handbook, Desired Plant Communities (DPC's) will be determined by the decision-makers and will meet minimum quality criteria established by the NRCS. The main purpose for including any description of a plant community here is to capture the current knowledge and experience at the time of this revision.

State and transition model

Additional community tables

Table 5. Community 1.1 plant community composition

Group	Common Name	Symbol	Scientific Name	Annual Production ()	Foliar Cover (%)
Grass/Grasslike					
1				135-224	
	needle and thread	HECO26	<i>Hesperostipa comata</i>	135-224	–
2				67-112	
	Indian ricegrass	ACHY	<i>Achnatherum hymenoides</i>	67-112	–
3				22-45	
	western wheatgrass	PASM	<i>Pascopyrum smithii</i>	22-45	–
4				45-90	
	Grass, perennial	2GP	<i>Grass, perennial</i>	0-22	–
	threeawn	ARIST	<i>Aristida</i>	0-22	–
	blue grama	BOGR2	<i>Bouteloua gracilis</i>	0-22	–
	threadleaf sedge	CAFI	<i>Carex filifolia</i>	0-22	–
	prairie sandreed	CALO	<i>Calamovilfa longifolia</i>	0-22	–
	Montana wheatgrass	ELAL7	<i>Elymus albicans</i>	0-22	–
	squirreltail	ELELE	<i>Elymus elymoides</i> ssp. <i>elymoides</i>	0-22	–
	prairie Junegrass	KOMA	<i>Koeleria macrantha</i>	0-22	–
	Sandberg bluegrass	POSE	<i>Poa secunda</i>	0-22	–
	bluebunch wheatgrass	PSSP6	<i>Pseudoroegneria spicata</i>	0-22	–
	sand dropseed	SPCR	<i>Sporobolus cryptandrus</i>	0-22	–
Forb					
5				45-90	
	Forb, perennial	2FP	<i>Forb, perennial</i>	0-22	–
	yarrow	ACHIL	<i>Achillea</i>	0-22	–
	textile onion	ALTE	<i>Allium textile</i>	0-22	–
	Franklin's sandwort	ARFR	<i>Arenaria franklinii</i>	0-22	–
	prairie sagewort	ARFR4	<i>Artemisia frigida</i>	0-22	–
	Missouri milkvetch	ASMI10	<i>Astragalus missouriensis</i>	0-22	–
	wavyleaf Indian paintbrush	CAAPM	<i>Castilleja applegatei</i> ssp. <i>martinii</i>	0-22	–
	bastard toadflax	COUM	<i>Comandra umbellata</i>	0-22	–
	little larkspur	DEBI	<i>Delphinium bicolor</i>	0-22	–
	cutleaf daisy	ERCO4	<i>Erigeron compositus</i>	0-22	–
	threadleaf fleabane	ERFI2	<i>Erigeron filifolius</i>	0-22	–
	parsnipflower buckwheat	ERHE2	<i>Eriogonum heracleoides</i>	0-22	–
	fleabane	ERIGE2	<i>Erigeron</i>	0-22	–
	scarlet beeblossom	GACO5	<i>Gaura coccinea</i>	0-22	–
	leafy wildparsley	MUDI	<i>Musineon divaricatum</i>	0-22	–

	plains pricklypear	OPPO	<i>Opuntia polyacantha</i>	0-22	–
	white locoweed	OXSES2	<i>Oxytropis sericea</i> var. <i>speciosa</i>	0-22	–
	spiny phlox	PHHO	<i>Phlox hoodii</i>	0-22	–
	lemon scurfpea	PSLA3	<i>Psoraleidium lanceolatum</i>	0-22	–
	scarlet globemallow	SPCO	<i>Sphaeralcea coccinea</i>	0-22	–
	smooth woodyaster	XYGL	<i>Xylorhiza glabriuscula</i>	0-22	–
	meadow deathcamas	ZIVE	<i>Zigadenus venenosus</i>	0-22	–
Shrub/Vine					
6				22-45	
	Wyoming big sagebrush	ARTRW8	<i>Artemisia tridentata</i> ssp. <i>wyomingensis</i>	22-45	–
7				0-22	
	winterfat	KRASC	<i>Krascheninnikovia</i>	0-22	–
8				22-45	
	Shrub (>.5m)	2SHRUB	<i>Shrub (>.5m)</i>	0-22	–
	silver sagebrush	ARCA13	<i>Artemisia cana</i>	0-22	–
	fourwing saltbush	ATCA2	<i>Atriplex canescens</i>	0-22	–
	yellow rabbitbrush	CHVI8	<i>Chrysothamnus viscidiflorus</i>	0-22	–
	rubber rabbitbrush	ERNA10	<i>Ericameria nauseosa</i>	0-22	–
	skunkbush sumac	RHTR	<i>Rhus trilobata</i>	0-22	–

Table 6. Community 2.1 plant community composition

Group	Common Name	Symbol	Scientific Name	Annual Production ()	Foliar Cover (%)
-------	-------------	--------	-----------------	----------------------	------------------

Table 7. Community 3.1 plant community composition

Group	Common Name	Symbol	Scientific Name	Annual Production ()	Foliar Cover (%)
-------	-------------	--------	-----------------	----------------------	------------------

Table 8. Community 4.1 plant community composition

Group	Common Name	Symbol	Scientific Name	Annual Production ()	Foliar Cover (%)
-------	-------------	--------	-----------------	----------------------	------------------

Animal community

Animal Community – Wildlife Interpretations Historic Climax Plant Community: The predominance of grasses in this plant community favors grazers and mixed-feeders, such as bison, elk, and antelope. Suitable thermal and escape cover for deer may be limited due to the low quantities of woody plants. However, topographical variations could provide some escape cover. When found adjacent to sagebrush dominated states, this plant community may provide brood rearing/foraging areas for sage grouse, as well as lek sites. Other birds that would frequent this plant community include western meadowlarks, horned larks, and golden eagles. Many grassland obligate small mammals would occur here. Perennial Grass/Big Sagebrush Plant Community: The combination of an overstory of sagebrush and an understory of grasses and forbs provide a very diverse plant community for wildlife. The crowns of sagebrush tend to break up hard crusted snow on winter ranges, so mule deer and antelope may use this state for foraging and cover year-round, as would cottontail and jack rabbits. It provides important winter, nesting, brood-rearing, and foraging habitat for sage grouse. Brewer's sparrows' nest in big sagebrush plants, and hosts of other nesting birds utilize stands in the 20-30% cover range. Mixed Shrub/Bare Ground Plant Community: This plant community can provide important winter foraging for elk, mule deer and antelope, as brush can approach 15% protein and 40-60% digestibility during that time. This community provides nesting habitat for upland game birds. However, it provides little foraging opportunities for upland game birds, as fewer forbs are available. Many grassland obligate small mammals would occur here. Threadleaf Sedge and/or Blue Grama Sod Plant Community: These communities provide limited foraging for antelope and other grazers. They may be used as a foraging site by sage grouse if proximal to woody cover and if the Historic Climax Plant Community or the Perennial Grass/Big Sagebrush Plant Community is limited. Generally, these are not target plant communities for wildlife habitat management.

Animal Community – Grazing Interpretations The following table lists suggested stocking rates for cattle under continuous season-long grazing under normal growing conditions. These are conservative estimates that should be used only as guidelines in the initial stages of the conservation planning process. Often, the current plant composition does not entirely match any particular plant community (as described in this ecological site description). Because of this, a field visit is recommended, in all cases, to document plant composition and production. More precise carrying capacity estimates should eventually be calculated using this information along with animal preference data, particularly when grazers other than cattle are involved. Under more intensive grazing management, improved harvest efficiencies can result in an increased carrying capacity. If distribution problems occur, stocking rates must be reduced to maintain plant

health and vigor. Plant Community Production Carrying Capacity* (lb./ac) (AUM/ac) Historical Climax Plant Community 225-600 .20 Perennial Grass/Big Sagebrush 180-480 .16 Mixed Shrub/Bare Ground 150-400 .10 Threadleaf Sedge and/or Blue Grama Sod 55-150 .05 * - Continuous, season-long grazing by cattle under average growing conditions. Grazing by domestic livestock is one of the major income-producing industries in the area. Rangeland in this area may provide yearlong forage for cattle, sheep, or horses. During the dormant period, the forage for livestock use needs to be supplemented with protein because the quality does not meet minimum livestock requirements.

Hydrological functions

Water is the principal factor limiting forage production on this site. This site is dominated by soils in hydrologic group B, with localized areas in hydrologic group C. Infiltration potential for this site varies from moderately rapid to rapid depending on soil hydrologic group and ground cover. Runoff varies from low to moderate. In many cases, areas with greater than 75% ground cover have the greatest potential for high infiltration and lower runoff. An example of an exception would be where short-grasses form a strong sod and dominate the site. Areas where ground cover is less than 50% have the greatest potential to have reduced infiltration and higher runoff (refer to Part 630, NRCS National Engineering Handbook for detailed hydrology information). Rills and gullies should not typically be present. Water flow patterns should be barely distinguishable if at all present. Pedestals are only slightly present in association with bunchgrasses. Litter typically falls in place, and signs of movement are not common. Chemical and physical crusts are rare to non-existent. Cryptogamic crusts are present, but only cover 1-2% of the soil surface.

Recreational uses

This site provides hunting opportunities for upland game species. The wide variety of plants which bloom from spring until fall have an esthetic value that appeals to visitors.

Wood products

No appreciable wood products are present on the site.

Other products

None noted.

Inventory data references

Information presented here has been derived from NRCS clipping data and other inventory data. Field observations from range trained personnel were also used. Those involved in developing this site include: Chris Krassin, Range Management Specialist, NRCS and Everett Bainter, Range Management Specialist, NRCS. Other sources used as references include: USDA NRCS Water and Climate Center, USDA NRCS National Range and Pasture Handbook, and USDA NRCS Soil Surveys from various counties.

Contributors

C. Krassin

Rangeland health reference sheet

Interpreting Indicators of Rangeland Health is a qualitative assessment protocol used to determine ecosystem condition based on benchmark characteristics described in the Reference Sheet. A suite of 17 (or more) indicators are typically considered in an assessment. The ecological site(s) representative of an assessment location must be known prior to applying the protocol and must be verified based on soils and climate. Current plant community cannot be used to identify the ecological site.

Author(s)/participant(s)	
Contact for lead author	
Date	07/01/2005

Approved by	
Approval date	
Composition (Indicators 10 and 12) based on	Annual Production

Indicators

1. Number and extent of rills: Rills should not be present

2. Presence of water flow patterns: Barely observable

3. Number and height of erosional pedestals or terracettes: Essentially non-existent

4. Bare ground from Ecological Site Description or other studies (rock, litter, lichen, moss, plant canopy are not bare ground):
Bare ground is 25-35% occurring in small areas throughout site

5. Number of gullies and erosion associated with gullies: Active gullies should not be present

6. Extent of wind scoured, blowouts and/or depositional areas: None

7. Amount of litter movement (describe size and distance expected to travel): Little to no plant litter movement. Plant litter remains in place and is not moved by erosional forces.

8. Soil surface (top few mm) resistance to erosion (stability values are averages - most sites will show a range of values):
Plant cover and litter is at 60% or greater of soil surface and maintains soil surface integrity. Soil Stability class is anticipated to be 4 or greater.

9. Soil surface structure and SOM content (include type of structure and A-horizon color and thickness): Use Soil Series description for depth and color of A-horizon

10. Effect of community phase composition (relative proportion of different functional groups) and spatial distribution on infiltration and runoff: Grass canopy and basal cover should reduce raindrop impact and slow overland flow providing increased time for infiltration to occur. Healthy deep rooted native grasses enhance infiltration and reduce runoff. Infiltration is Moderately Rapid to

Rapid.

11. Presence and thickness of compaction layer (usually none; describe soil profile features which may be mistaken for compaction on this site): No compaction layer or soil surface crusting should be present.
-

12. Functional/Structural Groups (list in order of descending dominance by above-ground annual-production or live foliar cover using symbols: >>, >, = to indicate much greater than, greater than, and equal to):

Dominant:

Sub-dominant:

Other:

Additional: Mid stature Bunch Grasses > Shrubs > Forbs > Mid Stature Rhizomatous Grasses > Short stature Grasses/Grasslike

13. Amount of plant mortality and decadence (include which functional groups are expected to show mortality or decadence):
Mid stature Bunch Grasses > Shrubs > Forbs > Mid Stature Rhizomatous Grasses > Short stature Grasses/Grasslike
-

14. Average percent litter cover (%) and depth (in): Average litter cover is 20-30% with depths of 0.1 to 0.25 inches
-

15. Expected annual annual-production (this is TOTAL above-ground annual-production, not just forage annual-production):
400 lbs/ac
-

16. Potential invasive (including noxious) species (native and non-native). List species which BOTH characterize degraded states and have the potential to become a dominant or co-dominant species on the ecological site if their future establishment and growth is not actively controlled by management interventions. Species that become dominant for only one to several years (e.g., short-term response to drought or wildfire) are not invasive plants. Note that unlike other indicators, we are describing what is NOT expected in the reference state for the ecological site: Threadleaf sedge, Blue grama, Big sagebrush, Yucca, Prickly Pear, Cheatgrass and other annuals, Exotics, and Species found on Noxious Weed List
-

17. Perennial plant reproductive capability: All species are capable of reproducing
-