

Ecological site R032XY278WY
Wetland (WL)
5-9" Wind River Basin Precipitation Zone

Accessed: 08/21/2026

General information

Provisional. A provisional ecological site description has undergone quality control and quality assurance review. It contains a working state and transition model and enough information to identify the ecological site.

Associated sites

R032XY228WY	Lowland (LL) 5-9" Wind River Basin Precipitation Zone
R032XY238WY	Saline Lowland (SL) 5-9" Wind River Basin Precipitation Zone
R032XY242WY	Saline Subirrigated (SS) 5-9" Wind River Basin Precipitation Zone

Similar sites

R032XY378WY	Wetland (WL) 10-14" East Precipitation Zone Wetland 10-14" East P.Z. has higher production.
-------------	--

Table 1. Dominant plant species

Tree	Not specified
Shrub	Not specified
Herbaceous	Not specified

Physiographic features

This site normally occurs on level to nearly level bottomlands near springs, seeps and sloughs.

Table 2. Representative physiographic features

Landforms	(1) Drainageway (2) Oxbow (3) Stream terrace
-----------	--

Flooding duration	Very brief (4 to 48 hours) to long (7 to 30 days)
Flooding frequency	Occasional to frequent
Ponding duration	Brief (2 to 7 days) to very long (more than 30 days)
Ponding frequency	Frequent
Elevation	1,370 – 2,010 m
Slope	0 – 10 %
Ponding depth	0 – 30 cm
Water table depth	0 – 50 cm
Aspect	Aspect is not a significant factor

Climatic features

Annual precipitation ranges from 5-9 inches per year. The normal precipitation pattern shows peaks in May and June and a secondary peak in September. This amounts to about 50% of the mean annual precipitation. Much of the moisture that falls in the latter part of the summer is lost by evaporation and much of the moisture that falls during the winter is lost by sublimation. Average snowfall is about 20 inches annually. Wide fluctuations may occur in yearly precipitation and result in more dry years than those with more than normal precipitation.

Temperatures show a wide range between summer and winter and between daily maximums and minimums, due to the high elevation and dry air, which permits rapid incoming and outgoing radiation. Cold air outbreaks from Canada in winter move rapidly from northwest to southeast and account for extreme minimum temperatures. Chinook winds may occur in winter and bring rapid rises in temperature. Extreme storms may occur during the winter, but most severely affect ranch operations during late winter and spring.

High winds are generally blocked from the basin by high mountains, but can occur in conjunction with an occasional thunderstorm.

Growth of native cool-season plants begins about April 1 and continues to about July 1. Cool weather and moisture in September may produce some green up of cool season plants that will continue to late October.

The following information is from the "Pavillion" climate station:

Minimum Maximum 5 yrs. out of 10 between

Frost-free period (days): 95 175 May 19 – September 19

Freeze-free period (days): 98 185 May 6 – October 3

Mean Annual Precipitation (inches): 2.50 12.54

Mean annual precipitation: 7.85 inches

Mean annual air temperature: 44.53 F (30.5 F Avg. Min. to 58.5 F Avg. Max.)

For detailed information visit the Natural Resources Conservation Service National Water and Climate Center at <http://www.wcc.nrcs.usda.gov/> website. Other climate station(s) representative of this precipitation zone include "Riverton", "Arminto", and "Lost Cabin".

Table 3 Representative climatic features

Frost-free period (average)	180 days
Freeze-free period (average)	190 days
Precipitation total (average)	230 mm

Influencing water features

Stream Type: C (Rosgen)

Soil features

This site consists of deep to very deep poorly drained soils formed in alluvium with a water table above the surface for part but not all of the growing season. They are on nearly level to slightly depressed areas with poor surface drainage. In some places the surface layers have a high organic matter content. The soil characteristics having the most influence on the plant community are depth to a water table at or near the surface for all of the growing season.

Major Soil Series correlated to this site include: Fluvaquents

Table 4. Representative soil features

Surface texture	(1) Clay (2) Clay loam (3) Loam
Family particle size	(1) Loamy
Drainage class	Poorly drained to excessively drained
Permeability class	Slow to moderate
Soil depth	50 – 150 cm
Surface fragment cover <=3"	Not specified
Surface fragment cover >3"	Not specified
Available water capacity (0-101.6cm)	5.59 – 16.76 cm
Calcium carbonate equivalent (0-101.6cm)	10 – 20 %

Electrical conductivity (0-101.6cm)	0 – 20 mmhos/cm
Sodium adsorption ratio (0-101.6cm)	0 – 20
Soil reaction (1:1 water) (0-101.6cm)	6.6 – 8.4
Subsurface fragment volume <=3" (Depth not specified)	Not specified
Subsurface fragment volume >3" (Depth not specified)	Not specified

Ecological dynamics

Potential vegetation on this site is dominated by plants that can tolerate soils which have a water table near the surface for most of the growing season and may be slightly saline. The expected potential composition for this site is about 80% grasses, 10% forbs and 10% woody plants. The composition and production will vary naturally due to historical use, fluctuating precipitation and fire frequency.

As this site deteriorates, species such as willows, rushes, and bulrush increase. Grasses such as Nebraska sedge and tufted hairgrass will decrease in frequency and production.

The Historic Climax Plant Community (description follows the plant community diagram) has been determined by study of rangeland relic areas, or areas protected from excessive disturbance. Trends in plant communities going from heavily grazed areas to lightly grazed areas, seasonal use pastures, and historical accounts have also been used.

The following is a State and Transition Model Diagram that illustrates the common plant communities (states) that can occur on the site and the transitions between these communities. The ecological processes will be discussed in more detail in the plant community narratives following the diagram.

Plant Community Narratives

Following are the narratives for each of the described plant communities. These plant communities may not represent every possibility, but they probably are the most prevalent and repeatable plant communities. The plant composition tables shown above have been developed from the best available knowledge at the time of this revision. As more data is collected, some of these plant communities may be revised or removed, and new ones may be added. None of these plant communities should necessarily be thought of as "Desired Plant Communities". According to the USDA NRCS National Range and Pasture Handbook, Desired Plant Communities (DPC's) will be determined by the decision-makers and will meet minimum quality criteria established by the NRCS. The main purpose for including any description of a plant community here is to capture the current knowledge and experience at the time of this revision.

State and transition model

Additional community tables

Table 5. Community 1.1 plant community composition

Group	Common Name	Symbol	Scientific Name	Annual Production ()	Foliar Cover (%)
Grass/Grasslike					

1				1261-1765	
	Nebraska sedge	CANE2	<i>Carex nebrascensis</i>	1261-1765	–
2				504-1261	
	water sedge	CAAQ	<i>Carex aquatilis</i>	504-1261	–
3				504-1261	
	beaked sedge	CARO6	<i>Carex rostrata</i>	504-1261	–
4				252-757	
5				252-757	
	Grass, perennial	2GP	<i>Grass, perennial</i>	0-252	–
	American sloughgrass	BESY	<i>Beckmannia syzigachne</i>	0-252	–
	reedgrass	CALAM	<i>Calamagrostis</i>	0-252	–
	northern reedgrass	CASTI3	<i>Calamagrostis stricta</i> ssp. <i>inexpansa</i>	0-252	–
	flatsedge	CYPER	<i>Cyperus</i>	0-252	–
	fowl mannagrass	GLST	<i>Glyceria striata</i>	0-252	–
	rush	JUNCU	<i>Juncus</i>	0-252	–
	reed canarygrass	PHAR3	<i>Phalaris arundinacea</i>	0-252	–
	bluegrass	POA	<i>Poa</i>	0-252	–
	bulrush	SCIRP	<i>Scirpus</i>	0-252	–
	alkali cordgrass	SPGR	<i>Spartina gracilis</i>	0-252	–
Forb					
6				252-504	
	Forb, perennial	2FP	<i>Forb, perennial</i>	0-252	–
	silverweed cinquefoil	ARAN7	<i>Argentina anserina</i>	0-252	–
	smooth horsetail	EQLA	<i>Equisetum laevigatum</i>	0-252	–
	Rocky Mountain iris	IRMI	<i>Iris missouriensis</i>	0-252	–
	rough bugleweed	LYAS	<i>Lycopus asper</i>	0-252	–
	waterhorehound	LYCOP4	<i>Lycopus</i>	0-252	–
	wild mint	MEAR4	<i>Mentha arvensis</i>	0-252	–
	common plantain	PLMA2	<i>Plantago major</i>	0-252	–
	alkali buttercup	RACY	<i>Ranunculus cymbalaria</i>	0-252	–
	sagebrush buttercup	RAGL	<i>Ranunculus glaberrimus</i>	0-252	–
	blue-eyed grass	SISYR	<i>Sisyrinchium</i>	0-252	–
	marsh hedgenettle	STPA	<i>Stachys palustris</i>	0-252	–
Shrub/Vine					
7				252-757	
	Shrub (>.5m)	2SHRUB	<i>Shrub (>.5m)</i>	0-252	–
	Woods' rose	ROWO	<i>Rosa woodsii</i>	0-252	–
	narrowleaf willow	SAEX	<i>Salix exigua</i>	0-252	–
	willow	SALIX	<i>Salix</i>	0-252	–

Table 6. Community 2.1 plant community composition

Group	Common Name	Symbol	Scientific Name	Annual Production ()	Foliar Cover (%)
-------	-------------	--------	-----------------	----------------------	------------------

Table 7. Community 3.1 plant community composition

Group	Common Name	Symbol	Scientific Name	Annual Production ()	Foliar Cover (%)
-------	-------------	--------	-----------------	----------------------	------------------

Table 8. Community 4.1 plant community composition

Group	Common Name	Symbol	Scientific Name	Annual Production ()	Foliar Cover (%)
-------	-------------	--------	-----------------	----------------------	------------------

Table 9. Community 5.1 plant community composition

Group	Common Name	Symbol	Scientific Name	Annual Production ()	Foliar Cover (%)
-------	-------------	--------	-----------------	----------------------	------------------

Table 10. Community 6.1 plant community composition

Group	Common Name	Symbol	Scientific Name	Annual Production ()	Foliar Cover (%)
-------	-------------	--------	-----------------	----------------------	------------------

Table 11. Community 7.1 plant community composition

Group	Common Name	Symbol	Scientific Name	Annual Production ()	Foliar Cover (%)
-------	-------------	--------	-----------------	----------------------	------------------

Animal community

Animal Community – Wildlife Interpretations Nebraska sedge/Tufted Hairgrass Plant Community: This plant community is very important for most wildlife in the area. Over 80% of all wildlife use this site to fulfill some part of their habitat needs. It provides forage and thermal and hiding cover for mule deer and moose. It provides nesting habitat for shorebirds, songbirds, and waterfowl as well as ground nesting birds such as harriers. The lush herbaceous material produces insects for sage grouse brood rearing and foraging. Dense ground cover provides escape cover, forage, and breeding areas for small mammals which draw predators such as raptors, red fox and coyote. Other birds that would frequent this plant community include red-wing blackbirds, sandhill cranes, western meadowlarks, neotropical migrants, and golden eagles. Mid-Sedge/Willow Plant Community: The abundant production and proximity to water make this state important for livestock and wildlife such as birds, mule deer, and whitetail deer. This plant community is useful for the wildlife that would use the Historic Climax Plant Community. The increase in willow production makes this even more attractive to some wildlife due to the increase in thermal and escape cover. It can provide foraging and nesting opportunities for upland game birds and songbirds. Low-Sedge/Baltic Rush Plant Community: The abundant production and proximity to water make this state important for livestock and wildlife such as birds, mule deer and whitetail deer. This plant community may be useful for the same large grazers that would use the Historic Climax Plant Community. However, the plant community composition is less diverse, and thus, less apt to meet the seasonal needs of these animals. It may provide some foraging opportunities for upland game birds and songbirds, when it occurs proximal to woody cover. Cattail/Low Sedge Plant Community: The abundant production and proximity to water make this state important for livestock and wildlife such as birds, mule deer, and whitetail deer. This plant community is useful for the same large grazers that would use the Historic Climax Plant Community. The increase in cattail production makes this even more attractive to some wildlife due to the increase in thermal and escape cover. It can provide foraging and nesting opportunities for upland game birds and songbirds. Sedge/Cattail Plant Community: The abundant production and proximity to water make this state important for livestock and wildlife such as birds, mule deer, and whitetail deer. This plant community is useful for the same large grazers that would use the Historic Climax Plant Community. The increase in cattail production makes this even more attractive to some wildlife due to the increase in thermal and escape cover. It can provide foraging and nesting opportunities for upland game birds and songbirds. Sedge/Saltcedar and/or Russian Olive Plant Community: The abundant production and proximity to water make this state important for livestock and wildlife such as birds, mule deer, and whitetail deer. This plant community is useful for the same large grazers that would use the Historic Climax Plant Community. The increase in tall shrubs production makes this even more attractive to some wildlife due to the increase in thermal and escape cover. It can provide foraging and nesting opportunities for upland game birds and songbirds. Some species utilize the Russian olive berries for food and are attracted to these colonized areas. Dense Saltcedar and/or Russian Olive Stand Plant Community: The proximity to water makes this state important for wildlife such as birds, mule deer, and whitetail deer. This is useful for the same large grazers that would use the Historic Climax Plant Community. The low production of herbaceous understory of this plant community decreases the foraging potential and cover for many wildlife species. The increase in tall shrubs, however, makes this an attractive site for thermal and escape cover for large grazers and upland birds. It can provide foraging and nesting opportunities for songbirds. Some species utilize the Russian olive berries for food and are attracted to these colonized areas. Animal Community – Grazing Interpretations The following table lists suggested stocking rates for cattle under continuous season-long grazing under normal growing conditions. These are conservative estimates that should be used only as guidelines in the initial stages of the conservation planning process. Often, the current plant composition does not entirely match any particular plant community (as described in this ecological site description). Because of this, a field visit is recommended, in all cases, to document plant composition and production. More precise carrying capacity estimates should eventually be calculated using this information along with animal preference data, particularly when grazers other than cattle are involved. Under more intensive grazing management, improved harvest efficiencies can result in an increased carrying capacity. If distribution problems occur, stocking rates must be reduced to maintain plant health and vigor. Plant Community Production Carrying Capacity* (lb./ac) (AUM/ac)

Historic Climax Plant Community	3000-6000	2.5
Mid-Sedge/Willow	2750-5500	1.8
Low-Sedge/Baltic Rush	1200-2500	0.8
Cattail/Low Sedge	2750-4000	1.0
Sedge/Cattail	3000-5000	1.2
Sedge/Saltcedar and/or Russian Olive	3000-5000	1.2
Dense Saltcedar and/or Russian Olive Stand	1750-2500	0.5

* - Continuous, season-long grazing by cattle under average growing conditions. Grazing by domestic livestock is one of the major income-producing industries in the area. Rangeland in this area may provide yearlong forage for cattle, sheep, or horses. During the dormant period, the forage for livestock use needs to be supplemented with protein because the quality does not meet minimum livestock requirements.

Hydrological functions

Climate is the principal factor limiting forage production on this site. This site is dominated by soils in hydrologic group B and C, with localized areas in hydrologic group D. Infiltration and runoff potential for this site varies from moderate to high depending on soil hydrologic group and water table. Runoff will be high on this site since the soil may be saturated. (Refer to Part 630, NRCS National Engineering Handbook for detailed hydraulic information. Rills and gullies should not typically be present. Water flow patterns should be barely distinguishable if at all present. Litter typically falls in place, and signs of movement are not common. Chemical and physical crusts are rare to non-existent. Cryptogamic crusts are present, but only cover 1-2% of the soil surface.

Recreational uses

This site provides hunting opportunities for upland game species. The wide variety of plants which bloom from spring until fall have an esthetic value that appeals to visitors.

Wood products

No appreciable wood products are present on the site.

Other products

none noted

Inventory data references

Information presented here has been derived from NRCS inventory data. Field observations from range trained personnel were also used. Those involved in developing this site include: Chris Krassin, Range Management Specialist, NRCS and Everet Bainter, Range Management Specialist, NRCS. Other sources used as references include USDA NRCS Water and Climate Center, USDA NRCS National Range and Pasture Handbook, USDI and USDA Interpreting Indicators of Rangeland Health Version 3, and USDA NRCS Soil Surveys from various counties.

Contributors

C. Krassin

Rangeland health reference sheet

Interpreting Indicators of Rangeland Health is a qualitative assessment protocol used to determine ecosystem condition based on benchmark characteristics described in the Reference Sheet. A suite of 17 (or more) indicators are typically considered in an assessment. The ecological site(s) representative of an assessment location must be known prior to applying the protocol and must be verified based on soils and climate. Current plant community cannot be used to identify the ecological site.

Author(s)/participant(s)	
Contact for lead author	
Date	07/01/2005
Approved by	
Approval date	
Composition (Indicators 10 and 12) based on	Annual Production

Indicators

1. Number and extent of rills: Rills should not be present

2. Presence of water flow patterns: Barely observable

3. Number and height of erosional pedestals or terracettes: Essentially non-existent

4. Bare ground from Ecological Site Description or other studies (rock, litter, lichen, moss, plant canopy are not bare ground):
Bare ground is less than 5%

5. Number of gullies and erosion associated with gullies: Active gullies should not be present

6. Extent of wind scoured, blowouts and/or depositional areas: None

7. Amount of litter movement (describe size and distance expected to travel): Little to no plant litter movement. Plant litter remains in place and is not moved by erosional forces.

8. Soil surface (top few mm) resistance to erosion (stability values are averages - most sites will show a range of values):
Plant cover and litter is at 95% or greater of soil surface and maintains soil surface integrity. Soil Stability class is anticipated to be 5 or greater.

9. Soil surface structure and SOM content (include type of structure and A-horizon color and thickness): Use Soil Series description for depth and color of A-horizon

10. Effect of community phase composition (relative proportion of different functional groups) and spatial distribution on infiltration and runoff: Grass canopy and basal cover should reduce raindrop impact and slow overland flow providing increased time for infiltration to occur. Healthy deep rooted native grasses enhance infiltration and reduce runoff. Infiltration varies from moderate to low and runoff is high since the soil is usually saturated.

11. Presence and thickness of compaction layer (usually none; describe soil profile features which may be mistaken for compaction on this site): No compaction layer or soil surface crusting should be present.

12. **Functional/Structural Groups** (list in order of descending dominance by above-ground annual-production or live foliar cover using symbols: >>, >, = to indicate much greater than, greater than, and equal to):

Dominant:

Sub-dominant:

Other:

Additional: Tall Grasses and Grasslike > Mid stature Grasses/Grasslike = Shrubs/Trees > Forbs

13. **Amount of plant mortality and decadence** (include which functional groups are expected to show mortality or decadence):
Very Low

14. **Average percent litter cover (%) and depth (in):** Average litter cover is 50-55% with depths of 0.75 to 1.5 inches

15. **Expected annual annual-production** (this is TOTAL above-ground annual-production, not just forage annual-production):
4500 lbs/ac

16. **Potential invasive (including noxious) species (native and non-native).** List species which BOTH characterize degraded states and have the potential to become a dominant or co-dominant species on the ecological site if their future establishment and growth is not actively controlled by management interventions. Species that become dominant for only one to several years (e.g., short-term response to drought or wildfire) are not invasive plants. Note that unlike other indicators, we are describing what is NOT expected in the reference state for the ecological site: Baltic rush, Bulrush, Arrowgrass, Horsetails, Kentucky Bluegrass, Saltcedar, Russian olive, Cattails, Annuals, other Exotics, and other Species found on Noxious Weed List.

17. **Perennial plant reproductive capability:** All species are capable of reproducing
