

Ecological site R034AY142WY

Saline Subirrigated

Green River and Great Divide Basins (SS)

Last updated: 2/24/2025
Accessed: 08/17/2026

General information

Provisional. A provisional ecological site description has undergone quality control and quality assurance review. It contains a working state and transition model and enough information to identify the ecological site.

Associated sites

R034AY138WY	Saline Lowland Green River and Great Divide Basins (SL) Saline Lowland
R034AY174WY	Subirrigated Green River and Great Divide Basins (Sb) Subirrigated
R034AY178WY	Wetland Green River and Great Divide Basins (WL) Wetland
R034AY128WY	Lowland Green River and Great Divide Basins (LL) Lowland

Similar sites

R034AY138WY	Saline Lowland Green River and Great Divide Basins (SL) Saline Lowland (SL) 7-9GR has more greasewood.
R034AY242WY	Saline Subirrigated Foothills and Basins West (SS) Saline Subirrigated (SS) 10-14W has higher production.

Table 1. Dominant plant species

Tree	Not specified
Shrub	Not specified
Herbaceous	Not specified

Physiographic features

This site occurs on nearly level land along perennial or intermittent streams, near seeps, sloughs, or springs. It is also found on broad, low lake terraces, lake plains, flood plains, on alluvial bottoms, and poorly-drained bottom lands adjacent to stream channels. These areas receive additional run-in water from higher sites and from a fluctuating water table, well within the root zone. Slopes are most commonly less than 3%.

Table 2. Representative physiographic features

Landforms	(1) Alluvial fan (2) Drainageway (3) Stream terrace
Flooding duration	Very brief (4 to 48 hours) to brief (2 to 7 days)
Flooding frequency	Occasional to frequent
Ponding frequency	None
Elevation	1,830 – 2,200 m
Slope	0 – 10 %
Ponding depth	0 cm
Water table depth	30 – 100 cm
Aspect	Aspect is not a significant factor

Climatic features

Annual precipitation ranges from 7-9 inches per year. Wide fluctuations may occur in yearly precipitation and result in more dry years than those with more than normal precipitation. Temperatures show a wide range between summer and winter and between daily maximums and minimums. This is predominantly due to the high elevation and dry air, which permits rapid incoming and outgoing radiation. Cold air outbreaks in winter move rapidly from northwest to southeast and account for extreme minimum temperatures. Extreme storms may occur during the winter, but most severely affect ranch operations during late winter and spring.

Daytime winds are generally stronger than nighttime and occasional strong storms may bring brief periods of high winds with gusts to more than 50 mph.

Growth of native cool season plants begins about April 15 and continues to about July 15. Some green up of cool season plants may occur in September if moisture is available.

For detailed information visit the Natural Resources Conservation Service National Water and Climate Center at <http://www.wcc.nrcs.usda.gov/cgibin/state.pl?state=wy> website. Other climate stations representative of this precipitation zone include "Bitter Creek", "Farson ", "Rock Springs FAA AP", and "Wamsutter" in Sweetwater County; "Church Buttes Gas PLT", and Mountain View" in Uinta County; "Fontenelle", "La Barge", and "Sage 4 NNW" in Lincoln County; and "Big Piney" in Sublette County.

Table 3 Representative climatic features

Frost-free period (average)	120 days
Freeze-free period (average)	130 days
Precipitation total (average)	230 mm

Influencing water features

Stream type: C (Rosgen)

Soil features

These are deep, saline, usually sodic soils on stream terraces. Sand and gravel may be at depths of 40 to 60 inches. The water table fluctuates during the growing season and is generally between 20-40 inches. Moderate to strong salt and alkali concentration in conjunction with moderately deep water tables characterize these soils. Salt crusts are commonly found on ridges and mounds during dry periods. Mottling or gleying may occur within 20-40 inches of the surface.

Major Soil Series correlated to this site include: Gunbarrel, Saleratus and phases of the Clowers, Leckman, and Pescar series.

Table 4. Representative soil features

Surface texture	(1) Loam (2) Fine sandy loam (3) Sandy loam
Family particle size	(1) Loamy
Drainage class	Somewhat poorly drained to moderately well drained
Permeability class	Moderate
Soil depth	40 – 150 cm
Surface fragment cover <=3"	0 – 10 %
Surface fragment cover >3"	Not specified
Available water capacity (0-101.6cm)	8.89 – 12.7 cm
Calcium carbonate equivalent (0-101.6cm)	0 – 10 %

Electrical conductivity (0-101.6cm)	10 – 20 mmhos/cm
Sodium adsorption ratio (0-101.6cm)	10 – 20
Soil reaction (1:1 water) (0-101.6cm)	10
Subsurface fragment volume <=3" (Depth not specified)	0 – 10 %
Subsurface fragment volume >3" (Depth not specified)	0 – 10 %

Ecological dynamics

As this site deteriorates from improper grazing management, species such as inland saltgrass and greasewood increase. Grasses such as alkali sacaton, basin wildrye, and Nuttall's alkaligrass will decrease in frequency and production. Basin wildrye will disappear from this site when hayed or with extended improper grazing during spring and summer.

These plant communities narratives may not represent every possibility, but they probably are the most prevalent and repeatable plant communities. The plant composition tables shown above have been developed from the best available knowledge at the time of this revision. As more data is collected, some of these plant communities may be revised or removed, and new ones may be added. None of these plant communities should necessarily be thought of as "Desired Plant Communities". According to the USDA NRCS National Range and Pasture Handbook, Desired Plant Communities (DPC's) will be determined by the decision-makers and will meet minimum quality criteria established by the NRCS. The main purpose for including any description of a plant community here is to capture the current knowledge and experience at the time of this revision.

The Reference Plant Community (description follows the plant community diagram) has been determined by study of rangeland relic areas, or areas protected from excessive disturbance. Trends in plant communities going from heavily grazed areas to lightly grazed areas, seasonal use pastures, and historical accounts have also been used.

The following is a State and Transition Model Diagram that illustrates the common plant communities (states) that can occur on the site and the transitions between these communities. The ecological processes will be discussed in more detail in the plant community narratives following the diagram.

State and transition model

Additional community tables

Table 5. Community 1.1 plant community composition

Group	Common Name	Symbol	Scientific Name	Annual Production ()	Foliar Cover (%)
Grass/Grasslike					
1				981-1261	
	alkali sacaton	SPAI	<i>Sporobolus airoides</i>	981-1261	–
2				420-841	
	basin wildrye	LECI4	<i>Leymus cinereus</i>	420-841	–
3				280-560	

	Nuttall's alkaligrass	PUNU2	<i>Puccinellia nuttalliana</i>	280-560	–
4				280-560	
	Grass, perennial	2GP	<i>Grass, perennial</i>	0-140	–
	inland sedge	CAIN11	<i>Carex interior</i>	0-140	–
	Nebraska sedge	CANE2	<i>Carex nebrascensis</i>	0-140	–
	saltgrass	DISP	<i>Distichlis spicata</i>	0-140	–
	mountain rush	JUARL	<i>Juncus arcticus ssp. littoralis</i>	0-140	–
	scratchgrass	MUAS	<i>Muhlenbergia asperifolia</i>	0-140	–
	western wheatgrass	PASM	<i>Pascopyrum smithii</i>	0-140	–
	Sandberg bluegrass	POSE	<i>Poa secunda</i>	0-140	–
Forb					
5				140-420	
	Forb, perennial	2FP	<i>Forb, perennial</i>	0-140	–
	milkvetch	ASTRA	<i>Astragalus</i>	0-140	–
	scouringrush horsetail	EQHY	<i>Equisetum hyemale</i>	0-140	–
	arrowgrass	TRIGL	<i>Triglochin</i>	0-140	–
Shrub/Vine					
6				28-140	
	greasewood	SAVE4	<i>Sarcobatus vermiculatus</i>	28-140	–
7				28-140	
	rubber rabbitbrush	ERNA10	<i>Ericameria nauseosa</i>	28-140	–

Table 6. Community 2.1 plant community composition

Group	Common Name	Symbol	Scientific Name	Annual Production ()	Foliar Cover (%)
-------	-------------	--------	-----------------	----------------------	------------------

Table 7. Community 3.1 plant community composition

Group	Common Name	Symbol	Scientific Name	Annual Production ()	Foliar Cover (%)
-------	-------------	--------	-----------------	----------------------	------------------

Table 8. Community 4.1 plant community composition

Group	Common Name	Symbol	Scientific Name	Annual Production ()	Foliar Cover (%)
-------	-------------	--------	-----------------	----------------------	------------------

Animal community

Animal Community – Wildlife Interpretations Alkali Sacaton/Basin Wildrye Plant Community (HCPC): This plant community is very important for many of the wildlife species in the area. With the presence of water at or near the soil surface, over 80% of all wildlife will use this site to fulfill some part of their habitat needs. It provides forage for mule deer and antelope. It provides nesting habitat for shorebirds, songbirds, and waterfowl as well as ground nesting birds such as harriers. The lush herbaceous material produces insects for sage grouse brood rearing and foraging. Dense ground cover provides escape cover, forage, and breeding areas for small mammals which draw predators such as raptors, red fox and coyote. Other birds that would frequent this plant community include red-wing blackbirds, sandhill cranes, western meadowlarks, and neotropical migrants. Alkali Muhly/Alkali Bluegrass Plant Community: This plant community may be beneficial for the same wildlife that would use the Historic Climax Plant Community. However, the plant community composition is less diverse, and thus, less apt to meet the seasonal needs of these animals. It may provide some foraging opportunities for sage grouse when it occurs proximal to woody cover. Foxtail Barley/Quackgrass Plant Community: This plant community exhibits a low level of plant species diversity. In most cases, it is not a desirable plant community to select as a wildlife habitat management objective. Salt Cedar/Inland Saltgrass Plant Community: This plant community exhibits a low level of plant species diversity due to the accumulation of salts in the soil. It may provide some thermal and escape cover for deer and moose if no other woody community is nearby, but in most cases it is not a desirable plant community to select as a wildlife habitat management objective. Animal Community – Grazing Interpretations The following table lists suggested stocking rates for cattle under continuous season-long grazing under normal growing conditions. These are conservative estimates that should be used only as guidelines in the initial stages of the conservation planning process. Often, the current plant composition does not entirely match any particular plant community (as described in this ecological site description). Because of this, a field visit is recommended, in all cases, to document plant composition and production. More precise carrying capacity estimates should eventually be calculated using this information along with animal preference data, particularly when grazers other than cattle are involved. Under more intensive grazing management, improved harvest efficiencies can result in an increased carrying capacity. Plant Community Production (lb./ac) and Carrying Capacity* (AUM/ac) Alkali Sacaton/Basin

Wildrye (HCPC) 2000-2800 lb./ac and .8 AUM/ac Alkali Muhly/Alkali Bluegrass 1000-1800 lb./ac and .5 AUM/ac Foxtail Barley/Quackgrass 500-2300 lb./ac and .3 AUM/ac Salt Cedar/Inland Saltgrass 200-800 lb./ac and .2 AUM/ac * - Continuous, season-long grazing by cattle under average growing conditions. Grazing by domestic livestock is one of the major income-producing industries in the area. Rangeland in this area may provide yearlong forage for cattle, sheep, or horses. During the dormant period, the forage for livestock use needs to be supplemented with protein because the quality does not meet minimum livestock requirements.

Hydrological functions

Salinity/alkalinity is the principal factor limiting forage production on this site. This site is dominated by soils in hydrologic groups C and D. Infiltration ranges from moderately slow to moderately well. Runoff potential for this site varies from moderate to high depending on soil hydrologic group and ground cover. In many cases, areas with greater than 75% ground cover have the greatest potential for high infiltration and lower runoff. Areas where ground cover is less than 50% have the greatest potential to have reduced infiltration and higher runoff (refer to Part 630, NRCS National Engineering Handbook for detailed hydrology information). Rills and gullies should not typically be present. Water flow patterns should be barely distinguishable if at all present. Pedestals are only slightly present in association with bunchgrasses. Litter typically falls in place, and signs of movement are not common. Chemical and physical crusts are sometimes present.

Recreational uses

This site provides a variety of hunting opportunities as well providing popular camping areas for recreationists. This site has a wide variety of forbs which bloom throughout spring and summer, providing esthetic values that appeal to visitors.

Wood products

No appreciable wood products are present on the site.

Other products

None noted.

Inventory data references

Inventory Data References (narrative) Information presented here has been derived from NRCS clipping data and other inventory data. Field observations from range trained personnel were also used. Those involved in developing this site include: Bill Christensen, Range Management Specialist, NRCS; Karen Clause, Range Management Specialist, NRCS; and Everet Bainter, Range Management Specialist, NRCS. Other sources used as references include: USDA NRCS Water and Climate Center, USDA NRCS National Range and Pasture Handbook, and USDA NRCS Soil Surveys from various counties. Inventory Data References Data Source Number of Records Sample Period State County SCS-RANGE-417 50 1966-1985 WY Sweetwater & others

Contributors

Karen Clause

Approval

Kirt Walstad, 2/24/2025

Rangeland health reference sheet

Interpreting Indicators of Rangeland Health is a qualitative assessment protocol used to determine ecosystem condition based on benchmark characteristics described in the Reference Sheet. A suite of 17 (or more) indicators are typically considered in an assessment. The ecological site(s) representative of an assessment location must be known prior to applying the protocol and must be verified based on soils and climate. Current plant community cannot be used to identify the ecological site.

Author(s)/participant(s)	K. Clause, B. Brazee, J. Haverkamp, E. Bainter
Contact for lead author	karen.clause@wy.usda.gov or 307-367-2257

Date	03/16/2007
Approved by	
Approval date	
Composition (Indicators 10 and 12) based on	Annual Production

Indicators

1. Number and extent of rills: Rare to nonexistent.

2. Presence of water flow patterns: Barely observable.

3. Number and height of erosional pedestals or terracettes: Rare to nonexistent.

4. Bare ground from Ecological Site Description or other studies (rock, litter, lichen, moss, plant canopy are not bare ground):
Bare ground can range from 0-5%.

5. Number of gullies and erosion associated with gullies: Active gullies should not be present.

6. Extent of wind scoured, blowouts and/or depositional areas: Minimal to nonexistent.

7. Amount of litter movement (describe size and distance expected to travel): Herbaceous litter not expected to move.

8. Soil surface (top few mm) resistance to erosion (stability values are averages - most sites will show a range of values): Soil Stability Index ratings range from 1 (interspaces) to 6 (under plant canopy), but average values should be 4.0 or greater.

9. Soil surface structure and SOM content (include type of structure and A-horizon color and thickness): Typically an A-horizon of greater than 2 inches (5 cm) with massive structure and color hues of 10YR or 2.5Y, values of 5-6, and chromas of 2-3. Organic matter is typically less than 1%.

10. Effect of community phase composition (relative proportion of different functional groups) and spatial distribution on infiltration and runoff: Plant community consists of 75-85% grasses, 15% forbs, and 0-10% shrubs. Dense plant canopy (70-100%) and litter plus moderate to rapid infiltration rates result in minimal to nonexistent runoff until soils are saturated. Basal cover is typically greater than 5% for this site and effectively reduces runoff on this site.

11. Presence and thickness of compaction layer (usually none; describe soil profile features which may be mistaken for compaction on this site): No compaction layer exists.

12. Functional/Structural Groups (list in order of descending dominance by above-ground annual-production or live foliar cover using symbols: >>, >, = to indicate much greater than, greater than, and equal to):

Dominant:

Sub-dominant:

Other:

Additional: warm season bunchgrasses>>tall, cool season bunchgrasses>>perennial forbs>mid-size, cool season bunchgrasses=cool season rhizomatous grasses=warm season rhizomatous grasses>perennial shrubs

13. Amount of plant mortality and decadence (include which functional groups are expected to show mortality or decadence): Decadence minimal to nonexistent.

14. Average percent litter cover (%) and depth (in): Litter ranges from 5-30% of total canopy measurement with total litter (including beneath the plant canopy) from 75-95% expected. Herbaceous litter depth typically ranges from 10-25 mm.

15. Expected annual annual-production (this is TOTAL above-ground annual-production, not just forage annual-production): English: 2000-2800 lb/ac (2500 lb/ac average); Metric: 2240-3136 kg/ha (2800 kg/ha average).

16. Potential invasive (including noxious) species (native and non-native). List species which BOTH characterize degraded states and have the potential to become a dominant or co-dominant species on the ecological site if their future establishment and growth is not actively controlled by management interventions. Species that become dominant for only one to several years (e.g., short-term response to drought or wildfire) are not invasive plants. Note that unlike other indicators, we are describing what is NOT expected in the reference state for the ecological site: Bare ground greater than 30% is the most common indicator of a threshold being crossed. Alkali muhly, inland saltgrass, and alkali bluegrass are common increasers. Annual weeds such as halogeton, kochia, and Russian thistle are common invasive species in disturbed sites.

17. Perennial plant reproductive capability: All species are capable of reproducing, except in drought years.
