

Ecological site R034AY260WY

Shallow Igneous Foothills and Basins West (SwIg)

Accessed: 08/21/2026

General information

Provisional. A provisional ecological site description has undergone quality control and quality assurance review. It contains a working state and transition model and enough information to identify the ecological site.

Associated sites

R034AY258WY	Shallow Clayey Foothills and Basins West (SwCy) Shallow Clayey
R034AY262WY	Shallow Loamy Foothills and Basins West (SwLy) Shallow Loamy

Similar sites

R034AY216WY	Igneous Foothills and Basins West (Ig) Igneous 10-14W has lower production and does not have bitterbrush as a major woody component.
R034AY263WY	Shallow Loamy Calcareous Foothills and Basins West (SwLyCa) Shallow Loamy, calcareous 10-14W does not have bitterbrush as a major woody component.
R043BY260WY	Shallow Igneous Foothills and Mountains West Shallow Igneous 15-19W has higher production.

Table 1. Dominant plant species

Tree	Not specified
Shrub	Not specified
Herbaceous	Not specified

Physiographic features

This site is found in an upland position on rolling to steep slopes and ridges. Slopes are commonly 15 to 30%.

Table 2. Representative physiographic features

Landforms	(1) Escarpment (2) Ridge
Flooding frequency	None
Ponding frequency	None
Elevation	1,980 – 2,290 m
Slope	10 – 70 %
Ponding depth	0 cm
Aspect	Aspect is not a significant factor

Climatic features

Annual precipitation ranges from 10-14 inches per year. Wide fluctuations may occur in yearly precipitation and result in more dry years than those with more than normal precipitation. Temperatures show a wide range between summer and winter and between daily maximums and minimums. This is predominantly due to the high elevation and dry air, which permits rapid incoming and outgoing radiation. Cold air outbreaks in winter move rapidly from northwest to southeast and account for extreme minimum temperatures. Extreme storms may occur during the winter, but most severely affect ranch operations during late winter and spring.

Daytime winds are generally stronger than nighttime and occasional strong storms may bring brief periods of high winds with gusts to more than 50 mph.

Growth of native cool season plants begins about April 15 and continues to about August 15. Some green up of cool season plants usually occurs in September depending upon fall moisture occurrences.

For detailed information visit the Natural Resources Conservation Service National Water and Climate Center at <http://www.wcc.nrcs.usda.gov/cgibin/state.pl?state=wy> website. Other climate stations representative of this precipitation zone include "Border 3 N " and Kemmerer Wtr Trtmt" in Lincoln County; "Evanston 1 E" in Uinta County; and "Merna" in Sublette County.

Table 3 Representative climatic features

Frost-free period (average)	70 days
Freeze-free period (average)	100 days
Precipitation total (average)	360 mm

Influencing water features

There are no water features associated with this site.

Soil features

These shallow, well drained soils are on bedrock-controlled mountain sides and foothills. Coarse fragment percentage is commonly more than 35 percent, but may be less in the surface layer. Parent material is residuum and colluvium from granite and schist with the soil texture modifiers of gravelly cobbly, and very cobbly.

Subsurface Texture Group: very cobbly sandy loam

Table 4. Representative soil features

Surface texture	(1) Gravelly loam (2) Gravelly sandy loam (3) Cobbly sandy loam
Family particle size	(1) Loamy
Drainage class	Well drained to somewhat excessively drained
Permeability class	Moderate to moderately rapid
Soil depth	30 – 50 cm
Surface fragment cover ≤3"	10 – 30 %
Surface fragment cover >3"	20 – 50 %
Available water capacity (0-101.6cm)	1.52 – 3.05 cm
Calcium carbonate equivalent (0-101.6cm)	Not specified
Soil reaction (1:1 water) (0-101.6cm)	5.6 – 7.2
Subsurface fragment volume ≤3" (Depth not specified)	20 – 30 %
Subsurface fragment volume >3" (Depth not specified)	30 – 50 %

Ecological dynamics

As this site deteriorates, species such as big sagebrush and green rabbitbrush will increase. Bluebunch wheatgrass, needleandthread, Indian ricegrass, and bitterbrush will decrease in frequency and production.

These plant communities narratives may not represent every possibility, but they probably are the most prevalent and repeatable plant communities. The plant composition tables shown above have been developed from the best available knowledge at the time of this revision. As more data is collected, some of these plant communities may be revised or removed, and new ones may be added. None of these plant communities should necessarily be thought of as “Desired Plant Communities”. According to the USDA NRCS National Range and Pasture Handbook, Desired Plant Communities (DPC’s) will be determined by the decision-makers and will meet minimum quality criteria established by the NRCS. The main purpose for including any description of a plant community here is to capture the current knowledge and experience at the time of this revision.

The Historic Climax Plant Community (description follows the plant community diagram) has been determined by study of rangeland relic areas, or areas protected from excessive disturbance. Trends in plant communities going from heavily grazed areas to lightly grazed areas, seasonal use pastures, and historical accounts have also been used.

The following is a State and Transition Model Diagram that illustrates the common plant communities (states) that can occur on the site and the transitions between these communities. The ecological processes will be discussed in more detail in the plant community narratives following the diagram.

State and transition model

Additional community tables

Table 5. Community 1.1 plant community composition

Group	Common Name	Symbol	Scientific Name	Annual Production ()	Foliar Cover (%)
Grass/Grasslike					
1				179-314	
	western wheatgrass	PASM	<i>Pascopyrum smithii</i>	179-314	–
2				45-90	
	bluebunch wheatgrass	PSSP6	<i>Pseudoroegneria spicata</i>	45-90	–
3				45-90	
	needle and thread	HECO26	<i>Hesperostipa comata</i>	45-90	–
4				45-90	
	Indian ricegrass	ACHY	<i>Achnatherum hymenoides</i>	45-90	–
5				9-90	
	Letterman's needlegrass	ACLE9	<i>Achnatherum lettermanii</i>	9-90	–
6				90-179	
	Grass, perennial	2GP	<i>Grass, perennial</i>	0-45	–
	needleleaf sedge	CADU6	<i>Carex duriuscula</i>	0-45	–
	plains reedgrass	CAMO	<i>Calamagrostis montanensis</i>	0-45	–
	squirreltail	ELEL5	<i>Elymus elymoides</i>	0-45	–
	prairie Junegrass	KOMA	<i>Koeleria macrantha</i>	0-45	–
	muttongrass	POFE	<i>Poa fendleriana</i>	0-45	–
	Sandberg bluegrass	POSE	<i>Poa secunda</i>	0-45	–
Forb					
7				45-90	
	Forb, perennial	2FP	<i>Forb, perennial</i>	0-45	–
	common yarrow	ACMI2	<i>Achillea millefolium</i>	0-45	–
	rosy pussytoes	ANRO2	<i>Antennaria rosea</i>	0-45	–
	milkvetch	ASTRA	<i>Astragalus</i>	0-45	–
	Indian paintbrush	CASTI2	<i>Castilleja</i>	0-45	–

	pale bastard toadflax	COUMP	<i>Comandra umbellata ssp. pallida</i>	0-45	–
	tapertip hawksbeard	CRAC2	<i>Crepis acuminata</i>	0-45	–
	fleabane	ERIGE2	<i>Erigeron</i>	0-45	–
	buckwheat	ERIOG	<i>Eriogonum</i>	0-45	–
	aster	EUCEP2	<i>Eucephalus</i>	0-45	–
	granite prickly phlox	LIPU11	<i>Linanthus pungens</i>	0-45	–
	stoneseed	LITHO3	<i>Lithospermum</i>	0-45	–
	bluebells	MERTE	<i>Mertensia</i>	0-45	–
	nailwort	PARON	<i>Paronychia</i>	0-45	–
	spiny phlox	PHHO	<i>Phlox hoodii</i>	0-45	–
	stonecrop	SEDUM	<i>Sedum</i>	0-45	–
	stemless mock goldenweed	STAC	<i>Stenotus acaulis</i>	0-45	–
	clover	TRIFO	<i>Trifolium</i>	0-45	–
Shrub/Vine					
8				45-135	
	antelope bitterbrush	PUTR2	<i>Purshia tridentata</i>	45-135	–
9				45-90	
	black sagebrush	ARNO4	<i>Artemisia nova</i>	45-90	–
10				45-90	
	big sagebrush	ARTR2	<i>Artemisia tridentata</i>	0-45	–
	yellow rabbitbrush	CHVI8	<i>Chrysothamnus viscidiflorus</i>	0-45	–
	winterfat	KRLA2	<i>Krascheninnikovia lanata</i>	0-45	–
	skunkbush sumac	RHTR	<i>Rhus trilobata</i>	0-45	–
	western snowberry	SYOC	<i>Symphoricarpos occidentalis</i>	0-45	–

Table 6. Community 2.1 plant community composition

Group	Common Name	Symbol	Scientific Name	Annual Production ()	Foliar Cover (%)
-------	-------------	--------	-----------------	----------------------	------------------

Table 7. Community 3.1 plant community composition

Group	Common Name	Symbol	Scientific Name	Annual Production ()	Foliar Cover (%)
-------	-------------	--------	-----------------	----------------------	------------------

Table 8. Community 4.1 plant community composition

Group	Common Name	Symbol	Scientific Name	Annual Production ()	Foliar Cover (%)
-------	-------------	--------	-----------------	----------------------	------------------

Animal community

Animal Community – Wildlife Interpretations Bitterbrush/Rhizomatous Wheatgrass Plant Community (HCPC): Suitable thermal and escape cover for most wildlife is limited due to the low height and density of woody plants. Bitterbrush and sagebrush provide important winter forage for mule deer and elk. Year-round habitat is provided for sage grouse and many other sagebrush obligate species such as the sage sparrow, Brewer's sparrow, sage thrasher, pygmy rabbit, sagebrush vole, horned lizard, and pronghorn antelope. Bitterbrush/Black Sagebrush Plant Community: This plant community may be beneficial for the same wildlife that would use the Historic Climax Plant Community. However, the plant community composition is less diverse, and thus, less apt to meet the seasonal needs of these animals. Big Sagebrush/Rhizomatous Wheatgrass Plant Community: This plant community may be beneficial for the same wildlife that would use the Historic Climax Plant Community. However, the plant community composition is less diverse, and thus, less apt to meet the seasonal needs of these animals. Cheatgrass Plant Community: This plant community exhibits a low level of plant species diversity. In most cases it is not a desirable plant community to select as a wildlife habitat management objective. Animal Community – Grazing Interpretations The following table lists suggested stocking rates for cattle under continuous season-long grazing under normal growing conditions. These are conservative estimates that should be used only as guidelines in the initial stages of the conservation planning process. Often, the current plant composition does not entirely match any particular plant community (as described in this ecological site description). Because of this, a field visit is recommended, in all cases, to document plant composition and production. More precise carrying capacity estimates should eventually be calculated using this information along with animal preference data, particularly when grazers other than cattle are involved. Under more intensive grazing management, improved harvest efficiencies can

result in an increased carrying capacity. Plant Community Production (lb./ac) and Carrying Capacity* (AUM/ac) Bitterbrush/Rhizomatous Wheatgrass (HCPC) 600-1100 lb./ac and .17 AUM/ac Bitterbrush/Black Sagebrush 400-900 lb./ac and .14 AUM/ac Big Sagebrush/Rhizomatous Wheatgrass 200-600 lb./ac and .07 AUM/ac Cheatgrass 50-250 lb./ac and .02 AUM/ac * - Continuous, season-long grazing by cattle under average growing conditions. Grazing by domestic livestock is one of the major income-producing industries in the area. Rangeland in this area may provide yearlong forage for cattle, sheep, or horses. During the dormant period, the forage for livestock use needs to be supplemented with protein because the quality does not meet minimum livestock requirements.

Hydrological functions

Water is the principal factor limiting forage production on this site. This site is highly variable and is dominated by soils in hydrologic group B and C, with localized areas in hydrologic group D. Infiltration ranges from slow to very rapid. Runoff potential for this site varies from moderate to high depending on soil hydrologic group, depth and degree of bedrock fracturing, slope, and ground cover (refer to Part 630, NRCS National Engineering Handbook for detailed hydrology information). Rills and gullies may be present, but should be small. Water flow patterns should be barely distinguishable. Pedestals are only slightly present in association with bunchgrasses such as bluebunch wheatgrass. Litter typically falls in place, and signs of movement are not common. Chemical and physical crusts are rare to non-existent. Cryptogammic crusts are present, but only cover 1-2% of the soil surface.

Recreational uses

This site provides hunting opportunities for upland game species.

Wood products

No appreciable wood products are present on the site.

Other products

None noted.

Inventory data references

Information presented here has been derived from NRCS clipping data and other inventory data. Field observations from range trained personnel were also used. Those involved in developing this site include: Bill Christensen, Range Management Specialist, NRCS; Karen Clause, Range Management Specialist, NRCS; and Everet Bainter, Range Management Specialist, NRCS. Other sources used as references include: USDA NRCS Water and Climate Center, USDA NRCS National Range and Pasture Handbook, and USDA NRCS Soil Surveys from various counties.

Contributors

Karen Clause

Rangeland health reference sheet

Interpreting Indicators of Rangeland Health is a qualitative assessment protocol used to determine ecosystem condition based on benchmark characteristics described in the Reference Sheet. A suite of 17 (or more) indicators are typically considered in an assessment. The ecological site(s) representative of an assessment location must be known prior to applying the protocol and must be verified based on soils and climate. Current plant community cannot be used to identify the ecological site.

Author(s)/participant(s)	K. Clause, E. Bainter
Contact for lead author	karen.clause@wy.usda.gov or 307-367-2257
Date	03/16/2007
Approved by	

Approval date	
Composition (Indicators 10 and 12) based on	Annual Production

Indicators

- 1. Number and extent of rills:** Some expected on this site. Where present, short and widely spaced.

- 2. Presence of water flow patterns:** Some can be observed.

- 3. Number and height of erosional pedestals or terracettes:** Rare to slight.

- 4. Bare ground from Ecological Site Description or other studies (rock, litter, lichen, moss, plant canopy are not bare ground):**
Bare ground can range from 20-50%.

- 5. Number of gullies and erosion associated with gullies:** Active gullies should not be present.

- 6. Extent of wind scoured, blowouts and/or depositional areas:** Rare to nonexistent.

- 7. Amount of litter movement (describe size and distance expected to travel):** Herbaceous litter expected to move in small amounts down-slope. Large woody debris from shrubs will show no movement.

- 8. Soil surface (top few mm) resistance to erosion (stability values are averages - most sites will show a range of values):** Soil Stability Index ratings range from 1 (interspaces) to 6 (under plant canopy), but average values should be 2.5 or greater.

- 9. Soil surface structure and SOM content (include type of structure and A-horizon color and thickness):** Soil data is limited for this site. Described A-horizons are about 4 inches (10 cm) with OM of .5 to 1%.

- 10. Effect of community phase composition (relative proportion of different functional groups) and spatial distribution on infiltration and runoff:** Plant community consists of 25-85% grasses, 10% forbs, and 10-65% shrubs. Unevenly distributed plant canopy (30-60%) and litter, but moderate to rapid infiltration rates result in slight to minimal runoff. Basal cover is typically less than 5% for this site and does very little to effect runoff on this site.

11. Presence and thickness of compaction layer (usually none; describe soil profile features which may be mistaken for compaction on this site): None.
-

12. Functional/Structural Groups (list in order of descending dominance by above-ground annual-production or live foliar cover using symbols: >>, >, = to indicate much greater than, greater than, and equal to):

Dominant:

Sub-dominant:

Other:

Additional: Mid-size, cool season bunchgrasses> perennial shrubs=cool season rhizomatous grasses>>perennial forbs>short, cool season bunchgrasses

13. Amount of plant mortality and decadence (include which functional groups are expected to show mortality or decadence): Minimal decadence, typically associated with shrub component.
-

14. Average percent litter cover (%) and depth (in): Litter ranges from 5-20% of total canopy measurement with total litter (including beneath the plant canopy) from 20-60% expected. Herbaceous litter depth typically ranges from 3-10mm. Woody litter can be up to a couple inches (4-6 cm).
-

15. Expected annual annual-production (this is TOTAL above-ground annual-production, not just forage annual-production): English: 700-1200 lb/ac (900 lb/ac average); Metric 784-1344 kg/ha (1008 kg/ha average).
-

16. Potential invasive (including noxious) species (native and non-native). List species which BOTH characterize degraded states and have the potential to become a dominant or co-dominant species on the ecological site if their future establishment and growth is not actively controlled by management interventions. Species that become dominant for only one to several years (e.g., short-term response to drought or wildfire) are not invasive plants. Note that unlike other indicators, we are describing what is NOT expected in the reference state for the ecological site: Bare ground greater than 75% and the presence of Cheatgrass are the most common indicators of a threshold being crossed. Rabbitbrush, Wyoming big sagebrush, Sandberg bluegrass, and goldenweed are common increasers. Annual weeds such as Cheatgrass and mustards are common invasive species in disturbed sites.
-

17. Perennial plant reproductive capability: All species are capable of reproducing, except in drought years.
-