

Ecological site R035XG124NM Hills

Last updated: 5/19/2025
Accessed: 06/10/2026

General information

Provisional. A provisional ecological site description has undergone quality control and quality assurance review. It contains a working state and transition model and enough information to identify the ecological site.

Ecological site concept

This site is characterized by rolling to steep hills and mountain footslopes. Slopes average 25 percent or more but range in extremes from 15 percent to 75 percent. Soils characterizing this site are typically shallow over acid igneous bedrock, although pockets of deeper soils also occur in saddles, between ledges, and lower side slopes. They may be loams, clay loams, or sandy loams, and are usually stony, gravelly or cobbly.

Table 1. Dominant plant species

Tree	(1) <i>Juniperus monosperma</i> (2) <i>Pinus edulis</i>
Shrub	(1) <i>Quercus gambelii</i> (2) <i>Cercocarpus montanus</i>
Herbaceous	(1) <i>Bouteloua gracilis</i> (2) <i>Schizachyrium scoparium</i>

Physiographic features

This site is characterized by rolling to steep hills and mountain footslopes. Slopes average 25 percent or more but range in extremes from 15 percent to 75 percent. Exposure or direction of slope is variable. Rock outcrops, exposed ledges, and occasional boulders are not uncommon. Elevations range from about 6000 to 7800 feet.

Table 2. Representative physiographic features

Landforms	(1) Hill (2) Mountain slope (3) Scarp slope
Flooding frequency	None
Ponding frequency	None
Elevation	1,830 – 2,380 m

Slope	20 – 80 %
Water table depth	180 cm
Aspect	Aspect is not a significant factor

Climatic features

Average annual precipitation varies from about 10 inches to just over 16 inches. Fluctuations ranging from about 5 inches to 25 inches are not uncommon. The overall climate is characterized by cold dry winters in which winter moisture is less than summer. As much as half or more of the annual precipitation can be expected to come during the period of July through September. Thus, fall conditions are often more favorable for good growth of cool-season perennial grasses, shrubs, and forbs than are those of spring.

The average frost-free season is about 120 days and extends from approximately mid-May to early or mid-September. Average annual air temperatures are 50 degrees F or lower and summer maximums rarely exceed 100 degrees F. Winter minimums typically approach or go below zero. Monthly mean temperatures exceed 70 degrees F for the period of July and August.

Rainfall patterns generally favor warm-season perennial vegetation, while the temperature regime tends to favor cool-season vegetation. This creates a somewhat complex community of plants on a given range site which is quite susceptible to disturbance and is at or near its productive potential only when both natural warm- and cool- season dominants are present.

Table 3 Representative climatic features

Frost-free period (average)	170 days
Freeze-free period (average)	250 days
Precipitation total (average)	410 mm

Influencing water features

This is an upland site, and is not associated with water features or wetlands. During heavy rain events, this site may receive run-on moisture from landforms above and contribute runoff to landforms below.

Soil features

Soils characterizing this site are typically shallow over acid igneous bedrock, although pockets of deeper soils also occur in saddles, between ledges, and lower side slopes. They may be loams, clay loams, or sandy loams, and are usually stony, gravelly or cobbly. Permeability is moderate to moderately slow, and the available water capacity is low due to shallow depth. Characteristic soils are Chimayo stony loam, Dusty gravelly loam, and Santa Fe very gravelly loam.

Table 4. Representative soil features

Surface texture	(1) Stony loam (2) Gravelly clay loam (3) Cobbly sandy loam
Family particle size	(1) Loamy

Drainage class	Well drained to excessively drained
Permeability class	Very slow to moderately rapid
Soil depth	30 – 180 cm
Surface fragment cover <=3"	20 – 30 %
Surface fragment cover >3"	20 – 30 %
Available water capacity (0-101.6cm)	2.54 – 12.7 cm
Calcium carbonate equivalent (0-101.6cm)	0 – 10 %
Electrical conductivity (0-101.6cm)	Not specified
Sodium adsorption ratio (0-101.6cm)	0 – 10
Soil reaction (1:1 water) (0-101.6cm)	6.1 – 8.4
Subsurface fragment volume <=3" (Depth not specified)	10 – 40 %
Subsurface fragment volume >3" (Depth not specified)	10 %

Ecological dynamics

Deterioration of the potential plant community due to inadequately managed grazing is most often typified by a decline in such desirable plants as sideoats grama, black grama, cool-season grasses, mountainmahogany, and winterfat. As these plants decline, they are replaced by pinyon, juniper, broom snakeweed, and lesser-value grasses such as threeawns and galleta. Because mechanical seeding and brush control are seldom justifiable on this site, the mixed use of both browsing and grazing kinds of livestock may be the best means of maintaining a healthy balance of woody and herbaceous vegetation.

State and transition model

Additional community tables

Table 5. Community 1.1 plant community composition

Group	Common Name	Symbol	Scientific Name	Annual Production ()	Foliar Cover (%)
Grass/Grasslike					
1				112-146	
	blue grama	BOGR2	<i>Bouteloua gracilis</i>	108-143	–
2				146-179	
	needle and thread	HECO26	<i>Hesperostipa comata</i>	143-178	–
	New Mexico feathergrass	HENE5	<i>Hesperostipa neomexicana</i>	143-178	–
	bullgrass	MUEM	<i>Muhlenbergia emersleyi</i>	143-178	–
	mountain muhly	MUMO	<i>Muhlenbergia montana</i>	143-178	–
	little bluestem	SCSC	<i>Schizachyrium scoparium</i>	143-178	–
3				0-34	
	black grama	BOER4	<i>Bouteloua eriopoda</i>	0-36	–
4				67-112	
	threeawn	ARIST	<i>Aristida</i>	72-108	–
	pine dropseed	BLTR	<i>Blepharoneuron tricholepis</i>	72-108	–
	squirreltail	ELEL5	<i>Elymus elymoides</i>	72-108	–
	Arizona fescue	FEAR2	<i>Festuca arizonica</i>	72-108	–
	sheep fescue	FEOV	<i>Festuca ovina</i>	72-108	–
5				34-67	
	sideoats grama	BOCU	<i>Bouteloua curtipendula</i>	36-72	–
6				11-34	
	threeawn	ARIST	<i>Aristida</i>	7-36	–
	common wolfstail	LYPH	<i>Lycurus phleoides</i>	7-36	–
	spike muhly	MUWR	<i>Muhlenbergia wrightii</i>	7-36	–
	James' galleta	PLJA	<i>Pleuraphis jamesii</i>	7-36	–
	sand dropseed	SPCR	<i>Sporobolus cryptandrus</i>	7-36	–
7				11-34	
	prairie Junegrass	KOMA	<i>Koeleria macrantha</i>	7-36	–
	green sprangletop	LEDU	<i>Leptochloa dubia</i>	7-36	–
	western wheatgrass	PASM	<i>Pascopyrum smithii</i>	7-36	–
Tree					
8				34-112	
	juniper	JUNIP	<i>Juniperus</i>	36-108	–
	twoneedle pinyon	PIED	<i>Pinus edulis</i>	36-108	–
Shrub/Vine					
9				11-67	
	Fendler's ceanothus	CEFE	<i>Ceanothus fendleri</i>	7-72	–
	mountain mahogany	CERCO	<i>Cercocarpus</i>	7-72	–
	Mexican cliffrose	PUME	<i>Purshia mexicana</i>	7-72	–
	antelope bitterbrush	PUTR2	<i>Purshia tridentata</i>	7-72	–
	oak	QUERC	<i>Quercus</i>	7-72	–
	skunkbush sumac	RHTR	<i>Rhus trilobata</i>	7-72	–
10				11-22	
	Apache plume	FAPA	<i>Fallugia paradoxa</i>	7-21	–

	broom snakeweed	GUSA2	<i>Gutierrezia sarothrae</i>	7-21	-
	winterfat	KRLA2	<i>Krascheninnikovia lanata</i>	7-21	-
Forb					
11				21-34	
	Forb, perennial	2FP	<i>Forb, perennial</i>	21-36	-
12				11-22	
	Forb, annual	2FA	<i>Forb, annual</i>	7-21	-

Table 6. Community 1.2 plant community composition

Group	Common Name	Symbol	Scientific Name	Annual Production ()	Foliar Cover (%)
-------	-------------	--------	-----------------	----------------------	------------------

Animal community

This range site provides habitats which support a resident animal community that is characterized by mule deer, gray fox, bobcat, desert cottontail, cliff chipmunk, white-hroated woodrat, pinyon mouse, harlequin quail, red-shafted flicker, scrub jay, pinyon jay, common raven, bridled titmouse, common bushtit, rufous-sided towhee, chipping sparrow, Eastern fence lizard, plateau whiptail, tree lizard, desert horned lizard, mountain patch-nose snake, and black-tailed rattlesnake.

Hydrological functions

Hydrology Functions: The runoff curve numbers are determined by field investigations using hydrologic cover conditions and hydrologic soil groups. Hydrologic Interpretations Soil Series-----Hydrologic Group Chimayo-----D

Recreational uses

This site offers good potential for hiking, horseback riding, hunting, nature observation, and photography. It has low to moderate potential for improved picnicking and camping sites, depending upon how steep the topography is. It provides natural beauty typical of the mountain foothills of the area in which it is formed.

Wood products

This site has a limited potential for wood products that is restricted almost entirely to fence posts and firewood production.

Other products

This site is well suited for grazing by multiple kinds and classes of livestock. Where slopes are steep, however, accessibility may become limited and stocking rates need to be properly adjusted. Deterioration of the potential plant community due to inadequately managed grazing is most often typified by a decline in such desirable plants as sideoats grama, black grama, cool-season grasses, mountainmahogany, and winterfat. As these plants decline, they are replaced by pinyon, juniper, broom snakeweed, and lesser-value grasses such as threeawns and galleta. Because mechanical seeding and brush control are seldom justifiable on this site, the mixed use of both browsing and grazing kinds of livestock may be the best means of maintaining a healthy balance of woody and herbaceous vegetation.

Other information

Guide to Suggested Initial Stocking Rate Acres per Animal Unit Month Similarity Index-----Ac/AUM 100 - 76-----3.6 - 4.8
75 - 51-----4.5 - 6.5 50 - 26-----6.0 - 12.5 25 - 0-----12.5 +

Contributors

- Brenda Simpson
- Christine Bishop
- Don Sylvester
- John Tunberg

Approval

Kendra Moseley, 5/19/2025

Rangeland health reference sheet

Interpreting Indicators of Rangeland Health is a qualitative assessment protocol used to determine ecosystem condition based on benchmark characteristics described in the Reference Sheet. A suite of 17 (or more) indicators are typically considered in an assessment. The ecological site(s) representative of an assessment location must be known prior to applying the protocol and must be verified based on soils and climate. Current plant community cannot be used to identify the ecological site.

Author(s)/participant(s)	
Contact for lead author	
Date	06/10/2026
Approved by	
Approval date	
Composition (Indicators 10 and 12) based on	Annual Production

Indicators

1. Number and extent of rills:

2. Presence of water flow patterns:

3. Number and height of erosional pedestals or terracettes:

4. Bare ground from Ecological Site Description or other studies (rock, litter, lichen, moss, plant canopy are not bare ground):

5. Number of gullies and erosion associated with gullies:

6. Extent of wind scoured, blowouts and/or depositional areas:

7. Amount of litter movement (describe size and distance expected to travel):

8. Soil surface (top few mm) resistance to erosion (stability values are averages - most sites will show a range of values):

9. Soil surface structure and SOM content (include type of structure and A-horizon color and thickness):

10. Effect of community phase composition (relative proportion of different functional groups) and spatial distribution on infiltration and runoff:

11. Presence and thickness of compaction layer (usually none; describe soil profile features which may be mistaken for compaction on this site):

12. Functional/Structural Groups (list in order of descending dominance by above-ground annual-production or live foliar cover using symbols: >>, >, = to indicate much greater than, greater than, and equal to):

Dominant:

Sub-dominant:

Other:

Additional:

13. Amount of plant mortality and decadence (include which functional groups are expected to show mortality or decadence):

14. Average percent litter cover (%) and depth (in):

15. Expected annual annual-production (this is TOTAL above-ground annual-production, not just forage annual-production):

16. Potential invasive (including noxious) species (native and non-native). List species which BOTH characterize degraded states and have the potential to become a dominant or co-dominant species on the ecological site if their future establishment and growth is not actively controlled by management interventions. Species that become dominant for only one to several years (e.g., short-term response to drought or wildfire) are not invasive plants. Note that unlike other indicators, we are describing what is NOT expected in the reference state for the ecological site:

17. Perennial plant reproductive capability:
