

Ecological site R042AD003NM

Draw, Dry Mixed Prairie

Accessed: 09/16/2026

General information

Provisional. A provisional ecological site description has undergone quality control and quality assurance review. It contains a working state and transition model and enough information to identify the ecological site.

Table 1. Dominant plant species

Tree	Not specified
Shrub	Not specified
Herbaceous	Not specified

Physiographic features

This site formed in calcareous alluvium and are on inset fans of fan piedmonts, draws or drainageways. The soils are frequently overflowed or receive runoff additional moisture from adjoining sites. Slopes range from 1 to 5 percent. Elevations range from 4700 to 6000 feet. Mean annual precipitation is about 13 inches and the mean annual temperature is about 60 degrees F.

Table 2. Representative physiographic features

Landforms	(1) Inset fan (2) Drainageway (3) Draw
Flooding duration	Very brief (4 to 48 hours)
Flooding frequency	Rare to occasional
Ponding frequency	None
Elevation	1,430 – 1,830 m
Slope	0 – 10 %
Aspect	Aspect is not a significant factor

Climatic features

Average precipitation for this site is approximately 12 to 14 inches. Variations of 5 inches are not uncommon. Approximately 75 percent occurs from May through October with most of the rainfall occurring from July to September. Most of the summer precipitation comes in the form of high intensity, short duration thunderstorms. Rain and snow of low intensity characterize the limited winter precipitation.

Temperatures are mild. Freezing temperatures are common at night from December through April, however, temperatures during the day are frequently above 50 degrees F. Occasionally in December to February brief periods of 0 degrees F. temperatures may be expected. During June to August some days may exceed 100 degrees F.

The mean annual precipitation figures are derived from rain gauge data collected by the BLM (1971 to 1990), and NOAA weather maps utilizing prism model estimation techniques. There are no permanent weather stations within the boundaries of the Land Resource Unit.

Table 3 Representative climatic features

Frost-free period (average)	180 days
Freeze-free period (average)	190 days
Precipitation total (average)	360 mm

Influencing water features

This site is not influenced by water from wetland or stream.

Soil features

The soils of this site are very deep. Surface texture is loam, or silt loam with less than 15 percent calcium carbonate. Upper subsoil textures are extremely gravelly loam, extremely gravelly clay loam, or extremely gravelly sandy clay loam with 20 to 40 percent calcium carbonate. Lower subsoil textures are loam, silt loam, clay loam, or silty clay loam with more than 40 percent calcium carbonate.

Minimum and maximum values listed below represent the characteristic soil for this site.

Characteristic Soil:

Oryx

Table 4. Representative soil features

Surface texture	(1) Loam
Family particle size	(1) Loamy
Drainage class	Well drained
Permeability class	Moderately slow
Soil depth	150 – 200 cm
Surface fragment cover <=3"	0 – 10 %

Surface fragment cover >3"	Not specified
Available water capacity (0-101.6cm)	4.32 – 7.87 cm
Calcium carbonate equivalent (0-101.6cm)	0 – 40 %
Electrical conductivity (0-101.6cm)	Not specified
Sodium adsorption ratio (0-101.6cm)	Not specified
Soil reaction (1:1 water) (0-101.6cm)	7.9 – 8.4
Subsurface fragment volume <=3" (Depth not specified)	0 – 10 %
Subsurface fragment volume >3" (Depth not specified)	0 – 30 %

Ecological dynamics

This site is associated with Limestone Hills, Igneous Hills, and Gravelly sites from which it receives and transports runoff water. This ecological site consists of two separate elements, the arroyo channel and its associated flood plain. The Draw site is an ephemeral stream floodplain with a gently sloping surface, broad enough that the channel covers only a part of the surface.

Along the channel it has the appearance of an elongated sinuous savannah with shrubs and trees dominant in aspect, however, grasses have the greatest annual production. There are an abundant variety of forbs that can occur in the understory. Vegetation is variable and is dependent on flood events, distance away from the channel, parent material, and amount of gravel and cobble in the soil profile. Sideoats grama is the dominant grass in the historic plant community. Cane bluestem, bush muhly, blue grama, and plains bristlegrass can also occur in significant numbers. Desert willow, Apache plume, brickellbush, littleleaf sumac, mariola, and mesquite are common woody species. Retrogression is characterized by a decrease in mid grasses such as sideoats grama, cane bluestem, Arizona cottontop, green sprangletop and an increase in blue grama and tobosa. Transition to the creosotebush-dominated state may occur as a result of continued loss of grass cover, erosion and lack of fire.

State and transition model

Figure 3. MLRA-42, SD-4, Draw

Additional community tables

Table 5. Community 1.1 plant community composition

Group	Common Name	Symbol	Scientific Name	Annual Production ()	Foliar Cover (%)
Grass/Grasslike					
1				605-706	
	sideoats grama	BOCU	<i>Bouteloua curtipendula</i>	404-504	–
	cane bluestem	BOBA3	<i>Bothriochloa barbinodis</i>	202-303	–
2				224-303	
	bush muhly	MUPO2	<i>Muhlenbergia porteri</i>	135-235	–
	blue grama	BOGR2	<i>Bouteloua gracilis</i>	101-168	–
3				202-242	
	big sacaton	SPWR2	<i>Sporobolus wrightii</i>	112-135	–
	tobosagrass	PLMU3	<i>Pleuraphis mutica</i>	84-112	–
	alkali sacaton	SPAI	<i>Sporobolus airoides</i>	84-112	–
4				168-202	
	plains bristlegass	SEVU2	<i>Setaria vulpiseta</i>	135-168	–
	Arizona cottontop	DICA8	<i>Digitaria californica</i>	56-84	–
	green sprangletop	LEDU	<i>Leptochloa dubia</i>	34-56	–
	spike dropseed	SPCO4	<i>Sporobolus contractus</i>	34-56	–
	sand dropseed	SPCR	<i>Sporobolus cryptandrus</i>	34-56	–
5				40-61	
	threeawn	ARIST	<i>Aristida</i>	40-61	–
	burrograss	SCBR2	<i>Scleropogon brevifolius</i>	40-61	–
Forb					
6				20-40	
	Forb, annual	2FA	<i>Forb, annual</i>	20-40	–
	common sunflower	HEAN3	<i>Helianthus annuus</i>	7-13	–
7				40-61	
	Forb, perennial	2FP	<i>Forb, perennial</i>	40-61	–
	croton	CROTO	<i>Croton</i>	20-40	–
	globemallow	SPHAE	<i>Sphaeralcea</i>	20-40	–
Shrub/Vine					
8				161-202	
	Apache plume	FAPA	<i>Fallugia paradoxa</i>	112-168	–
	brickellbush	BRICK	<i>Brickellia</i>	84-135	–
	littleleaf sumac	RHMI3	<i>Rhus microphylla</i>	56-84	–
9				40-101	
	creosote bush	LATR2	<i>Larrea tridentata</i>	40-101	–
	mariola	PAIN2	<i>Parthenium incanum</i>	40-101	–
	honey mesquite	PRGL2	<i>Prosopis glandulosa</i>	40-101	–
	goldeneye	VIGUI	<i>Viguiera</i>	13-27	–
Tree					
10				84-101	
	desert willow	CHLI2	<i>Chilopsis linearis</i>	84-101	–

Table 6. Community 2.1 plant community composition

Group	Common Name	Symbol	Scientific Name	Annual Production ()	Foliar Cover (%)
-------	-------------	--------	-----------------	----------------------	------------------

Table 7. Community 2.2 plant community composition

Group	Common Name	Symbol	Scientific Name	Annual Production ()	Foliar Cover (%)
-------	-------------	--------	-----------------	----------------------	------------------

Animal community

The Draw ecological sites exhibit high structural and vegetation diversity and provide thermal, nesting and hiding cover for many wildlife species. The site provides important high protein forage and browse and serves as corridors and conduits for water and wildlife movement. The draw sites are characterized by a diversity of species. A few of the species utilizing this site are mule deer, coyote, bobcat, fox, roadrunner, great-horned owl, northern mockingbird, gambel's quail and Texas horned lizard.

Hydrological functions

This site normally receives approximately 12-14 inches annual precipitation. Most summer rainfall occurs as brief sometimes-heavy thunderstorms. Soils are very deep and rated as being in hydrologic group B. Slopes range from 1- 5 percent. Permeability is moderately slow. Runoff is medium, and the hazard of water erosion is moderate. Available water capacity to a depth of 40 inches is very low to low. Soil Hydrologic Unit Oryx B

Recreational uses

This site offers recreation potential for hiking, horseback riding, camping, outdoor classrooms, nature observation, photography and hunting.

Wood products

This site has no significant value for wood products.

Other products

Grazing: The plant community on this site is well suited to grazing by both domestic livestock of all kinds and by wildlife at all seasons of the year. It is well suited to winter and early spring grazing, due to high protein levels in the forage. This site will respond well to a grazing system that rotates the season of use. When grazing sheep or goats and during calving season, predator control should be considered. Initial starting stocking rates will be determined with the landowner or decision-maker. They will be based on past use histories and type and condition of the vegetation. Calculations used to determine and initial starting stocking rate will also be based on forage preference ratings.

Type locality

Location 1: Bernalillo County, NM	
Township/Range/Section	T21 S. R12 E. S26
Latitude	32° 26'42"
Longitude	105° 41'17"
General legal description	Otero County, New Mexico; approximately 23.6 miles east and 4.3 miles north of Oro Grande; SW 1/4, SE 1/4 section 26, T. 21 S., R. 12 E.; USGS El Paso Draw, NM topographic quadrangle; UTM zone 13, 435289 E, 3589784 N; NAD 27.

Other references

1. U.S. Department of Agriculture, Forest Service, Rocky Mountain Research Station, Fire Sciences Laboratory (2002, September). Fire Effects Information System, [Online]. Available: <http://www.fs.fed.us/database/feis/> [accessed 9/23/02].
2. 9. U.S. Department of Agriculture, Natural Resources Conservation Service. 2001. Soil Quality Information Sheet. Rangeland Soil Quality—Physical and Biological Soil Crusts. Rangeland Sheet 7, [Online]. Available: <http://www.statlab.iastate.edu/survey/SQL/range.html>

Contributors

David Trujillo
Jason Martin

Rangeland health reference sheet

Interpreting Indicators of Rangeland Health is a qualitative assessment protocol used to determine ecosystem condition based on benchmark characteristics described in the Reference Sheet. A suite of 17 (or more) indicators are typically considered in an assessment. The ecological site(s) representative of an assessment location must be known prior to applying the protocol and must be verified based on soils and climate. Current plant community cannot be used to identify the ecological site.

Author(s)/participant(s)	
Contact for lead author	
Date	
Approved by	
Approval date	
Composition (Indicators 10 and 12) based on	Annual Production

Indicators

1. Number and extent of rills:

2. Presence of water flow patterns:

3. Number and height of erosional pedestals or terracettes:

4. Bare ground from Ecological Site Description or other studies (rock, litter, lichen, moss, plant canopy are not bare ground):

5. Number of gullies and erosion associated with gullies:

6. Extent of wind scoured, blowouts and/or depositional areas:

7. Amount of litter movement (describe size and distance expected to travel):

8. Soil surface (top few mm) resistance to erosion (stability values are averages - most sites will show a range of values):

9. Soil surface structure and SOM content (include type of structure and A-horizon color and thickness):

10. Effect of community phase composition (relative proportion of different functional groups) and spatial distribution on infiltration and runoff:

11. Presence and thickness of compaction layer (usually none; describe soil profile features which may be mistaken for compaction on this site):

12. Functional/Structural Groups (list in order of descending dominance by above-ground annual-production or live foliar cover using symbols: >>, >, = to indicate much greater than, greater than, and equal to):

Dominant:

Sub-dominant:

Other:

Additional:

13. Amount of plant mortality and decadence (include which functional groups are expected to show mortality or decadence):

14. Average percent litter cover (%) and depth (in):

15. Expected annual annual-production (this is TOTAL above-ground annual-production, not just forage annual-production):

16. Potential invasive (including noxious) species (native and non-native). List species which BOTH characterize degraded states and have the potential to become a dominant or co-dominant species on the ecological site if their future

establishment and growth is not actively controlled by management interventions. Species that become dominant for only one to several years (e.g., short-term response to drought or wildfire) are not invasive plants. Note that unlike other indicators, we are describing what is NOT expected in the reference state for the ecological site:

17. Perennial plant reproductive capability:
