

**Ecological site F043AY519WA**  
**Warm-Frigid, Xeric, Loamy Slopes, low AWC subsoils**  
**(Douglas-Fir/Warm Dry Shrub)**  
***Pseudotsuga menziesii* / *Physocarpus malvaceus* - *Symphoricarpos albus***

Last updated: 10/14/2020

Accessed: 04/21/2026

**General information**

**Provisional.** A provisional ecological site description has undergone quality control and quality assurance review. It contains a working state and transition model and enough information to identify the ecological site.

**MLRA notes**

Major Land Resource Area (MLRA): 043A–Northern Rocky Mountains

Description of MLRAs can be found in: United States Department of Agriculture, Natural Resources Conservation Service. 2006. Land Resource Regions and Major Land Resource Areas of the United States, the Caribbean, and the Pacific Basin. U.S. Department of Agriculture Handbook 296.

**LRU notes**

Most commonly found in LRU 43A03 (Columbia-Colville Valleys). This LRU is composed predominantly of low elevation valley floors and the slopes of foothills or low mountains adjacent to the Columbia and Colville rivers. The LRU is in the portion of the Northern Rocky Mountains that was subjected to continental glaciation. The soils tend to be loamy mollisols and inceptisols with mixed or thin ash surfaces. Till and outwash are the dominant parent materials. Soil climate is a mesic or frigid temperature regime and xeric moisture regime with average annual precipitation around 495 mm (19 inches) and an average annual air temperature around 8.2 degrees C (47 degrees F). Elevation ranges from about 370 to 1030 m (1,200 to 3,380 feet). Also found in adjacent areas of 43A01, 43A02, 43A04 and 44A02. Climate parameters were obtained from PRISM and other models for the area. Landscape descriptors are derived from USGS DEM products and their derivatives.

**Classification relationships**

Relationship to Other Established Classifications: United States National Vegetation Classification (2008) - A3392 Douglas fir- P. Pine / Shrub Understory Central Rocky Mt. Forest & Woodland Alliance Washington Natural Heritage Program. Ecosystems of Washington State, A Guide to Identification, Rocchio and Crawford, 2015 - Northern Rocky Mt. Dry-Mesic Montane Mixed Conifer Forest (D. Fir – Pine) Description of Ecoregions of the United States, USFS PN # 1391, 1995 - M333 Northern Rocky Mt. Forest-Steppe-Coniferous Forest-Alpine Meadow Province Level III and IV Ecoregions of WA, US EPA, June 2010 - 15x Okanogan Highland Dry Forest. 15w Western Selkirk Maritime Forest. 15r Okanogan – Colville Xeric Valleys & Foothills. This ecological site includes the following USDA Forest Service Plant Associations: PSME/PHMA, PSME/PHMA-LIBOL and PSME/SYAL (Douglas-fir Series). (Williams et. al. 1995)

**Ecological site concept**

This site consists of hillslopes and mountain slopes with the following characteristics: loamy soil materials; a water table (perched or apparent) more than 75 cm (30 in) below the soil surface during the April to October period; cumulative available water capacity to 100 cm (40 inches) of less than 7.5 cm (3 inches); PSME/PHMA or PSME/SYAL habitat type.

Table 1. Dominant plant species

Tree	(1) <i>Pseudotsuga menziesii</i> var. <i>glauca</i> (2) <i>Pinus ponderosa</i>
Shrub	(1) <i>Physocarpus malvaceus</i> (2) <i>Symphoricarpos albus</i>

Herbaceous	(1) <i>Calamagrostis rubescens</i> (2) <i>Bromus vulgaris</i>
------------	--

### Physiographic features

Physiographic Features

Landscapes: Foothills, Valleys, Mountains

Landform: sideslopes, foot slopes, lake terraces, outwash terraces

Elevation (m): Total range = 290 to 1510 m

(950 to 4,950 feet)

Central tendency = 635 to 985 m

(2,080 to 3,230 feet)

Slope (percent): Total range = 0 to 75 percent

Central tendency = 6 to 35 percent

Aspect:

68-205-338

Central tendency:

138-205-265

**Table 2. Representative physiographic features**

Hillslope profile	(1) Summit (2) Backslope
Landforms	(1) Mountains > Mountain slope (2) Foothills > Hillslope
Flooding frequency	None
Ponding frequency	None
Elevation	730 – 1,030 m
Slope	20 – 50 %
Water table depth	200 cm
Aspect	W, SE, S, SW

**Table 3. Representative physiographic features (actual ranges)**

Flooding frequency	Not specified
Ponding frequency	Not specified

Elevation	390 – 1,490 m
Slope	0 – 90 %
Water table depth	200 cm

### Climatic features

#### Climatic Features

Frost-free period (days): Total range = 105 to 140 days

Central tendency = 105 to 115 days

Mean annual precipitation (cm): Total range = 265 to 1160 mm

(10 to 47 inches)

Central tendency = 490 to 760 mm

(19 to 30 inches)

MAAT (C): Total range = 4.3 to 10.2

(40 to 50 F)

Central tendency = 6.5 to 8.0

(44 to 46 F)

Climate Stations: Newport

**Table 4 Representative climatic features**

Frost-free period (characteristic range)	110-120 days
Freeze-free period (characteristic range)	
Precipitation total (characteristic range)	480-760 mm
Frost-free period (actual range)	110-140 days
Freeze-free period (actual range)	
Precipitation total (actual range)	250-1,190 mm

### Influencing water features

Water Table Depth (cm): >200 cm

Flooding:

Frequency: None

Duration: None

Ponding:

Frequency: None

Duration: None

## Soil features

### Representative Soil Features

This ecological subsite is associated with several soil series (e.g. Treble, Thout, Oxerine, Raisio, and Mineral). The soil components can be grouped into: Vitrandic Haploxerolls, Vitrandic Haploxerepts, Vitrandic Dystroxerepts and Andic Haploxerepts. These soils have developed in mixed Mazama tephra, loess and other deposits over till, and/or residuum and colluvium from granitic and metasedimentary rock. The soils range from moderately deep to very deep and have low available water capacity to a depth of 1 m. The soils are mostly well-drained (~90% by area).

#### Parent Materials:

Kind: Tephra (volcanic ash) mixed with loess and other material

Origin: mixed

Kind: residuum and colluvium, and/or till

Origin: Granite, Metasedimentary rock

#### Surface Texture:

(1) Ashy Sandy loam

(2) Ashy Silt loam

(3) Ashy Very fine sandy loam

Fragment content of surface: 22 to 61 percent (median = 27%)

#### Subsurface Texture Group: Loamy

Fragment content of subsurface (25 to 100cm): 38 to 80 percent (median = 55%)

Surface Fragments: about 10% of components have a stony or bouldery surface

Drainage Class: Well drained

Saturated Hydraulic conductivity: Moderately high to High

Soil Depth: ~35% of components have no restriction within 150 cm

Lithic contacts when present are at 25 to 90cm (median = 60cm)

Paralithic contacts when present are at 40 to 65cm (median = 53 cm)

Densic contacts when present are at 40 to 65cm (median = 53 cm)

Calcium Carbonate Equivalent (percent): 0

Soil Reaction (1:1 Water): 6.1 to 7.8

Available Water Capacity (total in 100cm): 5.68-7.64cm (median = 6.24cm)

Table 5. Representative soil features

Parent material	(1) Volcanic ash (2) Till (3) Colluvium – granite and gneiss (4) Residuum – granite and gneiss
Surface texture	(1) Ashy very fine sandy loam (2) Ashy silt loam (3) Ashy sandy loam
Drainage class	Well drained
Permeability class	Moderately rapid

Depth to restrictive layer	80 cm
Soil depth	80 cm
Surface fragment cover >3"	Not specified
Available water capacity (0-101.6cm)	6.22 cm
Calcium carbonate equivalent (0-152.4cm)	Not specified
Soil reaction (1:1 water) (0-152.4cm)	Not specified
Subsurface fragment volume <=3" (25.4-101.6cm)	60 %

**Table 6. Representative soil features (actual values)**

Drainage class	Well drained
Permeability class	Moderate to rapid
Depth to restrictive layer	20 – 200 cm
Soil depth	20 – 200 cm
Surface fragment cover >3"	0 %
Available water capacity (0-101.6cm)	5.72 – 7.62 cm
Calcium carbonate equivalent (0-152.4cm)	0 %

Soil reaction (1:1 water) (0-152.4cm)	6.1 – 7.8
Subsurface fragment volume <=3" (25.4-101.6cm)	40 – 80 %

### Ecological dynamics

A description of vegetation dynamics and a state and transition model can be found in Ecological Site Group EX043AESG05

### State and transition model

### Additional community tables

Table 7. Representative site productivity

Common Name	Symbol	Site Index Low	Site Index High	CMAI Low	CMAI High	Age Of CMAI	Site Index Curve Code	Site Index Curve Basis	Citation
ponderosa pine	<i>PIPO</i>	90	114	85	130	40	–	–	
Rocky Mountain Douglas-fir	<i>PSMEG</i>	61	76	58	93	120	–	–	
Rocky Mountain Douglas-fir	<i>PSMEG</i>	70	85	61	92	100	–	–	
western larch	<i>LAOC</i>	65	75	64	83	50	–	–	

### Approval

Curtis Talbot, 10/14/2020

### Rangeland health reference sheet

**Interpreting Indicators of Rangeland Health** is a qualitative assessment protocol used to determine ecosystem condition based on benchmark characteristics described in the Reference Sheet. A suite of 17 (or more) indicators are typically considered in an assessment. The ecological site(s) representative of an assessment location must be known prior to applying the protocol and must be verified based on soils and climate. Current plant community cannot be used to identify the ecological site.

Author(s)/participant(s)	
Contact for lead author	
Date	04/21/2026
Approved by	
Approval date	
Composition (Indicators 10 and 12) based on	Annual Production

## Indicators

1. Number and extent of rills:

---

2. Presence of water flow patterns:

---

3. Number and height of erosional pedestals or terracettes:

---

4. Bare ground from Ecological Site Description or other studies (rock, litter, lichen, moss, plant canopy are not bare ground):

---

5. Number of gullies and erosion associated with gullies:

---

6. Extent of wind scoured, blowouts and/or depositional areas:

---

7. Amount of litter movement (describe size and distance expected to travel):

---

8. Soil surface (top few mm) resistance to erosion (stability values are averages - most sites will show a range of values):

---

9. Soil surface structure and SOM content (include type of structure and A-horizon color and thickness):

---

10. Effect of community phase composition (relative proportion of different functional groups) and spatial distribution on infiltration and runoff:

---

11. Presence and thickness of compaction layer (usually none; describe soil profile features which may be mistaken for compaction on this site):

---

12. Functional/Structural Groups (list in order of descending dominance by above-ground annual-production or live foliar cover using symbols: >>, >, = to indicate much greater than, greater than, and equal to):

Dominant:

Sub-dominant:

Other:

**Additional:**

---

**13. Amount of plant mortality and decadence (include which functional groups are expected to show mortality or decadence):**

---

**14. Average percent litter cover (%) and depth ( in):**

---

**15. Expected annual annual-production (this is TOTAL above-ground annual-production, not just forage annual-production):**

---

**16. Potential invasive (including noxious) species (native and non-native). List species which BOTH characterize degraded states and have the potential to become a dominant or co-dominant species on the ecological site if their future establishment and growth is not actively controlled by management interventions. Species that become dominant for only one to several years (e.g., short-term response to drought or wildfire) are not invasive plants. Note that unlike other indicators, we are describing what is NOT expected in the reference state for the ecological site:**

---

**17. Perennial plant reproductive capability:**

---