

Ecological site R043AY508ID

Subalpine Meadows

30-45" PZ Cryic

Okanogan Plateau

Last updated: 10/15/2020
Accessed: 08/30/2026

General information

Provisional. A provisional ecological site description has undergone quality control and quality assurance review. It contains a working state and transition model and enough information to identify the ecological site.

MLRA notes

Major Land Resource Area (MLRA): 043A–Northern Rocky Mountains

Major Land Resource Area (MLRA): 043A–Northern Rocky Mountains Description of MLRAs can be found in: United States Department of Agriculture, Natural Resources Conservation Service. 2006. Land Resource Regions and Major Land Resource Areas of the United States, the Caribbean, and the Pacific Basin. U.S. Department of Agriculture Handbook 296.

LRU notes

Major land resource area (MLRA): 043A-Northern Rocky Mountains Modal LRU – 43A01 - Okanogan Plateau This LRU is composed predominantly of glaciated mountains and foothills. The soils tend to be loamy to sandy andisols, inceptisols, and mollisols with mixed or distinct ash surfaces. Recent alluvium, till, outwash and residuum from granitic or metamorphic rock are the dominant parent materials. Soil climate is a frigid or cryic temperature regime and xeric moisture regime with average annual precipitation around 450 mm (18 inches). Others where occurring: 43A04 Selkirk Mountains 43A02 Western Selkirk Highlands

Classification relationships

This ES fits into the National Vegetation Standard's Rocky Mountain-Vancouverian Grassland & Meadow Group and is most closely related to Washington State's Natural Heritage Program's Rocky Mountain Subalpine-Montane Mesic Meadow. (Compare to previous Washington range sites: R043AY702WA, MOUNTAIN PARK 24+ PZ; R043AY704WA SUBALPINE PARK 24+ PZ,)

Ecological site concept

Ecological Site Concept: This ES is found on well drained, loamy sites on mountains with adequate water holding capacity, cryic temperatures and a plant community dominated by forbs such as fleabane, mertensia and aster.

Table 1. Dominant plant species

Tree	Not specified
Shrub	(1) <i>Dasiphora fruticosa</i> (2) <i>Symphoricarpos oreophilus</i>
Herbaceous	(1) <i>Erigeron</i> (2) <i>Aster</i>

Physiographic features

Physiographic Features

This ecological site occurs mainly on hillslopes, lower mountain slopes, outwash terraces and stream terraces. Parent materials are primarily till, outwash or colluvium and residuum from metamorphic rock with a mantle of volcanic ash and loess.

Landscapes: Mountains

Landforms: mountain slopes and avalanche chutes

Elevation:

Total range = 675 to 2200 m

(2,215 to 7,215 feet)

Central tendency = 1190 to 1660 m

(3,895 to 5,435 feet)

Slope (percent):

Total range = 0 to 80 percent

Central tendency = 25 to 45 percent

Water Table Depth:

>200cm

(>80 in)

Flooding:

Frequency: None

Duration: None

Ponding:

Frequency: None

Duration: None

Aspect: 165-190-325

Central tendency: 140-190-225

Table 2. Representative physiographic features

Landforms	(1) Mountains > Mountain slope (2) Mountains > Avalanche chute
Elevation	1,190 – 1,660 m
Slope	30 – 50 %
Water table depth	0 cm
Aspect	W, SE, S, SW

Table 3. Representative physiographic features (actual ranges)

Elevation	680 – 2,200 m
Slope	0 – 80 %
Water table depth	0 cm

Climatic features

Climatic Features

During the spring and summer, a circulation of air around a high-pressure center brings a prevailing westerly and northwesterly flow of comparatively dry, cool and stable air into the region. As the air moves inland, it becomes warmer and drier which results in a dry season beginning in the late spring and reaching a peak in mid-summer. In the fall and winter, a circulation of air around two pressure centers over the ocean brings a prevailing southwesterly and westerly flow of air into the Pacific Northwest. This air from over the ocean is moist and near the temperature of the water. Condensation occurs as the air moves inland over the cooler land and rises along the windward slopes of the mountains or highlands. This results in a wet season beginning in October, reaching a peak in winter, then gradually decreasing in the spring.

The elevation within the LRU varies from approximately 1,000 feet in the lower river valleys to about 7,200 feet in the higher mountains. The annual precipitation increases from 9 inches in the valleys to over 48 inches in the highest mountains. Winter season snowfall averages about 43 inches. Both rainfall and snowfall increase with elevation. Snow can be expected after the first of November and to remain on the ground from the first of December until March or April.

In January, the average maximum temperature is near 29° F and the minimum temperature is 14° F. Minimum temperatures from -5° to -20°F are recorded almost every winter and temperatures ranging to -25° F have been recorded. In July, the average maximum temperature is 81° and the minimum temperature 47° F. Maximum temperatures reach 100° F on a few afternoons each summer and temperatures as high as 108° F have been recorded. The average date of the last freezing temperatures can be expected by early-June and after early-September in the warmer areas.

(Compiled from WRCC: Climate of Washington and available station data)

Frost-free period (days):

Total range = 75 to 125 days

Central tendency = 90 to 105 days

Mean annual precipitation (cm):

Total range = 425 to 1590 mm

(17 to 63 inches)

Central tendency = 525 to 1120 mm

(21 to 44 inches)

MAAT (C)

Total range = 1.5 to 7.9

(35 to 46 F)

Central tendency = 3.8 to 5.8

(39 to 42 F)

Climate stations: none

Influencing water features

Water Table Depth:

>200cm

(>80 in)

Flooding:

Frequency: None

Duration: None

Ponding:

Frequency: None

Duration: None

Soil features

Representative Soil Features

This ecological site is associated with the Brickel, Crocamp, Pelican, Tenas, and Vitrandic Dystrocryepts components. The soils are Haploxerandic Humicryepts, Vitrandic Dystrocryepts, Vitrandic Haplocryolls, Vitrandic Haploxerolls, and Vitrixerandic Humicryepts. These soils have developed in till, colluvium and residuum from granite, metamorphic and andesite rocks. Most sites have a surface influenced by volcanic ash and loess. Volcanic ash layers are mixed at lower elevations and form distinct, moderately thicker layers at higher

elevations.

Parent Materials:

Kind: volcanic ash, loess, till,

Origin: mixed

Kind: colluvium, residuum

Origin: granite, metamorphics, andesite

Surface Texture: (2mm fraction)

(1) Ashy-Loam

(2) Ashy-Sandy Loam

(3) Ashy-Fine Sandy Loam

Surface Fragments

Table 4. Representative soil features

Parent material	(1) Volcanic ash (2) Loess (3) Till (4) Colluvium – igneous and metamorphic rock
Surface texture	(1) Ashy loam (2) Ashy sandy loam (3) Ashy fine sandy loam
Drainage class	Well drained
Permeability class	Moderately rapid
Depth to restrictive layer	80 – 110 cm
Surface fragment cover ≤ 3 "	Not specified
Surface fragment cover > 3 "	Not specified
Available water capacity (0-101.6cm)	9.4 cm
Calcium carbonate equivalent (0-152.4cm)	Not specified

Electrical conductivity (0-152.4cm)	Not specified
Sodium adsorption ratio (0-152.4cm)	Not specified
Soil reaction (1:1 water) (0-152.4cm)	6.1

Table 5. Representative soil features (actual values)

Drainage class	Well drained
Permeability class	Moderate to moderately rapid
Depth to restrictive layer	50 – 150 cm
Surface fragment cover <=3"	0 %
Surface fragment cover >3"	0 %
Available water capacity (0-101.6cm)	7.62 – 10.92 cm
Calcium carbonate equivalent (0-152.4cm)	0 %
Electrical conductivity (0-152.4cm)	0 mmhos/cm
Sodium adsorption ratio (0-152.4cm)	0
Soil reaction (1:1 water) (0-152.4cm)	5.6 – 7.3

Ecological dynamics

State and transition model

Figure 1. Narrative

Figure 2. STM

Additional community tables

References

. (Date accessed). USNVC [United States National Vegetation Classification]. 2019. United States National Vegetation Classification Database, V2.03. Federal Geographic Data Committee, Vegetation Subcommittee, Washington DC.. USNVC: <http://usnvc.org/>.

Gerald, R. 2004. NRCS - Washington; Interim Ecological Site Descriptions for Rangeland.

Rocchio, J.F. and R.C. Crawford. 2015. Ecological systems of Washington State. A guide to identification. Washington Department of Natural Resources.. Natural Heritage Report.. Washington Department of Natural Resources, Natural Heritage Program, Olympia, WA. 1–397.

Contributors

Stephanie Shoemaker
Brian Gardner

Approval

Curtis Talbot, 10/15/2020

Rangeland health reference sheet

Interpreting Indicators of Rangeland Health is a qualitative assessment protocol used to determine ecosystem condition based on benchmark characteristics described in the Reference Sheet. A suite of 17 (or more) indicators are typically considered in an assessment. The ecological site(s) representative of an assessment location must be known prior to applying the protocol and must be verified based on soils and climate. Current plant community cannot be used to identify the ecological site.

Author(s)/participant(s)	
Contact for lead author	
Date	08/30/2026
Approved by	
Approval date	
Composition (Indicators 10 and 12) based on	Annual Production

Indicators

1. Number and extent of rills:

2. Presence of water flow patterns:

3. Number and height of erosional pedestals or terracettes:

4. Bare ground from Ecological Site Description or other studies (rock, litter, lichen, moss, plant canopy are not bare ground):

5. Number of gullies and erosion associated with gullies:

6. Extent of wind scoured, blowouts and/or depositional areas:

7. Amount of litter movement (describe size and distance expected to travel):

8. Soil surface (top few mm) resistance to erosion (stability values are averages - most sites will show a range of values):

9. Soil surface structure and SOM content (include type of structure and A-horizon color and thickness):

10. Effect of community phase composition (relative proportion of different functional groups) and spatial distribution on infiltration and runoff:

11. Presence and thickness of compaction layer (usually none; describe soil profile features which may be mistaken for compaction on this site):

12. Functional/Structural Groups (list in order of descending dominance by above-ground annual-production or live foliar cover using symbols: >>, >, = to indicate much greater than, greater than, and equal to):

Dominant:

Sub-dominant:

Other:

Additional:

13. Amount of plant mortality and decadence (include which functional groups are expected to show mortality or decadence):

14. Average percent litter cover (%) and depth (in):

15. Expected annual annual-production (this is TOTAL above-ground annual-production, not just forage annual-production):

16. Potential invasive (including noxious) species (native and non-native). List species which BOTH characterize degraded states and have the potential to become a dominant or co-dominant species on the ecological site if their future establishment and growth is not actively controlled by management interventions. Species that become dominant for only one to several years (e.g., short-term response to drought or wildfire) are not invasive plants. Note that unlike other indicators, we are describing what is NOT expected in the reference state for the ecological site:

17. Perennial plant reproductive capability:
