

# Ecological site R043AY510ID

## Poorly Drained Floodplains

### 23-25" PZ Mesic

## Columbia-Colville Valleys

Last updated: 10/15/2020

Accessed: 04/21/2026

### General information

**Provisional.** A provisional ecological site description has undergone quality control and quality assurance review. It contains a working state and transition model and enough information to identify the ecological site.

### MLRA notes

Major Land Resource Area (MLRA): 043A–Northern Rocky Mountains

Major Land Resource Area (MLRA): 043A–Northern Rocky Mountains Description of MLRAs can be found in: United States Department of Agriculture, Natural Resources Conservation Service. 2006. Land Resource Regions and Major Land Resource Areas of the United States, the Caribbean, and the Pacific Basin. U.S. Department of Agriculture Handbook 296.

### LRU notes

Major land resource area (MLRA): 043A-Northern Rocky Mountains Modal LRU – 43A03 - Columbia-Colville Valleys This LRU is composed predominantly of glaciated valleys and foothills. The soils tend to be loamy to sandy Haploxerepts and Haploxerolls with mixed or distinct ash surfaces. Recent alluvium, till, outwash and residuum from granitic or metamorphic rock are the dominant parent materials. Soil climate is a mesic or frigid temperature regime and xeric moisture regime with average annual precipitation around 490 mm (19 inches). Others where occurring: 44A01 Spokane-Rathdrum Outwash Plains 43A04 Selkirk Mountains

### Classification relationships

This ES fits into the National Vegetation Standard's and is most closely related to Washington State's Natural Heritage Program's. (Compare to previous Washington range site: R044XY601WA, WET MEADOW 16-24 PZ)

### Ecological site concept

Ecological Site Concept: This ES is found on poorly to somewhat poorly drained, loamy sites on floodplains. A water table is within 75 cm of the surface at some time during the May-Oct period. They have mesic temperatures and a plant community dominated by sedges and grasses.

Table 1. Dominant plant species

Tree	Not specified
Shrub	Not specified
Herbaceous	(1) <i>Carex</i> (2) <i>Calamagrostis stricta</i>

### Physiographic features

#### Physiographic Features

This ecological site occurs mainly on floodplains and drainageways. Parent materials are mixed alluvium with a mantle of volcanic ash

and loess.

Landscapes: Valleys and scablands

Landforms: depressions, floodplains, drainageways

Elevation:

Total range = 390 to 1040 m

(1,285 to 3,405 feet)

Central tendency = 560 to 750 m

(1,835 to 2,465 feet)

Slope (percent):

Total range = 0 to 3 percent

Central tendency = 1 to 2 percent

Water Table Depth:

0 to 92 cm; median = 31 cm

(0 to 36 inches; median = 12inches)

Flooding:

Frequency: occasional to frequent

Duration: brief to long

Ponding:

Frequency: None to frequent

Duration: None to long

Aspect: NA

**Table 2. Representative physiographic features**

Landforms	(1) Valley > Flood plain (2) Valley > Depression (3) Scabland > Drainageway
Flooding duration	Not specified
Flooding frequency	Not specified
Ponding duration	Not specified
Ponding frequency	Not specified
Elevation	560 – 750 m
Slope	0 %
Water table depth	30 cm
Aspect	Aspect is not a significant factor

**Table 3. Representative physiographic features (actual ranges)**

Flooding duration	Brief (2 to 7 days) to long (7 to 30 days)
Flooding frequency	Occasional to frequent
Ponding duration	Long (7 to 30 days)
Ponding frequency	None to frequent
Elevation	390 – 1,040 m
Slope	0 %
Water table depth	0 – 90 cm

### **Climatic features**

#### Climatic Features

During the spring and summer, a circulation of air around a high-pressure center brings a prevailing westerly and northwesterly flow of comparatively dry, cool and stable air into the region. As the air moves inland, it becomes warmer and drier which results in a dry season beginning in the late spring and reaching a peak in mid-summer. In the fall and winter, a circulation of air around two pressure centers over the ocean brings a prevailing southwesterly and westerly flow of air into the Pacific Northwest. This air from over the ocean is moist and near the temperature of the water. Condensation occurs as the air moves inland over the cooler land and rises along the windward slopes of the mountains or highlands. This results in a wet season beginning in October, reaching a peak in winter, then gradually decreasing in the spring.

The elevation within the LRU varies from approximately 1,000 feet in the lower river valleys to about 4,900 feet on low mountains and foothills. The annual precipitation increases from 9 inches in the valleys to over 38 inches in the higher terrain. Winter season snowfall averages about 43 inches. Both rainfall and snowfall increase with elevation. Snow can be expected after the first of November and to remain on the ground from the first of December until March or April.

In January, the average maximum temperature is near 29° F and the minimum temperature is 14° F. Minimum temperatures from -5° to -20°F are recorded almost every winter and temperatures ranging to -25° F have been recorded. In July, the average maximum temperature is 81° and the minimum temperature 47° F. Maximum temperatures reach 100° F on a few afternoons each summer and temperatures as high as 108° F have been recorded. The average date of the last freezing temperatures can be expected by early-June and after early-September in the warmer areas.

(Compiled from WRCC: Climate of Washington and available station data)

#### Frost-free period (days):

Total range = 100 to 140 days

Central tendency = 115 to 125 days

#### Mean annual precipitation (cm):

Total range = 360 to 855 mm

(14 to 34 inches)

Central tendency = 455 to 615 mm

(18 to 24 inches)

#### MAAT (C)

Total range = 5.8 to 9.9

(42 to 50 F)

Central tendency = 7.6 to 8.7

(46 to 48 F)

Climate stations: none

## Influencing water features

Water Table Depth:

0 to 92 cm; median = 31 cm

(0 to 36 inches; median = 12inches)

Flooding:

Frequency: occasional to frequent

Duration: brief to long

Ponding:

Frequency: None to frequent

Duration: None to long

## Soil features

Representative Soil Features

This ecological site is associated with the Cocolalla, Konner, Peone, and Wethey components. The soils are Aquandic Endoaquepts, Aquandic Endoaquolls, Aquic Xerofluvents, Cumulic Endoaquolls. These soils have developed in mixed alluvium. Most sites have a surface influenced by volcanic ash and loess. Volcanic ash layers are mixed at lower elevations and form distinct, moderately thick layers at higher elevations.

Parent Materials:

Kind: volcanic ash, loess

Origin: mixed

Kind: alluvium

Origin: mixed

Surface Texture: (2mm fraction)

(1) Ashy-Silt Loam

(2) Silty Clay Loam

(3) Loamy Sand

Surface Fragments

Table 4. Representative soil features

Parent material	(1) Volcanic ash (2) Loess (3) Alluvium
Surface texture	(1) Ashy silt loam (2) Silty clay loam (3) Loamy sand
Drainage class	Poorly drained
Permeability class	Moderate
Depth to restrictive layer	Not specified

Surface fragment cover <=3"	Not specified
Surface fragment cover >3"	Not specified
Available water capacity (0-101.6cm)	19.81 cm
Calcium carbonate equivalent (0-152.4cm)	Not specified
Electrical conductivity (0-152.4cm)	Not specified
Sodium adsorption ratio (0-152.4cm)	Not specified
Soil reaction (1:1 water) (0-152.4cm)	6.7

Table 5. Representative soil features (actual values)

Drainage class	Somewhat poorly drained to poorly drained
Permeability class	Slow to very rapid
Depth to restrictive layer	0 cm
Surface fragment cover <=3"	0 – 10 %
Surface fragment cover >3"	0 %
Available water capacity (0-101.6cm)	7.62 – 22.1 cm
Calcium carbonate equivalent (0-152.4cm)	0 %

Electrical conductivity (0-152.4cm)	0 mmhos/cm
Sodium adsorption ratio (0-152.4cm)	0
Soil reaction (1:1 water) (0-152.4cm)	5.6 – 7.3

## Ecological dynamics

## State and transition model

Figure 1. Narrative

Figure 2. STM

## Additional community tables

## References

. (Date accessed). USNVC [United States National Vegetation Classification]. 2019. United States National Vegetation Classification Database, V2.03. Federal Geographic Data Committee, Vegetation Subcommittee, Washington DC.. USNVC: <http://usnvc.org/>.

Gerald, R. 2004. NRCS - Washington; Interim Ecological Site Descriptions for Rangeland.

Kovalchik, B.L. and R.R. Clausnitzer. 2004. Classification and Management of Aquatic, Riparian, and Wetland Sites on the National Forests of Eastern Washington: Series Description. General Technical Report PNW-GTR-593. United States Department of Agriculture, Forest Service, Pacific Northwest Research Station. 1–354.

Rocchio, J.F. and R.C. Crawford. 2015. Ecological systems of Washington State. A guide to identification. Washington Department of Natural Resources.. Natural Heritage Report.. Washington Department of Natural Resources, Natural Heritage Program, Olympia, WA. 1–397.

## Contributors

Stephanie Shoemaker  
Brian Gardner

## Approval

Curtis Talbot, 10/15/2020

## Rangeland health reference sheet

**Interpreting Indicators of Rangeland Health** is a qualitative assessment protocol used to determine ecosystem condition based on benchmark characteristics described in the Reference Sheet. A suite of 17 (or more) indicators are typically considered in an assessment. The ecological site(s) representative of an assessment location must be known prior to applying the protocol and must be verified based on soils and climate. Current plant community cannot be used to identify the ecological site.

Author(s)/participant(s)	
Contact for lead author	
Date	04/21/2026
Approved by	
Approval date	
Composition (Indicators 10 and 12) based on	Annual Production

### Indicators

**1. Number and extent of rills:**

---

**2. Presence of water flow patterns:**

---

**3. Number and height of erosional pedestals or terracettes:**

---

**4. Bare ground from Ecological Site Description or other studies (rock, litter, lichen, moss, plant canopy are not bare ground):**

---

**5. Number of gullies and erosion associated with gullies:**

---

**6. Extent of wind scoured, blowouts and/or depositional areas:**

---

**7. Amount of litter movement (describe size and distance expected to travel):**

---

**8. Soil surface (top few mm) resistance to erosion (stability values are averages - most sites will show a range of values):**

---

**9. Soil surface structure and SOM content (include type of structure and A-horizon color and thickness):**

---

10. **Effect of community phase composition (relative proportion of different functional groups) and spatial distribution on infiltration and runoff:**

---

11. **Presence and thickness of compaction layer (usually none; describe soil profile features which may be mistaken for compaction on this site):**

---

12. **Functional/Structural Groups (list in order of descending dominance by above-ground annual-production or live foliar cover using symbols: >>, >, = to indicate much greater than, greater than, and equal to):**

**Dominant:**

**Sub-dominant:**

**Other:**

**Additional:**

---

13. **Amount of plant mortality and decadence (include which functional groups are expected to show mortality or decadence):**

---

14. **Average percent litter cover (%) and depth ( in):**

---

15. **Expected annual annual-production (this is TOTAL above-ground annual-production, not just forage annual-production):**

---

16. **Potential invasive (including noxious) species (native and non-native). List species which BOTH characterize degraded states and have the potential to become a dominant or co-dominant species on the ecological site if their future establishment and growth is not actively controlled by management interventions. Species that become dominant for only one to several years (e.g., short-term response to drought or wildfire) are not invasive plants. Note that unlike other indicators, we are describing what is NOT expected in the reference state for the ecological site:**

---

17. **Perennial plant reproductive capability:**

---