

Ecological site EX043B23C172

Stony Upland (StU)

Absaroka Subalpine Zone

Last updated: 10/04/2019

Accessed: 08/30/2026

General information

Provisional. A provisional ecological site description has undergone quality control and quality assurance review. It contains a working state and transition model and enough information to identify the ecological site.

MLRA notes

Major Land Resource Area (MLRA): 043B—Central Rocky Mountains

Major Land Resource Unit (MLRA) 43B: Central Rocky Mountains 43B – Central Rocky Mountains – The Central Rocky Mountains extends from northern Montana to southern extent of Wyoming and from Idaho to central Wyoming. The southern extent of 43B is comprised of a combination of metamorphic, igneous, and sedimentary mountains and foothills. Climatic changes across this extent are broad and create several unique breaks in the landscape. Further information regarding MLRAs, refer to: United States Department of Agriculture, Natural Resources Conservation Service. 2006. Land Resource Regions and Major Land Resource Areas of the United States, the Caribbean, and the Pacific Basin. U.S. Department of Agriculture Handbook 296. Available electronically at: http://www.nrcs.usda.gov/wps/portal/nrcs/detail/soils/ref/?cid=nrcs142p2_053624#handbook.

LRU notes

Land Resource Unit (LRU) 43B23C: Absaroka Subalpine Zone Based on the shifts in geology, precipitation patterns and other climatic factors, as well as elevation and vegetation, the Absaroka Range was divided into LRU 23. Further division of this LRU is necessary due to the gradient moving from the foothills to the summit, as well as aspect shifts (north/east face versus south/west face). Subset C is the high elevation zone noted for dense timber interspersed with open parks and longer persisting snowpack (within timberline). Precipitation can range from 18 to 20 plus inches and is more noted for the duration of snow cover and shorter growing season. To verify or identify Subset C (the referenced subset for this ecological site), refer to the Wyoming LRU matrix key contained within the Ecological Site Key. This LRU/Subset occurs on the eastern divide of the Absaroka Range. This LRU starts north of Clark, WY and runs to the Thermopolis, WY area. Once the Absaroka Range merges with the Owl Creek and Wind River Ranges, the climatic patterns and elevational changes shifts the plant community and creates a break in the LRU/Subset. The extent of soils currently correlated to this ecological site does not fit within the digitized boundary. Many of the noted soils are provisional and will be reviewed and corrected in mapping update projects. Other map units are correlated as small inclusions within other MLRA's/LRU's based on elevation, landform, and biological references. Moisture Regime: Typic Ustic Temperature Regime: Cryic Dominant Cover: Rangeland – Sagebrush Steppe (major species is Mountain Big Sagebrush) Representative Value (RV) Effective Precipitation: 20+ inches (508 mm) RV Frost-Free Days: 31-65 days

Classification relationships

Relationship to Other Established Classification Systems: National Vegetation Classification System (NVC): 2 Shrub & Herb Vegetation Class 2.B Temperate & Boreal Grassland & Shrubland Subclass 2.B.2 Temperate Grassland & Shrubland Formation 2.B.2.Na Western North American Grassland & Shrubland Division M048 Central Rocky Mountain Montane-Foothill Grassland & Shrubland Macro-group G273 Central Rocky Mountain Lower Montane, Foothill & Valley Grassland Group Ecoregions (EPA): Level I: 6 North Western Forested Mountains Level II: 6.2 Western Cordillera Level III: 6.2.10 Middle Rockies Level IV: 6.2.17ao – Absaroka Volcanic Subalpine Zone 6.2.17i – Absaroka – Gallatin Volcanic Mountains

Ecological site concept

- Site receives no additional water.
- Slope is 30%
- Soils are:
 - o Textures range from fine sandy loam to clay loam in top 4" (10 cm) of mineral soil surface
 - o All subsurface horizons in the particle size control section have a weighted average of >18% but 35% clay. (The particle size control section is the segment of the profile from either the start of an argillic horizon for 50 cm's or from 25-100 cm's).
 - o Moderately deep to very deep (20-80+ in. (50-200+ cm) o > 5% stone and boulder cover and 20% cobble and gravel cover
 - o Skeletal (? 35% rock fragments) within 20" (50 cm) of mineral soil surface
 - o Non-saline, sodic, or saline-sodic

Associated sites

R043BY162WY	Shallow Loamy High Mountains Shallow Loamy
R043BY170WY	Steep Stony High Mountains Steep Stony

Similar sites

R043BY108WY	Coarse Upland High Mountains Coarse Upland (CU) 20+M has higher production, larger coarse fragments (boulders), and different shrub species.
R043BY170WY	Steep Stony High Mountains Steep Stony (St) 20+M has higher production and different shrub species.

Table 1. Dominant plant species

Tree	Not specified
Shrub	(1) <i>Artemisia tridentata ssp. vaseyana</i>
Herbaceous	(1) <i>Pseudoroegneria spicata</i> (2) <i>Leucopoa kingii</i>

Legacy ID

R043BX772WY

Physiographic features

This site usually occurs on gentle to steep mountain slopes, valley bottoms, and outwash fans.

Table 2. Representative physiographic features

Landforms	(1) Mountain range > Mountain slope (2) Mountain range > Alluvial fan (3) Mountain range > Ridge
Runoff class	Negligible to medium
Elevation	1,980 – 3,660 m

Slope	0 – 30 %
Aspect	Aspect is not a significant factor

Climatic features

Annual precipitation and modeled relative effective annual precipitation range from 18 to 35 inches (457 – 889 mm). The normal precipitation pattern is evenly distributed through the year and averages over 20 inches. Annual snowfall averages 150 to 200 inches annually. Wide fluctuations may occur in yearly precipitation and result in more dry years than those with more than normal precipitation.

Because of the varied topography, the wind will vary considerably for different parts of the area. Prevailing winds are from the southwest, and strong winds are less frequent than over other areas of Wyoming. Occasional storms, however, can bring brief periods of high winds with gusts exceeding 50 mph.

Temperatures show a wide range between summer and winter and between daily maximums and minimums, due to the high elevation and dry air, which permits rapid incoming and outgoing radiation. Cold air outbreaks from Canada in winter move rapidly from northwest to southeast and account for extreme minimum temperatures. Chinook winds may occur in winter and bring rapid rises in temperature. High winds are generally blocked by high mountains but occur in conjunction with thunderstorms, which are common in late summer. Growth of native cool-season plants begins about June 1, but can be as late as July 15, and continues until the beginning of September.

For detailed information visit the Natural Resources Conservation Service National Water and Climate Center at <http://www.wcc.nrcs.usda.gov/>. Climate station representative of this precipitation zone include: “Cooke City 2W” and “Tower Falls”. The following graphs and charts are a collective sample representing the averaged normals and 30-year annual rainfall data for the selected weather stations from 1981 to 2010.

Table 3 Representative climatic features

Frost-free period (characteristic range)	0-0 days
Freeze-free period (characteristic range)	20-50 days
Precipitation total (characteristic range)	460-580 mm
Frost-free period (actual range)	0-0 days
Freeze-free period (actual range)	20-50 days
Precipitation total (actual range)	430-610 mm
Frost-free period (average)	0 days
Freeze-free period (average)	40 days
Precipitation total (average)	530 mm

- (2) TOWER FALLS [USC00489025], Yellowstone National Park, WY

Influencing water features

The characteristics of these upland soils have no influence from ground water (water table below 60 inches (150 cm)) and have minimal influence from surface water/overland flow. There may be isolated features that are affected by snow pack that persists longer than surrounding areas due to position on the landform (shaded/protected pockets).

Soil features

The soils of this site are moderately deep to deep (greater than 20" to bedrock), well-drained, and stony and/or bouldery. Coarse fragments are greater than 35 percent, by volume, within the first 20 inches of soil, usually increasing with depth. Roots penetrate the soil material to at least 20 inches in most places.

Table 4. Representative soil features

Parent material	(1) Alluvium – igneous, metamorphic and sedimentary rock (2) Colluvium
Surface texture	(1) Gravelly, cobbly, stony loam
Family particle size	(1) Fine-loamy (2) Loamy-skeletal
Drainage class	Well drained to somewhat excessively drained
Permeability class	Moderately slow to moderately rapid
Soil depth	50 – 150 cm
Surface fragment cover <=3"	0 – 20 %
Surface fragment cover >3"	0 – 10 %
Available water capacity (0-101.6cm)	5.08 – 10.16 cm
Calcium carbonate equivalent (0-101.6cm)	0 – 20 %
Electrical conductivity (0-101.6cm)	Not specified

Sodium adsorption ratio (0-101.6cm)	Not specified
Soil reaction (1:1 water) (0-101.6cm)	6.6 – 8.4
Subsurface fragment volume <=3" (Depth not specified)	10 – 30 %
Subsurface fragment volume >3" (Depth not specified)	0 – 20 %

Ecological dynamics

As this site deteriorates, species such as rhizomatous wheatgrass, timber oatgrass, and low sagebrush will increase. Bluebunch wheatgrass, spike fescue, and bitterbrush will decrease in frequency and production. This site has relatively low productivity potential, and is not well suited to grazing improvement practices unless treated as part of a larger unit containing more productive areas.

The Historic Climax Plant Community (description follows the plant community diagram) has been determined by study of rangeland relic areas, or areas protected from excessive disturbance. Trends in plant communities going from heavily grazed areas to lightly grazed areas, seasonal use pastures, and historical accounts have also been used.

The following is a State and Transition Model Diagram that illustrates the common plant communities (states) that can occur on the site and the transitions between these communities. The ecological processes will be discussed in more detail in the plant community narratives following the diagram.

State and transition model

Additional community tables

Table 5. Community 1.1 plant community composition

Group	Common Name	Symbol	Scientific Name	Annual Production ()	Foliar Cover (%)
Grass/Grasslike					
1				280-392	
	bluebunch wheatgrass	PSSP6	<i>Pseudoroegneria spicata</i>	280-392	–
2				168-280	
	spike fescue	LEK12	<i>Leucopoa kingii</i>	168-280	–
3				56-112	
	Idaho fescue	FEID	<i>Festuca idahoensis</i>	56-112	–
4				112-224	
	Grass, perennial	2GP	<i>Grass, perennial</i>	0-56	–
	Letterman's needlegrass	ACLE9	<i>Achnatherum lettermanii</i>	0-56	–
	Columbia needlegrass	ACNE9	<i>Achnatherum nelsonii</i>	0-56	–
	western needlegrass	ACOC3	<i>Achnatherum occidentale</i>	0-56	–
	bentgrass	AGROS2	<i>Agrostis</i>	0-56	–
	timber oatgrass	DAIN	<i>Danthonia intermedia</i>	0-56	–

	onespike danthonia	DAUN	<i>Danthonia unispicata</i>	0-56	–
	squirreltail	ELELE	<i>Elymus elymoides</i> ssp. <i>elymoides</i>	0-56	–
	thickspike wheatgrass	ELLAL	<i>Elymus lanceolatus</i> ssp. <i>lanceolatus</i>	0-56	–
	slender wheatgrass	ELTR7	<i>Elymus trachycaulus</i>	0-56	–
	prairie Junegrass	KOMA	<i>Koeleria macrantha</i>	0-56	–
	oniongrass	MEBU	<i>Melica bulbosa</i>	0-56	–
	mountain muhly	MUMO	<i>Muhlenbergia montana</i>	0-56	–
	muttongrass	POFE	<i>Poa fendleriana</i>	0-56	–
	spike trisetum	TRSP2	<i>Trisetum spicatum</i>	0-56	–
Forb					
5				56-112	
	Forb, perennial	2FP	<i>Forb, perennial</i>	0-56	–
	common yarrow	ACMI2	<i>Achillea millefolium</i>	0-56	–
	agoseris	AGOSE	<i>Agoseris</i>	0-56	–
	pussytoes	ANTEN	<i>Antennaria</i>	0-56	–
	sandwort	ARENA	<i>Arenaria</i>	0-56	–
	milkvetch	ASTRA	<i>Astragalus</i>	0-56	–
	balsamroot	BALSA	<i>Balsamorhiza</i>	0-56	–
	Indian paintbrush	CASTI2	<i>Castilleja</i>	0-56	–
	hawksbeard	CREPI	<i>Crepis</i>	0-56	–
	fleabane	ERIGE2	<i>Erigeron</i>	0-56	–
	buckwheat	ERIOG	<i>Eriogonum</i>	0-56	–
	stoneseed	LITHO3	<i>Lithospermum</i>	0-56	–
	desertparsley	LOMAT	<i>Lomatium</i>	0-56	–
	lupine	LUPIN	<i>Lupinus</i>	0-56	–
	creeping barberry	MARE11	<i>Mahonia repens</i>	0-56	–
	bluebells	MERTE	<i>Mertensia</i>	0-56	–
	owl's-clover	ORTHO	<i>Orthocarpus</i>	0-56	–
	ragwort	PACKE	<i>Packera</i>	0-56	–
	phlox	PHLOX	<i>Phlox</i>	0-56	–
	stonecrop	SEDUM	<i>Sedum</i>	0-56	–
	ragwort	SENEC	<i>Senecio</i>	0-56	–
	aster	SYMPH4	<i>Symphyotrichum</i>	0-56	–
	clover	TRIFO	<i>Trifolium</i>	0-56	–
	American vetch	VIAM	<i>Vicia americana</i>	0-56	–
Shrub/Vine					
6				56-112	
	antelope bitterbrush	PUTR2	<i>Purshia tridentata</i>	56-112	–
7				56-112	
	little sagebrush	ARAR8	<i>Artemisia arbuscula</i>	56-112	–
8				56-168	
	Shrub, deciduous	2SD	<i>Shrub, deciduous</i>	0-56	–
	Shrub, evergreen	2SE	<i>Shrub, evergreen</i>	0-56	–
	Tree, deciduous	2TD	<i>Tree, deciduous</i>	0-56	–
	Tree, evergreen	2TE	<i>Tree, evergreen</i>	0-56	–
	threetip sagebrush	ARTR4	<i>Artemisia tripartita</i>	0-56	–

	mountain big sagebrush	ARTRV	<i>Artemisia tridentata ssp. vaseyana</i>	0-56	–
	snowberry	SYMPH	<i>Symphoricarpos</i>	0-56	–

Table 6. Community 2.1 plant community composition

Group	Common Name	Symbol	Scientific Name	Annual Production ()	Foliar Cover (%)
-------	-------------	--------	-----------------	----------------------	------------------

Table 7. Community 3.1 plant community composition

Group	Common Name	Symbol	Scientific Name	Annual Production ()	Foliar Cover (%)
-------	-------------	--------	-----------------	----------------------	------------------

Animal community

Animal Community – Wildlife Interpretations Bluebunch Wheatgrass/Bitterbrush Plant Community (HCPC): When blown clear, this plant community provides limited winter forage for large grazers when snow depth prevents foraging on other sites. Otherwise, it is mostly used by wildlife in transit to other habitats. Low Sage/Rhizomatous Wheatgrass Plant Community: This plant community may be useful for the same wildlife that would use the Historic Climax Plant Community. However, the plant community composition is less diverse, and thus, less apt to meet the seasonal needs of these animals. Three-tip Sage Plant Community: This plant community exhibits a low level of plant species diversity. In most cases, it is not a desirable plant community to select as a wildlife habitat management objective. Animal Community – Grazing Interpretations The following table lists suggested stocking rates for cattle under continuous season-long grazing under normal growing conditions. These are conservative estimates that should be used only as guidelines in the initial stages of the conservation planning process. Often, the current plant composition does not entirely match any particular plant community (as described in this ecological site description). Because of this, a field visit is recommended, in all cases, to document plant composition and production. More precise carrying capacity estimates should eventually be calculated using this information along with animal preference data, particularly when grazers other than cattle are involved. Under more intensive grazing management, improved harvest efficiencies can result in an increased carrying capacity. If distribution problems occur, stocking rates must be reduced to maintain plant health and vigor. Plant Community Production Carrying Capacity* (lb./ac) (AUM/ac) Bluebunch Wheatgrass/Bitterbrush (HCPC) 750-1200 0.3 Low Sage/Rhizomatous Wheatgrass 500-1000 0.2 Three-tip Sage 200-800 0.1 * - Continuous, season-long grazing by cattle under average growing conditions. Grazing by domestic livestock is one of the major income-producing industries in the area. Rangeland in this area may provide yearlong forage for cattle, sheep, or horses. During the dormant period, the forage for livestock use needs to be supplemented with protein because the quality does not meet minimum livestock requirements.

Hydrological functions

Water is the principal factor limiting forage production on this site. This site is dominated by soils in hydrologic group B and C, with localized areas in hydrologic group D. Infiltration ranges from moderately slow to moderate. Runoff potential for this site varies from low to moderate depending on soil hydrologic group and ground cover (refer to Part 630, NRCS National Engineering Handbook for detailed hydrology information). Rills and gullies should not typically be present. Water flow patterns should be barely distinguishable if at all present. Pedestals are only slightly present in association with bunchgrasses. Litter typically falls in place, and signs of movement are not common. Chemical and physical crusts are rare to non-existent. Cryptogamic crusts are present, but only cover 1-2% of the soil surface.

Recreational uses

This site provides hunting opportunities for upland game species.

Wood products

No appreciable wood products are present on the site.

Inventory data references

Information presented here has been derived from NRCS clipping data and other inventory data. Field observations from range trained personnel were also used. Those involved in developing this site include: Bill Christensen, Range Management Specialist, NRCS; Karen Clause, Range Management Specialist, NRCS; and Everett Bainter, Range Management Specialist, NRCS. Other sources used as references include USDA NRCS Water and Climate Center, USDA NRCS National Range and Pasture Handbook, USDI and USDA Interpreting Indicators of Rangeland Health Version 3, and USDA NRCS Soil Surveys from various counties. Information presented here has been derived from NRCS inventory data, Field observations from range trained personnel, and the existing range site descriptions. Those involved in developing the Loamy range site include: Karen Clause, Range Management Specialist, NRCS and Everett Bainter, Range Management Specialist. Those involved in the development of the new concept for Loamy and Loamy Calcareous Ecological site include: Ray Gullion, Area Range Management Specialist, NRCS; Jim Wolf, Resource Manager, USDI-BLM; Jack Mononi, Range Management Specialist, USDI-BLM; Daniel Wood, MLRA Soil Survey Leader, NRCS; Jane Karinen, Soil Data Quality Specialist, NRCS; and Marji Patz, Ecological Site Specialist, NRCS. Inventory Data References: Ocular field estimations observed by trained personnel were completed at each site. Then sites were selected where a 100-foot tape was stretched, and the following sample procedures were

completed by inventory staff. For full sampling protocol and guidelines with forms please refer to the Wyoming ESI Operating Procedures, compiled in 2012 for the Powell and Rock Springs Soil Survey Office, USDA-NRCS. • Double Sampling Production Data (4.8 square foot hoop used to estimate 10 points, clipped a minimum of 2 of these estimated points, with two 21-foot X 21-foot square extended shrub plots). • Line Point Intercept (over story and understory captured with soil cover). Height of herbaceous and woody cover is collected every three feet along established transect.) • Continuous Line Intercept (Woody Canopy Cover, with minimum gap of 0.2 of a foot for all woody species and succulents. Intercept height collected at each measurement.), • Gap Intercept (Basal Gap measured with a minimum gap requirement of 0.7 foot.), • Sample Point (10 – 1-meter square point photographs taken at set distances on transect. Red using the sample point computer program established by the High Plains Agricultural Research Center, WY). • Soil Stability (Slake Test – surface and subsurface samples collected and processed according to the soil stability guidelines provided by the Jornada Research Center, NM.)

Other references

Baker, William L. 2006. Fire and Restoration of Sagebrush Ecosystems. *Wildlife Society Bulletin* 34(1): 177-185.

Bestelmeyer, B., and J. R. Brown. 2005. State-and-transition models 101: a fresh look at vegetation change. *The Quivira Coalition Newsletter*, Vol. 7, No. 3.

Bestelmeyer, B., J. R. Brown, K. M. Havstad, B. Alexander, G. Chavez, J. E. Herrick. 2003. Development and use of state and transition models for rangelands. *Journal of Range Management* 56(2):114-126.

Bestelmeyer, B., J. E. Herrick, J. R. Brown, D. A. Trujillo, and K. M. Havstad. 2004. Land management in the American Southwest: a state-and-transition approach to ecosystem complexity. *Environmental Management* 34(1):38-51.

Herrick, J. E., J. W. Van Zee, K. M. Havstad, L. M. Burkett, and W. G. Whitford. 2005. Monitoring manual for grassland, shrubland and savanna Ecosystems. Volume I Quick Start. USDA - ARS Jornada Experimental Range, Las Cruces, New Mexico.

Herrick, J. E., J. W. Van Zee, K. M. Havstad, L. M. Burkett, and W. G. Whitford. 2005. Monitoring manual for grassland, shrubland and savanna Ecosystems. Volume II: Design, supplementary methods and interpretation. USDA - ARS Jornada Experimental Range, Las Cruces, New Mexico.

NRCS. 2014. (electronic) National Water and Climate Center. Available online at <http://www.wcc.nrcs.usda.gov/>

NRCS. 2014. (electronic) Field Office Technical Guide. Available online at http://efotg.nrcs.usda.gov/efotg_locator.aspx?map=WY NRCS. 2009. Plant Guide: Cheatgrass. Prepared by Skinner et al., National Plant Data Center.

Pellant, M., P. Shaver, D. A. Pyke, and J. E. Herrick. 2005. Interpreting indicators of rangeland health. Version 4. Technical Reference 1734-6.

USDI-BLM. Ricketts, M. J., R. S. Noggles, and B. Landgraf-Gibbons. 2004. Pryor Mountain Wild Horse Range Survey and Assessment. USDA-Natural Resources Conservation Service.

Schoeneberger, P. J., D. A. Wysocki, E. C. Benham, and Soil Survey Staff. 2012. Field book for describing and sampling soils, Version 3.0. Natural Resources Conservation Service, National Soil Survey Center, Lincoln, NE. (<http://soils.usda.gov/technical/fieldbook/>)

Stringham, T. K. and W. C. Krueger. 2001. States, transitions, and thresholds: Further refinement for rangeland applications. Agricultural Experiment Station, Oregon State University. Special Report 1024.

Stringham, T. K., W. C. Kreuger, and P. L. Shaver. 2003. State and transition modeling: an ecological process approach. *Journal of Range Management* 56(2):106-113.

United States Department of Agriculture. Soil Survey Division Staff. 1993. Soil Survey Manual, United States Department of Agriculture Handbook No. 18, Chapter 3: Examination and Description of Soils. Pg.192-196.

USDA, NRCS. 1997. National Range and Pasture Handbook. (<http://www.glti.nrcs.usda.gov/technical/publications/nrph.html>)

Trlica, M. J. 1999. Grass growth and response to grazing. Colorado State University. Cooperative Extension. Range. Natural Resource Series. No. 6.108.

U.S. Department of Agriculture, Natural Resources Conservation Service (USDA/NRCS). 2007. The PLANTS Database (<http://plants.usda.gov>). National Plant Data Center, Baton Rouge, LA 70874-4490 USA.

U.S. Department of Agriculture, Natural Resources Conservation Service (USDA/NRCS), Soil Survey Staff. 2010. Keys to Soil Taxonomy, Eleventh Edition, 2010.

Contributors

K. Clause

Approval

Scott Woodall, 10/04/2019

Rangeland health reference sheet

Interpreting Indicators of Rangeland Health is a qualitative assessment protocol used to determine ecosystem condition based on benchmark characteristics described in the Reference Sheet. A suite of 17 (or more) indicators are typically considered in an assessment. The ecological site(s) representative of an assessment location must be known prior to applying the protocol and must be verified based on soils and climate. Current plant community cannot be used to identify the ecological site.

Author(s)/participant(s)	K. Clause, E. Bainter
Contact for lead author	karen.clause@wy.usda.gov or 307-367-2257
Date	03/16/2007
Approved by	
Approval date	
Composition (Indicators 10 and 12) based on	Annual Production

Indicators

1. **Number and extent of rills:** Rare to nonexistent. Where present, short and widely spaced.

2. **Presence of water flow patterns:** Barely observable.

3. **Number and height of erosional pedestals or terracettes:** Rare to nonexistent.

4. **Bare ground from Ecological Site Description or other studies (rock, litter, lichen, moss, plant canopy are not bare ground):**
Bare ground can range from 0-25%.

5. Number of gullies and erosion associated with gullies: Active gullies should not be present.

6. Extent of wind scoured, blowouts and/or depositional areas: Rare to nonexistent.

7. Amount of litter movement (describe size and distance expected to travel): Herbaceous and large woody litter not expected to move.

8. Soil surface (top few mm) resistance to erosion (stability values are averages - most sites will show a range of values): Soil Stability Index ratings range from 2 (interspaces) to 6 (under plant canopy), but average values should be 3.5 or greater.

9. Soil surface structure and SOM content (include type of structure and A-horizon color and thickness): Soil data is limited for this site. Soil OM of 4-16% is expected.

10. Effect of community phase composition (relative proportion of different functional groups) and spatial distribution on infiltration and runoff: Plant community consists of 55-80% grasses, 10% forbs, and 10-35% shrubs. Evenly distributed plant canopy (50-75%) and litter plus moderate infiltration rates result in minimal runoff. Basal cover is typically less than 15% and marginally affects runoff on this site. Surface rock fragments of 20-50% provide stability to the site, but reduce infiltration.

11. Presence and thickness of compaction layer (usually none; describe soil profile features which may be mistaken for compaction on this site): None.

12. Functional/Structural Groups (list in order of descending dominance by above-ground annual-production or live foliar cover using symbols: >>, >, = to indicate much greater than, greater than, and equal to):

Dominant: Mid-size, cool season bunchgrasses

Sub-dominant: perennial shrubs perennial forbs

Other: cool season rhizomatous grasses=short cool season bunchgrasses

Additional:

13. Amount of plant mortality and decadence (include which functional groups are expected to show mortality or decadence): Minimal decadence, typically associated with shrub component.

14. Average percent litter cover (%) and depth (in): Litter ranges from 15-20% of total canopy measurement with total litter (including beneath the plant canopy) from 40-60% expected. Herbaceous litter depth typically ranges from 3-10mm. Woody litter can be up to a couple inches (4-6 cm).

15. **Expected annual annual-production (this is TOTAL above-ground annual-production, not just forage annual-production):**
English: 750-1200 lb/ac (1000 lb/ac average); Metric 840-1344 kg/ha (1120 kg/ha average).
-

16. **Potential invasive (including noxious) species (native and non-native).** List species which BOTH characterize degraded states and have the potential to become a dominant or co-dominant species on the ecological site if their future establishment and growth is not actively controlled by management interventions. Species that become dominant for only one to several years (e.g., short-term response to drought or wildfire) are not invasive plants. Note that unlike other indicators, we are describing what is NOT expected in the reference state for the ecological site: Bare ground greater than 40% is the most common indicator of a threshold being crossed. Rabbitbrush, Sandberg bluegrass, buckwheat, yarrow, and phlox are common increasers. Annual weeds such as cheatgrass and mustards are common invasive species in disturbed sites.
-

17. **Perennial plant reproductive capability:** All species are capable of reproducing, except in extreme drought years.
-