

# Ecological site F043CY601OR

## Cold Wet Conifer Mountains and Plateaus (ABLA/VASC-VAME)

Last updated: 4/04/2025  
Accessed: 08/16/2026

---

### General information

**Provisional.** A provisional ecological site description has undergone quality control and quality assurance review. It contains a working state and transition model and enough information to identify the ecological site.

### MLRA notes

Major Land Resource Area (MLRA): 043C–Blue and Seven Devils Mountains

This MLRA covers the Blue and Seven Devils Mountains of Oregon, Washington and Idaho. The area is characterized by thrust and block-faulted mountains and deep canyons composed of sedimentary, metasedimentary, and volcanic rocks. Elevations range from 1,300 to 9,800 feet (395 to 2,990 meters). The climate is characterized by cold, wet winters and cool, dry summers. Annual precipitation, mostly in the form of snow, averages 12 to 43 inches (305 to 1,090 millimeters) yet ranges as high as 82 inches (2,085 millimeters) at upper elevations. Soil temperature regimes are predominately Frigid to Cryic and soil moisture regimes are predominately Xeric to Udic. Mollisols and Andisols are the dominant soil orders. Ecologically, forests dominate but shrub and grass communities may occur on south aspects and lower elevations as well as in alpine meadow environments. Forest composition follows moisture, temperature and elevational gradients and typically ranges from ponderosa pine and Douglas-fir plant associations at lower elevations, grand fir at middle elevations and subalpine fir and Engelmann spruce at upper elevations. Historical fire regimes associated with these forest types range from frequent surface fires in ponderosa pine - Douglas Fir forest types to mixed and stand replacing fire regimes in grand fir and subalpine fir types. A large percentage of the MLRA is federally owned and managed by the U.S. Forest Service for multiple uses.

### Classification relationships

Plant Assoc. Of Blue and Ochoco Mountains (R6 E TP-036-92) Subalpine fir/grouse huckleberry (blueberry) - CES411 (modal) Lodgepole pine(subalpine fir)/big huckleberry - CLS514 Lodgepole pine(subalpine fir)/big huckleberry/pinegrass - CLS516 Lodgepole pine(subalpine fir)/elk sedge - CLG322 Lodgepole pine(subalpine fir)/grouse huckleberry - CLS418 Lodgepole pine(subalpine fir)/western needlegrass - CLG11 Subalpine fir/big huckleberry - CES311 Subalpine fir/elk sedge - CAG111 Subalpine fir/heartleaf arnica - CEF412 Subalpine fir/queencup beadlily - CES314 Subalpine fir/twinflower - CES414 Subalpine fir/western (false) bugbane - CEF331 Plant Assoc. Of Wallowa-Snake Province (R6 E 255-86) Subalpine fir/grouse huckleberry/Jacob's ladder - CES415 (modal) Engelmann spruce/queencup beadlily - CEM222 Lodgepole pine (subalpine fir)/blue(big) huckleberry - CLF211 Lodgepole pine(subalpine fir)/big huckleberry - CLS515 Lodgepole pine(subalpine fir)/grouse huckleberry/Jacob's ladder - CLS415 Subalpine fir/big huckleberry - CES315 Subalpine fir/pinegrass - CEG312 Subalpine fir/queencup beadlily - CES131 Subalpine fir/skunkleaf polemonium - CEF411 Subalpine fir/twinflower - CEF221 Subalpine fir/twistedstalk - CEF311 U.S. National Vegetation Classification (NVC) Standard Rocky Mountain Subalpine Dry-Mesic Spruce-Fir Forest & Woodland - Group-219 Rocky Mountain Mesic-Wet Subalpine Fir-Engelmann Spruce Forest - Alliance-3614 Forest Service Ecological Sub-region Blue Mountains - M332 LANDFIRE BpS model Rocky Mountain Subalpine Mesic-Wet Spruce-Fir Forest and Woodland - 0910560

### Ecological site concept

This Ecological Site occurs mainly on forested backslopes and summits of higher elevation mountain slopes, and plateaus. As one of the highest forest types in MLRA 43C, this site is often adjacent to subalpine meadows and parklands. In higher elevation mountain ranges such as the Elkhorns, Strawberries or Wallows, forests and subalpine woodlands dominated by whitebark pine may occur above and adjacent to this site. This site has a cryic temperature regime and udic moisture regime. The climate is characterized by cold winters during which deep snowpacks accumulate, and cool, dry summers. Parent materials are derived from basalt or other igneous extrusive geologies (andesite, dacite, ignimbrites, etc.) with a thick mantle of volcanic ash and loess. Other geologies such as granitics or metavolcanics may also be present. These are well drained soils with adequate available water capacity. Subalpine fir generally represents late seral stages however Engelmann spruce may be codominant, and other species such as grand fir, western larch, Douglas-fir, whitebark pine and lodgepole pine may be present. Major understory species may range from grouse whortleberry, thinleaf huckleberry, and twinflower, to elk sedge and Columbia brome. This is a provisional ecological site that groups characteristics at a broad scale with little to no field verification and is subject to extensive review and revision before final approval. All data herein was developed using existing information and literature and should be considered provisional and contingent upon field validation prior to use in conservation planning.

## Associated sites

|                    |                                                                                                                                                                            |
|--------------------|----------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| <b>R043CY801OR</b> | <b>Cold Dry Subalpine Grasslands (FEVI)</b><br><br>Shallower soils, often on exposed, rocky positions with lower moisture holding capacity                                 |
| <b>F043CY503OR</b> | <b>Mountain Riparian Forest (PIEN/ALIN)</b><br><br>Occupying adjacent moderate to high-energy riparian areas                                                               |
| <b>F043CY603OR</b> | <b>Cool Wet Conifer Mountains and Plateaus (ABGR/VAME/LIBO)</b><br><br>Occupying adjacent soils with warmer temperatures due to aspect or landscape position               |
| <b>F043CY605OR</b> | <b>Cool Moist Conifer Mountains and Plateaus (PSME-PIPO/CARU)</b><br><br>Occupying adjacent soils with warmer and drier micro climates due to aspect or landscape position |

## Similar sites

|                    |                                                                                                                                            |
|--------------------|--------------------------------------------------------------------------------------------------------------------------------------------|
| <b>F043CY603OR</b> | <b>Cool Wet Conifer Mountains and Plateaus (ABGR/VAME/LIBO)</b><br><br>Frigid soil temperature regime, subalpine fir uncommon or absent    |
| <b>R043CY801OR</b> | <b>Cold Dry Subalpine Grasslands (FEVI)</b><br><br>Shallower soils, often on exposed, rocky positions with lower moisture holding capacity |

**Table 1. Dominant plant species**

|            |                                                                     |
|------------|---------------------------------------------------------------------|
| Tree       | (1) <i>Abies lasiocarpa</i>                                         |
| Shrub      | (1) <i>Vaccinium scoparium</i><br>(2) <i>Vaccinium membranaceum</i> |
| Herbaceous | Not specified                                                       |

## Physiographic features

As one of the highest elevation forest types in MLRA 43c, this site occurs primarily on upper elevation backslopes of mountain slopes and plateaus. Sites that occur on summit positions may be more exposed to desiccation by wind. Slopes are most commonly 7.5 - 45% but may be as steep as 65%. Slope profiles range from concave, to convex to linear across and down slope. Elevations are typically 5,750 to 6,550 (1,750 to 2,000 m) but may range from 4,900 to 9,000 ft (1,500 to 2,750 m). This site does not experience flooding or ponding and no water table is present within the upper two meters of soil.

**Table 2. Representative physiographic features**

|           |                                                           |
|-----------|-----------------------------------------------------------|
| Landforms | (1) Mountains > Mountain slope<br>(2) Mountains > Plateau |
|-----------|-----------------------------------------------------------|

|                    |                            |
|--------------------|----------------------------|
| Flooding frequency | None                       |
| Ponding frequency  | None                       |
| Elevation          | 1,750 – 2,000 m            |
| Slope              | 10 – 50 %                  |
| Ponding depth      | 0 cm                       |
| Water table depth  | 250 cm                     |
| Aspect             | W, NW, N, NE, E, SE, S, SW |

**Table 3. Representative physiographic features (actual ranges)**

|                    |                 |
|--------------------|-----------------|
| Flooding frequency | Not specified   |
| Ponding frequency  | Not specified   |
| Elevation          | 1,490 – 2,740 m |
| Slope              | 0 – 70 %        |
| Ponding depth      | Not specified   |
| Water table depth  | Not specified   |

### Climatic features

The climate of this site is characterized by high elevation, intermountain patterns. Winters are cold and wet with deep accumulations of snow persisting into late spring or early summer. Summers are cool and dry with occasional convective thunderstorms bringing measurable precipitation. Compared to continental systems influencing interior Rocky Mountain landscapes, these storms are less frequent and therefore summer precipitation is lower. This site has a Cryic soil temperature regime and a Udic soil moisture regime typically experiencing fewer than 45 dry days per year. Mean annual precipitation is typically 30 - 52 in. (1,045 – 1,415 mm) but ranges from 25 – 65 in. (710 – 1,590 mm). Mean annual temperatures are typically 36 – 39°F (2 - 4 °C) but may range from 34 – 41°F (1 - 5°C). Frost free days average 15 to 70 per year. Climate graphs are populated from the closest available weather stations and are included to represent general trends rather than representative values.

**Table 4 Representative climatic features**

|                                          |            |
|------------------------------------------|------------|
| Frost-free period (characteristic range) | 20-70 days |
|------------------------------------------|------------|

|                                            |              |
|--------------------------------------------|--------------|
| Freeze-free period (characteristic range)  |              |
| Precipitation total (characteristic range) | 760-1,320 mm |
| Frost-free period (average)                | 40 days      |
| Freeze-free period (average)               |              |
| Precipitation total (average)              | 970 mm       |

- (1) MEACHAM [USW00024152], Pendleton, OR
- (2) AUSTIN 3 S [USC00350356], Prairie City, OR
- (3) SENECA [USC00357675], Seneca, OR

### Influencing water features

This site is not influenced by water from a wetland or stream.

### Soil features

Soils that typify this site concept are moderately deep to very deep and well drained. Parent materials are diverse and typically composed of materials ranging from colluvium, residuum and glacial till derived from basalt, andesite, granite and argillite beneath thick mantles of volcanic ash. Surface textures are commonly silt loams and may be significantly ashy or stony. The family particle size is typically ashy over loamy skeletal or loamy skeletal but may also be ashy over loamy. See Troutmeadows, Mountemily, and Mudlakebasin for modal series concepts.

**Table 5. Representative soil features**

|                      |                                                                                                                                                         |
|----------------------|---------------------------------------------------------------------------------------------------------------------------------------------------------|
| Parent material      | (1) Colluvium – volcanic rock<br>(2) Residuum – volcanic rock<br>(3) Residuum – argillite<br>(4) Residuum – granite<br>(5) Volcanic ash – volcanic rock |
| Surface texture      | (1) Gravelly silt loam<br>(2) Ashy silt loam<br>(3) Stony silt loam                                                                                     |
| Family particle size | (1) Ashy over loamy-skeletal<br>(2) Ashy over loamy<br>(3) Loamy-skeletal                                                                               |

|                                                   |                                     |
|---------------------------------------------------|-------------------------------------|
| Drainage class                                    | Well drained                        |
| Permeability class                                | Moderately slow to moderately rapid |
| Depth to restrictive layer                        | 50 – 200 cm                         |
| Soil depth                                        | 50 – 200 cm                         |
| Surface fragment cover <=3"                       | 0 – 50 %                            |
| Surface fragment cover >3"                        | 0 – 50 %                            |
| Available water capacity<br>(0-101.6cm)           | 11.68 – 26.42 cm                    |
| Soil reaction (1:1 water)<br>(0-101.6cm)          | 5.1 – 7.3                           |
| Subsurface fragment volume <=3"<br>(10.2-152.4cm) | 10 – 30 %                           |
| Subsurface fragment volume >3"<br>(10.2-152.4cm)  | 10 – 30 %                           |

### Ecological dynamics

Occurring in the high elevations of the Blue and Wallowa mountains, the reference plant community of this site at maturity is represented by a subalpine fir dominated forest. Engelman spruce (*Picea Englemanii*) may be subdominant or codominant at maturity, with other conifers such as lodgepole pine (*Pinus contorta*), western larch (*Larix occidentalis*) and Douglas-fir (*Pseudotsuga menziesii*) more common in younger forests. Douglas-fir and grand fir (*Abies grandis*) may be more common on warmer microsites within this concept, while whitebark pine (*Pinus albicaulis*) may be present toward the coldest, upper elevation limits of the site. The modal concept for this site is the USFS plant association, subalpine fir/grouse huckleberry (ABLA/VASC) yet the breadth of the concept also includes plant associations such as subalpine fir/grouse huckleberry/Jacob's ladder, subalpine fir/big huckleberry, subalpine fir/twinflower and subalpine fir/pinegrass. The understory of mid development and mature forests is often characterized by shrubs and herbaceous species such as grouse whortleberry (*Vaccinium scoparium*), thinleaf huckleberry (*Vaccinium membranaceum*), prince's pine (*Chimaphila umbellata*), twinflower (*Linnea borealis*), heartleaf arnica (*Arnica cordifolia*), pinegrass (*Calamagrostis rubescens*), elk sedge (*Carex geyeri*), Ross' sedge (*Carex rossii*), Columbia brome (*Bromus vulgaris*) and Oregon boxwood (*Pachistima myrsinites*).

These forest sites were historically subject to relatively infrequent, stand replacing and mixed severity wildfires approximating Landfire fire regime 3,35-100+ year frequency, mixed severity (Landfire 2007). Both mixed and stand replacement fires occurred at roughly equal frequencies, with an average estimate of 125 for any fire event. Surface fires were largely absent from this forest type. Fire frequency is influenced by the cold, wet climate and intensity is influenced by the low fire resistance of the dominant tree species. The two most common conifers on this site at maturity, subalpine fir and Engelman spruce, share very low resistance to fire with high mortality occurring even with low intensity fires. This is the result of shallow rooting, thin bark and low hanging branches that characterize both species. Following fire, lodgepole pine will often dominate young tree regeneration due to its prolific seed production and establishment on newly burned sites. Engelmann spruce also produces high amounts of viable seed and may be common following disturbance,

whereas subalpine fir often regenerates later in the subcanopy. The superior shade tolerance of Engelmann spruce and subalpine fir in comparison to lodgepole pine, will help to allow them to dominate the canopy overtime in the absence of fire. This is common within the reference conditions, yet sites experiencing more frequent fire will tend toward forest compositions with higher proportions of lodgepole pine, sometimes persisting for 100s of years.

Western larch is also common following disturbance and given adequate light will grow quickly. With thick bark at maturity, western larch is highly resistant to fire and large individuals may persist in mature stands. Douglas-fir is common on warmer microsites such as southern aspects or the lower elevations of this site. Like western larch, older individuals are highly fire resistant and may survive in mature stands. Grand fir may be present on lower elevation sites and is especially common during early regeneration and young forest phases. As a shade tolerant conifer, it may also regenerate in mature stands where sites are favorable.

Many of the common understory shrubs and graminoids species of this site are adapted to resprout following fire. Both big huckleberry and grouse huckleberry will resprout, with big huckleberry sometimes following after grouse huckleberry. Pinegrass and elk sedge will resprout from rhizomes in addition to some forbs such as heartleaf arnica which may increase following fire. Understory composition will change overtime with canopy closure and subsequent alteration of surface microclimates. Relative cover of grouse huckleberry will decrease with canopy closure as big huckleberry increases somewhat.

In addition to fire disturbance, this ecologic site is also impacted by periodic windthrow, and by significant insect and disease disturbances that affect major tree species. Of particular interest are spruce beetles which can kill both mature subalpine fir and Engelman spruce trees, and mountain pine beetle which targets lodgepole pine, especially mature trees. The interactions of the disturbances are described in the Landfire BPS model 0710550, Rocky Mountain Subalpine Dry-Mesic Spruce-Fir Forest and Woodland. The state and transition model below is largely based on the dynamics described in this BPS.

More protected sites with less droughty or windswept aspects may include dynamics that include those described in the Landfire BPS model 0710560 Rocky Mountain Subalpine Mesic-Wet Spruce-Fir Forest and Woodland, which has a somewhat longer replacement fire return interval and includes root diseases as a disturbance agent. Note that Balsam wooly adelgid is not present in this historical reference model since it is an introduced insect, despite currently contributing to high mortality of true firs.

Emerging evidence is suggesting that climate change is leading to hotter and drier conditions in western forests that will increase fire frequency and extent and lengthen fire seasons (Halofsky et al. 2020). While these changing conditions are likely to impact lower elevation forests first, catastrophic fire may carry through to higher elevation forests. However, when combined with the interacting impacts of drought and insect outbreaks, it is possible that this ecological system will experience unpredictable ecosystem shifts and additional alternative states. Further, while the effects of climate change may contribute to transitions of this forest type to species adapted to warmer and drier climates, further research is required to confidently model these impacts. For this reason, this scenario has not yet been included as a potential alternative state.

State and transition model

Additional community tables

Table 6. Community 1.1 plant community composition

| Group | Common Name | Symbol | Scientific Name | Annual Production () | Foliar Cover (%) |
|-------|-------------|--------|-----------------|----------------------|------------------|
|-------|-------------|--------|-----------------|----------------------|------------------|

Table 7. Community 1.2 plant community composition

| Group | Common Name | Symbol | Scientific Name | Annual Production () | Foliar Cover (%) |
|-------|-------------|--------|-----------------|----------------------|------------------|
|-------|-------------|--------|-----------------|----------------------|------------------|

Table 8. Community 1.3 plant community composition

| Group | Common Name | Symbol | Scientific Name | Annual Production () | Foliar Cover (%) |
|-------|-------------|--------|-----------------|----------------------|------------------|
|-------|-------------|--------|-----------------|----------------------|------------------|

Table 9. Community 1.4 plant community composition

| Group | Common Name | Symbol | Scientific Name | Annual Production () | Foliar Cover (%) |
|-------|-------------|--------|-----------------|----------------------|------------------|
|-------|-------------|--------|-----------------|----------------------|------------------|

Table 10. Community 1.5 plant community composition

| Group | Common Name | Symbol | Scientific Name | Annual Production () | Foliar Cover (%) |
|-------|-------------|--------|-----------------|----------------------|------------------|
|-------|-------------|--------|-----------------|----------------------|------------------|

References

. (Date accessed). **Fire Effects Information System**. <http://www.fs.fed.us/database/feis/>.

. 2021 (Date accessed). **USDA PLANTS Database**. <http://plants.usda.gov>.

## Other references

Agee, J.K., 1993. Fire Ecology of Pacific Northwest Forests. Inland Press, Washington, DC.

Anderson, Michelle D. 2003. *Pinus contorta* var. *latifolia*. In: Fire Effects Information System, [Online]. U.S. Department of Agriculture, Forest Service, Rocky Mountain Research Station, Fire Sciences Laboratory (Producer). Available: <https://www.fs.fed.us/database/feis/plants/tree/pincon/all.html> [2020, June 17].

Arno, S. (2000). Fire in western forest ecosystems (chapter 5). In: J.K Brown and J.K. Smith (eds), pp. 97-120, Wildland fire in ecosystems: effects of fire on flora. USDA Forest Service, General Technical Report RMRS-GTR-42-vol 2.

Cochran, P. H., J.M. Geist, D.L Clemens, Rodrick R. Clausnitzer, and David C. Powell. 1994. Suggested Stocking Levels for Forest Stands in Northeastern Oregon and Southeastern Washington. Research Note PNW-RN-513. Bend, OR. U.S. Department of Agriculture, Forest Service. 21 p.

Halofsky, J.E., Peterson, D.L. & Harvey, B.J. Changing wildfire, changing forests: the effects of climate change on fire regimes and vegetation in the Pacific Northwest, USA. *fire ecol* 16, 4 (2020). <https://doi.org/10.1186/s42408-019-0062-8>

Hood, Sharon; Abrahamson, Ilana; and Cansler, C. Alina. 2018. Fire resistance and regeneration characteristics of Northern Rockies tree species. In: Fire Effects Information System, [Online]. U.S. Department of Agriculture, Forest Service, Rocky Mountain Research Station, Missoula Fire Sciences Laboratory (Producer). Available: <https://www.fs.fed.us/database/feis/pdfs/other/FireResistRegen.html>.

Johnson, CG, Jr.; Clausnitzer, R.R. 1992. Plant Association of the Blue and Ochoco Mountains. Tech. Publ. R6-ERW-TP-036-92. Portland, OR; U.S. Department of Agriculture, Forest Service, Pacific Northwest Region, Wallow-Whitman National Forest. 164 p.

Johnson, CG, Jr.; Simon, Steven A. Plant Association of the Walla-Snake Province, Wallowa-Whitman National Forest. R6-ECOL-TP-225A-86. U.S. Department of Agriculture, Forest Service, Pacific Northwest Region, Wallow-Whitman National Forest. 400 p.

Juran, Ashley G. 2017. Fire regimes of conifer forests in the Blue Mountains. In: Fire Effects Information System, [Online]. U.S. Department of Agriculture, Forest Service, Rocky Mountain Research Station, Missoula Fire Sciences Laboratory (Producer).

LANDFIRE: LANDFIRE National Vegetation Dynamics Models. (2007, January – last update). BpS 10550 (2013, October) and 10560 (2007, October). [Homepage of the LANDFIRE Project, U.S. Department of Agriculture, Forest Service; U.S. Department of Interior], [Online]. Available [accessed 2020, March 27].

Miller, M. 2000. Chapter 2: Fire Autecology. In: Brown, J.K.; Smith, J.K., editors. Wildland fire in ecosystems: effects of fire on flora. Gen. Tech. Report RMRS-GTR-42- vol. 2. Ogden, UT: Rocky Mountain Research Station, Forest Service, U.S. Department of Agriculture. 257 p.

NatureServe. 2020. "Rocky Mountain Subalpine Dry-Mesic Spruce-Fir Forest & Woodland". NatureServe, Arlington, Virginia. (Accessed: September 15, 2020).

Powell, D.C.; Johnson, C.G.; Crowe, E.A.; Wells, A.; Swanson, D.K. 2007. Potential vegetation hierarchy for the Blue Mountains section of northeastern Oregon, southeastern Washington, and west-central Idaho. Gen. Tech. Rep. PNW-GTR-709. Portland, OR: U.S. Department of Agriculture, Forest Service, Pacific Northwest Research Station. 87 p.

Uchytel, Ronald J. 1991. *Abies lasiocarpa*. In: Fire Effects Information System, [Online]. U.S. Department of Agriculture, Forest Service, Rocky Mountain Research Station, Fire Sciences Laboratory (Producer). Available: <https://www.fs.fed.us/database/feis/plants/tree/abilas/all.html> [2020, June 17].

Uchytel, Ronald J. 1991. *Picea engelmannii*. In: Fire Effects Information System, [Online]. U.S. Department of Agriculture, Forest Service, Rocky Mountain Research Station, Fire Sciences Laboratory (Producer). Available: <https://www.fs.fed.us/database/feis/plants/tree/piceng/all.html> [2020, June 17].

United States National Vegetation Classification (2008) - A3643 Rocky Mountain Dry-Mesic Subalpine Fir-Engelmann Spruce Forest.

## Contributors

Andrew Neary - Concept development for 2020 PES initiative  
Kurt Moffit - Initial PES grouping

## Approval

Kirt Walstad, 4/04/2025

## Rangeland health reference sheet

**Interpreting Indicators of Rangeland Health** is a qualitative assessment protocol used to determine ecosystem condition based on benchmark characteristics described in the Reference Sheet. A suite of 17 (or more) indicators are typically considered in an assessment. The ecological site(s) representative of an assessment location must be known prior to applying the protocol and must be verified based on soils and climate. Current plant community cannot be used to identify the ecological site.

|                                             |                   |
|---------------------------------------------|-------------------|
| Author(s)/participant(s)                    |                   |
| Contact for lead author                     |                   |
| Date                                        | 02/12/2025        |
| Approved by                                 |                   |
| Approval date                               |                   |
| Composition (Indicators 10 and 12) based on | Annual Production |

## Indicators

### 1. Number and extent of rills:

---

### 2. Presence of water flow patterns:

---

### 3. Number and height of erosional pedestals or terracettes:

---

### 4. Bare ground from Ecological Site Description or other studies (rock, litter, lichen, moss, plant canopy are not bare ground):

---

### 5. Number of gullies and erosion associated with gullies:

---

### 6. Extent of wind scoured, blowouts and/or depositional areas:

---

7. Amount of litter movement (describe size and distance expected to travel):

---

8. Soil surface (top few mm) resistance to erosion (stability values are averages - most sites will show a range of values):

---

9. Soil surface structure and SOM content (include type of structure and A-horizon color and thickness):

---

10. Effect of community phase composition (relative proportion of different functional groups) and spatial distribution on infiltration and runoff:

---

11. Presence and thickness of compaction layer (usually none; describe soil profile features which may be mistaken for compaction on this site):

---

12. Functional/Structural Groups (list in order of descending dominance by above-ground annual-production or live foliar cover using symbols: >>, >, = to indicate much greater than, greater than, and equal to):

Dominant:

Sub-dominant:

Other:

Additional:

---

13. Amount of plant mortality and decadence (include which functional groups are expected to show mortality or decadence):

---

14. Average percent litter cover (%) and depth ( in):

---

15. Expected annual annual-production (this is TOTAL above-ground annual-production, not just forage annual-production):

---

16. Potential invasive (including noxious) species (native and non-native). List species which BOTH characterize degraded states and have the potential to become a dominant or co-dominant species on the ecological site if their future establishment and growth is not actively controlled by management interventions. Species that become dominant for only one to several years (e.g., short-term response to drought or wildfire) are not invasive plants. Note that unlike other indicators, we are describing what is NOT expected in the reference state for the ecological site:

---

17. Perennial plant reproductive capability:

---