

Ecological site F043CY608OR

Cool Dry Conifer Foothills and Mountains (PIPO/FEID-PSSPS)

Last updated: 3/31/2025
Accessed: 08/31/2026

General information

Provisional. A provisional ecological site description has undergone quality control and quality assurance review. It contains a working state and transition model and enough information to identify the ecological site.

MLRA notes

Major Land Resource Area (MLRA): 043C–Blue and Seven Devils Mountains

This MLRA covers the Blue and Seven Devils Mountains of Oregon, Washington and Idaho. The area is characterized by thrust and block-faulted mountains and deep canyons composed of sedimentary, metasedimentary, and volcanic rocks. Elevations range from 1,300 to 9,800 feet (395 to 2,990 meters). The climate is characterized by cold, wet winters and cool, dry summers. Annual precipitation, mostly in the form of snow, averages 12 to 43 inches (305 to 1,090 millimeters) yet ranges as high as 82 inches (2,085 millimeters) at upper elevations. Soil temperature regimes are predominately Frigid to Cryic and soil moisture regimes are predominately Xeric to Udic. Mollisols and Andisols are the dominant soil orders. Ecologically, forests dominate but shrub and grass communities may occur on south aspects and lower elevations as well as in alpine meadow environments. Forest composition follows moisture, temperature and elevational gradients and typically ranges from ponderosa pine and Douglas-fir plant associations at lower elevations, grand fir at middle elevations and subalpine fir and Engelmann spruce at upper elevations. Historical fire regimes associated with these forest types range from frequent surface fires in ponderosa pine - Douglas Fir forest types to mixed and stand replacing fire regimes in grand fir and subalpine fir types. A large percentage of the MLRA is federally owned and managed by the U.S. Forest Service for multiple uses.

Classification relationships

Plant Assoc. of Blue and Ochoco Mountains (R6 E TP-036-92) Ponderosa pine/bluebunch wheatgrass - CPG111 (modal) Ponderosa pine/Idaho fescue - CPG112 (modal) Ponderosa pine/bitterbrush/Idaho fescue-bluebunch wheatgrass - CPS226 Ponderosa pine/mountain big sagebrush/Idaho fescue-bluebunch wheatgrass - CPS131 Ponderosa pine/mountain mahogany/Idaho fescue-bluebunch wheatgrass - CPS234 Ponderosa pine/mountain mahogany/Wheeler bluegrass - CPS233 Ponderosa pine/mountain snowberry - CPS525 Plant Assoc. of Wallowa-Snake Province (R6 E 255-86) Ponderosa pine/bitterbrush-bluebunch wheatgrass - CPS231 Ponderosa pine/bluebunch wheatgrass - CPG132 Ponderosa pine/Idaho fescue - CPG131 USDA Forest Service Ecological Sub-region M332 "Blue Mountains" LANDFIRE BpS model 0710531 Northern Rocky Mountain Ponderosa Pine Woodland and Savanna – Mesic Natureserve Terrestrial Ecological System (2020) Northern Rocky Mountain Ponderosa Pine Woodland and Savanna, CES306.030e

Ecological site concept

This site represents a commonly occurring ponderosa pine (*Pinus ponderosa*) site in the foothills of the Blue, Ochoco and Wallowa mountains of Oregon. The overstory is composed of almost exclusively ponderosa pine and the understory is dominated by grasses such as Idaho fescue (*Festuca idahoensis*), and bluebunch wheatgrass (*Pseudoregnaria spicata*). Western juniper (*Juniperus occidentalis*) may be occasionally present as well as antelope bitterbrush (*Purshia tridentata*). This site exists toward the lower moisture threshold of forested communities in these mountains and as such often transitions into shrublands, woodlands and grasslands. These may be grasslands dominated by Idaho fescue and bluebunch wheatgrass shrublands dominated by big sagebrush (*Artemisia tridentata*) and antelope bitterbrush, woodlands dominated by western juniper (*Juniperus occidentalis*) and mixed woodlands dominated by juniper with occasional ponderosa pine. Forested sites on adjacent protected aspects and with ashier soils with few coarse fragments, experience higher soil moisture retention facilitating higher productivity of ponderosa pine, higher shrub cover and occasional components of Douglas-fir (*Pseudotsuga menziesii*). The soil moisture regime of this site is xeric and the soil temperature regime is frigid. Disturbance was historically influenced by a fire regime characterized by relatively frequent surface fires. These frequent fires, as well as bark beetles, were historically a critical element of the disturbance regime of this site, acting to thin crowded understories and allow fire resistant ponderosa stands to attain an open, savanna-like forest structure. This is a provisional ecological site that groups characteristics at a broad scale with little to no field verification and is subject to extensive review and revision before final approval. All data herein was developed using existing information and literature and should be considered provisional and contingent upon field validation prior to use in conservation planning.

Associated sites

F043CY609OR	Warm Dry Conifer Foothills and Mountains (PIPO-PSME/SYAL/CAGE) Downslope positions with warmer soil climates
R043CY808OR	Claypan (ARAR8/FEID-PSSPS) Adjacent scablands with very shallow soils
F043CY607OR	Cool Moist Conifer Foothills and Mountains (PIPO-PSME/SYAL) Occupying adjacent soils with somewhat deeper depth to a restrictive layer and higher moisture retention

Similar sites

F043CY607OR	Cool Moist Conifer Foothills and Mountains (PIPO-PSME/SYAL) Higher available water content, somewhat deeper soils, lower coarse fragment content, andic soil properties more common
F043CY609OR	Warm Dry Conifer Foothills and Mountains (PIPO-PSME/SYAL/CAGE) Downslope positions with warmer soil climates, mesic soil temperature regime

Table 1. Dominant plant species

Tree	(1) <i>Pinus ponderosa</i>
Shrub	Not specified
Herbaceous	(1) <i>Festuca idahoensis</i> (2) <i>Pseudoroegneria spicata</i>

Physiographic features

This site occurs on forested backslopes and summits of foothill and mountain slopes. This site typically occurs on south and east aspects with slopes typically ranging from 7.5 - 45%. Elevations are typically between 4,000 to 5,200 (1,225 to 1,575 m) but may range from 3,300 to 6,000 (1,000 to 1,825 m). This site does not experience flooding or ponding and no water table is present within the upper two meters of soil.

Table 2. Representative physiographic features

Landforms	(1) Mountains > Mountain slope (2) Foothills > Hillside or mountainside
Flooding frequency	None

Ponding frequency	None
Elevation	1,220 – 1,610 m
Slope	10 – 50 %
Ponding depth	0 cm
Water table depth	250 cm
Aspect	E, SE, S, SW

Table 3. Representative physiographic features (actual ranges)

Flooding frequency	Not specified
Ponding frequency	Not specified
Elevation	1,010 – 1,830 m
Slope	0 – 70 %
Ponding depth	Not specified
Water table depth	Not specified

Climatic features

This site experiences an intermountain climate with cool, wet winters and warm, dry summers. Mean annual precipitation is typically 16 - 23 inches (405 - 585 mm) but can range from 14 to 26 inches (355 - 660 mm). Mean annual temperatures are typically 43°F (6°C) but range from 39 - 45 °F (3 - 7°C). The soil temperature regime is Frigid and the soil moisture regime is Xeric typically experiencing 60 – 90 dry days per year. Frost free days average 40 to 120 per year. Climate graphs are populated from the closest available weather station and are included to represent general trends rather than representative values.

Table 4 Representative climatic features

Frost-free period (characteristic range)	40-120 days
Freeze-free period (characteristic range)	
Precipitation total (characteristic range)	410-580 mm

Frost-free period (average)	70 days
Freeze-free period (average)	
Precipitation total (average)	460 mm

- (1) MASON DAM [USC00355258], Baker City, OR
- (2) AUSTIN 3 S [USC00350356], Prairie City, OR

Influencing water features

This site is not influenced by water from a wetland or stream.

Soil features

Soils on this site are typically shallow to moderately deep (but range to very deep). Surface textures are varied but typically range from very gravelly ashy silt loam to very gravelly ashy sandy loam. Subsurface coarse fragment content typically ranging from very gravelly to extremely gravelly or stony. The family particle size is typically loamy-skeletal but vary. These soils are typically derived from basalt colluvium and residuum and other volcanic rock and commonly have some influence of volcanic ash throughout yet lack a deep ash cap in the surface horizon. These are well drained, droughty soils, typified by an average representative available water capacity of 0.5 - 1.2 in the 0 – 10 inch depth. See Fivebit and Gaib for modal series concepts.

Table 5. Representative soil features

Parent material	(1) Colluvium – volcanic rock (2) Residuum – volcanic rock (3) Volcanic ash – volcanic rock
Surface texture	(1) Very gravelly sandy loam (2) Very gravelly silt loam
Family particle size	(1) Loamy-skeletal (2) Clayey-skeletal (3) Fine
Drainage class	Well drained
Permeability class	Slow to moderate
Depth to restrictive layer	30 – 100 cm
Soil depth	30 – 100 cm

Surface fragment cover <=3"	0 – 50 %
Surface fragment cover >3"	0 – 50 %
Available water capacity (0-101.6cm)	3.56 – 12.7 cm
Soil reaction (1:1 water) (0-101.6cm)	5.6 – 7.3
Subsurface fragment volume <=3" (10.2-152.4cm)	10 – 50 %
Subsurface fragment volume >3" (10.2-152.4cm)	10 – 40 %

Table 6. Representative soil features (actual values)

Drainage class	Not specified
Permeability class	Not specified
Depth to restrictive layer	30 – 200 cm
Soil depth	30 – 200 cm
Surface fragment cover <=3"	Not specified
Surface fragment cover >3"	Not specified
Available water capacity (0-101.6cm)	Not specified
Soil reaction (1:1 water) (0-101.6cm)	Not specified

Subsurface fragment volume <=3" (10.2-152.4cm)	Not specified
Subsurface fragment volume >3" (10.2-152.4cm)	Not specified

Ecological dynamics

The modal plant associations that define this site concept are the widespread Ponderosa pine / Idaho Fescue and Ponderosa pine / Bluebunch wheatgrass associations (PIPO/FEID and PIPO/PSSP as described in Johnson and Clausnitzer 1992, and Johnson and Simon 1987). These communities exist toward the warm and dry end of the ponderosa pine forests of MLRA 43c in Oregon. As such, productivity is lower than other ponderosa pine forests in these mountains. In its reference phase, these mature ponderosa forests would have included a perennial grass component of Idaho fescue and bluebunch wheatgrass, with bluebunch wheatgrass and cover of bare ground increasing on soils with greater surface coarse fragments. Other herbaceous species include Sandberg bluegrass (*Poa secunda*), squirreltail (*Elymus elymoides*), common yarrow (*Achillea millefolium*) and tailcup lupine (*Lupinus caudatus*). This site concept also includes less locally common ponderosa pine communities that include components of shrub and tree species adapted to droughty soils and exposures such as antelope bitterbrush, western juniper and big sagebrush. These species are especially common on the southernmost extent of this site and less common in locations further north where exposure to Columbia river weather patterns instead favors minor components of shrubs such as rose (*Rosa* sp.) and snowberry (*Symphoricarpos* sp.). Mountain mahogany (*Cercocarpus ledifolius*) may also occur in this plant community on upper slope positions and ridgetops with rocky soils.

Ponderosa pine forests were historically subject to frequent surface fires primarily ignited by lightning strikes and Native American cultural burning practices. These fires would have likely occurred at less than 35 year intervals and approximated Landfire fire regime group 1 (0-35 year frequency, surface severity). These low intensity fires would have decreased the density of young regenerating understory conifers which are less tolerant of fire when young and may otherwise act as ladder fuels to ignite crown fires and lead to stand replacing events. Overtime, frequent low intensity fires, as well as occasional mixed severity fires, would have favored the development of mature, uneven-aged ponderosa pine stands with open canopies. Mixed severity fires, where direct overstory mortality was patchy, were less common than surface fires, and true stand replacement fires were rare. Fire-resistant ponderosa is well-adapted to these conditions, developing increasing fire resistance with age by growing thick bark and self-thinning lower limbs. With longer time between fire, increased development of understory fuels such as young pine, juniper and down wood, along with the development of a closed canopy, can promote an increased frequency of stand replacing fires and insect outbreaks. In addition to direct impacts of wildfire, disturbance factors that resulted in the death of mature pine included injury from lightning strikes, wind events, weather extremes, and the collective influence of various damage agents such as bark beetles, pine engraver, mistletoe, and other adapted insects and diseases associated with injury or competitive stressors.

Understory species in these forests were adapted to respond to this fire driven disturbance regime. Many cool season perennial grasses, such as bluebunch wheatgrass, would resprout from crowns following fire. Understory forbs, such as common yarrow, may resprout from rhizomes while others, such as lupine, can effectively respond to post fire conditions that may delay the reestablishment of other species, such as low soil nutrient levels, by fixing their own nitrogen.

As a forested site with a grass dominated understory, much of these forests were historically subject to livestock grazing, especially cattle and sheep. Prolonged historical use by these ungulates may have altered the composition of understory herbaceous and shrub communities. A decrease in palatable shrubs such as bitterbrush, and perennial grasses such as Idaho fescue may have paralleled increases in forbs such as lupine and yarrow and exotic annual grasses such as cheatgrass (*Bromus tectorum*), Medusahead rye (*Taeniatherum caput-medusae*), and North Africa grass (*Ventenata dubia*). Increases in exotic grasses have been associated with other impacts as well, including off highway vehicle use and proximity to human development. At high levels, the impacts of these invasions on nutrient cycling, wildlife habitat and fire cycles may be severe.

Current stand conditions in this ecological site do not, in most instances, conform to the historic range of variability (as represented by the reference state below) in terms of the expected frequency and severity of future wildfire events. Since the arrival of Euro-American settlers to the region in the late 1880's, the character and function of these forests have changed. Logging, grazing, conversion to other uses, and fire exclusion have impacted the natural processes of this fire-dependent ecosystem. Depending on the severity and degree of impact, alternative states (which function outside of the parameters of the reference state), have developed. Within this forest type across the Blue Mountains, fire regimes are considered to have experienced "high" departure from historical conditions (Natureserve 2020).

The state and transition model below represents a generalized and simplified version of forest change in response to major disturbance types in this ecological site. It does not attempt to model the potential effects of climate change on ecosystem function or process. Emerging evidence is suggesting that climate change is leading to hotter and drier conditions in western forests that will increase fire frequency and extent and lengthen fire seasons (Halofsky et al. 2020). When combined with the interacting impacts of fire suppression,

drought, and insect outbreaks, it is possible that this ecological system will experience unpredictable ecosystem shifts and additional alternative states. For warm and dry sites, this may include the possibility of regeneration failure following wildfire disturbance (Halofsky et al. 2020). As evidence increases, this model will likely undergo alterations and updates to reflect our emerging understanding.

State and transition model

Additional community tables

Table 7. Community 1.1 plant community composition

Group	Common Name	Symbol	Scientific Name	Annual Production ()	Foliar Cover (%)
-------	-------------	--------	-----------------	----------------------	------------------

Table 8. Community 1.2 plant community composition

Group	Common Name	Symbol	Scientific Name	Annual Production ()	Foliar Cover (%)
-------	-------------	--------	-----------------	----------------------	------------------

Table 9. Community 1.3 plant community composition

Group	Common Name	Symbol	Scientific Name	Annual Production ()	Foliar Cover (%)
-------	-------------	--------	-----------------	----------------------	------------------

Table 10. Community 1.4 plant community composition

Group	Common Name	Symbol	Scientific Name	Annual Production ()	Foliar Cover (%)
-------	-------------	--------	-----------------	----------------------	------------------

Table 11. Community 1.5 plant community composition

Group	Common Name	Symbol	Scientific Name	Annual Production ()	Foliar Cover (%)
-------	-------------	--------	-----------------	----------------------	------------------

References

. (Date accessed). **Fire Effects Information System**. <http://www.fs.fed.us/database/feis/>.

. 2021 (Date accessed). **USDA PLANTS Database**. <http://plants.usda.gov>.

Other references

Agee, J.K., 1993. Fire Ecology of Pacific Northwest Forests. Inland Press, Washington, DC.

Arno, S. (2000). Fire in western forest ecosystems (chapter 5). In: J.K Brown and J.K. Smith (editors), pp. 97-120, Wildland fire in ecosystems: effects of fire on flora. USDA Forest Service, General Technical Report RMRS-GTR-42-vol 2.

Churchhill, Derek J.; Carnwath, Gunnar C.; Larson, Andrew J.; Jeronimo, Sean A. 2017. Historical forest structure, composition and spatial pattern in dry conifer forests of the western Blue Mountains, Oregon. Gen. Tech. Rep. PNW-GTR-956. Portland, OR: U.S. Department of Agriculture, Forest Service, Pacific Northwest Research Station. 93 p.

Crane, M. F.; Fischer, William C. 1986. Fire ecology of the forest habitat types of central Idaho. Gen. Tech. Rep. INT-218. Ogden, UT: U.S. Department of Agriculture, Forest Service, Intermountain Research Station. 86 p.

Fitzgerald, Stephen A. 2005. Fire Ecology of Ponderosa Pine and the Rebuilding of Fire-Resilient Ponderosa Pine Ecosystems (pg. 197-225). In "Proceedings of the Symposium on Ponderosa Pine: Issues, Trends, and Management", Gen. Tech. Rep. PSW-GTR-198. Albany, CA: Pacific Southwest Research Station, Forest Service, U.S. Department of Agriculture. 281 p.

Graham, Russell T.; Jain, Theresa B. 2005. Ponderosa Pine Ecosystems (pg. 1-32). In "Proceedings of the Symposium on Ponderosa Pine: Issues, Trends, and Management", Gen. Tech. Rep. PSW-GTR-198. Albany, CA: Pacific Southwest Research Station, Forest Service, U.S. Department of Agriculture. 281 p.

Hall, Frederick C.; Bryant, Larry; Clausnitzer, Rod (and others). 1995. Definitions and Codes of Seral Status and Structure of Vegetation. General Technical Report PNW-GTR-363. Portland, OR: USDA Forest Service, Pacific Northwest Research Station. 39 p.

Halofsky, J.E., Peterson, D.L. & Harvey, B.J. Changing wildfire, changing forests: the effects of climate change on fire regimes and vegetation in the Pacific Northwest, USA. fire ecol 16, 4 (2020). <https://doi.org/10.1186/s42408-019-0062-8>

Johnson, CG, Jr.; Simon, Steven A. 1987. Plant Association of the Walla-Snake Province, Wallowa-Whitman National Forest. R6-ECOL-

TP-225A-86. U.S. Department of Agriculture, Forest Service, Pacific Northwest Region, Wallow-Whitman National Forest. 400 p.

Johnson, CG, Jr.; Clausnitzer, R.R. 1992. Plant Association of the Blue and Ochoco Mountains. Tech. Publ. R6-ERW-TP-036-92. Portland, OR; U.S. Department of Agriculture, Forest Service, Pacific Northwest Region, Wallow-Whitman National Forest. 164 p.

Juran, Ashley G. 2017. Fire regimes of conifer forests in the Blue Mountains. In: Fire Effects Information System, [Online]. U.S. Department of Agriculture, Forest Service, Rocky Mountain Research Station, Missoula Fire Sciences Laboratory (Producer).

LANDFIRE: LANDFIRE National Vegetation Dynamics Models. (2007, January – last update). BpS 10450 and 10531. [Homepage of the LANDFIRE Project, U.S. Department of Agriculture, Forest Service; U.S. Department of Interior], [Online]. Available [2007, February 8].

Larson, A.J., Churchill, D.C., 2012. Tree spatial patterns in fire-frequent forests of western North America, including mechanisms of pattern formation and implications for designing fuel reduction and restoration treatments. *Forest Ecology and Management* 267, 74–92.

McDonald, G. I.; Harvey, A. E.; Tonn, J. R. 2000. "Fire, Competition and Forest Pests: Landscape Treatment to Sustain Ecosystem Function", in Neuenschwander, Leon F.; Ryan, Kevin C., tech. eds. *Proceedings from the Joint Fire Science Conference and Workshop: crossing the millennium: integrating spatial technologies and ecological principles for a new age in fire management*; the Grove Hotel, Boise, Idaho, June 15-17, 1999. Moscow, Idaho: University of Idaho, 2000: 195-211.

Miller, M. 2000. Chapter 2: Fire Autecology. In: Brown, J.K.; Smith, J.K., editors. *Wildland fire in ecosystems: effects of fire on flora*. Gen. Tech. Report RMRS-GTR-42- vol. 2. Ogden, UT: Rocky Mountain Research Station, Forest Service, U.S. Department of Agriculture. 257 p.

NatureServe. 2020. "Terrestrial Ecological System: Northern Rocky Mountain Ponderosa Pine Woodland and Savanna." NatureServe, Arlington, Virginia. (Accessed: September 15, 2020).

Powell, 1996 (rev 2012). A Stage is a Stage is a Stage ... Or Is It? White Paper F14-SO-WP-Silv-10. Pendleton, OR. U.S. Department of Agriculture, Forest Service, Umatilla National Forest. 14 p.

Powell, D.C. 2006 (rev 2017). Seral Status for Tree Species of the Blue and Ochoco Mountains. Tech. Publ. F14-SO-WP-SILV-58. Pendleton, OR. U.S. Department of Agriculture, Forest Service, Pacific Northwest Region, Umatilla National Forest. 17 p.

Powell, D.C.; Johnson, C.G.; Crowe, E.A.; Wells, A.; Swanson, D.K. 2007. Potential vegetation hierarchy for the Blue Mountains section of northeastern Oregon, southeastern Washington, and west-central Idaho. Gen. Tech. Rep. PNW-GTR-709. Portland, OR: U.S. Department of Agriculture, Forest Service, Pacific Northwest Research Station. 87 p.

Contributors

Kurt Moffit - Initial PES grouping

Andrew Neary - Concept development for 2020 PES initiative

Approval

Kirt Walstad, 3/31/2025

Rangeland health reference sheet

Interpreting Indicators of Rangeland Health is a qualitative assessment protocol used to determine ecosystem condition based on benchmark characteristics described in the Reference Sheet. A suite of 17 (or more) indicators are typically considered in an assessment. The ecological site(s) representative of an assessment location must be known prior to applying the protocol and must be verified based on soils and climate. Current plant community cannot be used to identify the ecological site.

Author(s)/participant(s)	
Contact for lead author	
Date	02/12/2025

Approved by	
Approval date	
Composition (Indicators 10 and 12) based on	Annual Production

Indicators

1. Number and extent of rills:

2. Presence of water flow patterns:

3. Number and height of erosional pedestals or terracettes:

4. Bare ground from Ecological Site Description or other studies (rock, litter, lichen, moss, plant canopy are not bare ground):

5. Number of gullies and erosion associated with gullies:

6. Extent of wind scoured, blowouts and/or depositional areas:

7. Amount of litter movement (describe size and distance expected to travel):

8. Soil surface (top few mm) resistance to erosion (stability values are averages - most sites will show a range of values):

9. Soil surface structure and SOM content (include type of structure and A-horizon color and thickness):

10. Effect of community phase composition (relative proportion of different functional groups) and spatial distribution on infiltration and runoff:

11. Presence and thickness of compaction layer (usually none; describe soil profile features which may be mistaken for compaction on this site):

12. Functional/Structural Groups (list in order of descending dominance by above-ground annual-production or live foliar cover using symbols: >>, >, = to indicate much greater than, greater than, and equal to):

Dominant:

Sub-dominant:

Other:

Additional:

13. Amount of plant mortality and decadence (include which functional groups are expected to show mortality or decadence):

14. Average percent litter cover (%) and depth (in):

15. Expected annual annual-production (this is TOTAL above-ground annual-production, not just forage annual-production):

16. Potential invasive (including noxious) species (native and non-native). List species which BOTH characterize degraded states and have the potential to become a dominant or co-dominant species on the ecological site if their future establishment and growth is not actively controlled by management interventions. Species that become dominant for only one to several years (e.g., short-term response to drought or wildfire) are not invasive plants. Note that unlike other indicators, we are describing what is NOT expected in the reference state for the ecological site:

17. Perennial plant reproductive capability:
