

# Ecological site EX044B01A138

## Shallow Droughty (SwDr)

### 10-14" PZ Frigid

Last updated: 2/11/2025

Accessed: 07/11/2026

---

#### General information

**Provisional.** A provisional ecological site description has undergone quality control and quality assurance review. It contains a working state and transition model and enough information to identify the ecological site.

#### MLRA notes

Major Land Resource Area (MLRA): 044B–Central Rocky Mountain Valleys

Major Land Resource Area (MLRA) 44B, Central Rocky Mountain Valleys, is nearly 3.7 million acres of southwest Montana and borders two MLRAs: 43B Central Rocky Mountains and Foothills and 46 Northern and Central Rocky Mountain Foothills. The major watersheds of this MLRA are those of the Missouri and Yellowstone Rivers and their associated headwaters such as the Beaverhead, Big Hole, Jefferson, Ruby, Madison, Gallatin, and Shields Rivers. These waters allow for extensive irrigation for crop production in an area that would generally only be compatible with rangeland and grazing. The Missouri River and its headwaters are behind several reservoirs that supply irrigation water, hydroelectric power, and municipal water. Limited portions of the MLRA are west of the Continental Divide along the Clark Fork River. The primary land use of this MLRA is production agriculture (grazing, small grain production, and hay), but there is some limited mining. Urban development is high with large expanses of rangeland converted to subdivisions for a rapidly growing population. The MLRA consists of one Land Resource Unit (LRU) and seven climate based LRU subsets. These subsets are based on a combination of Relative Effective Annual Precipitation (REAP) and frost free days. Each subset expresses a distinct set of plants that differentiate it from other LRU subsets. Annual precipitation ranges from a low of 9 inches to a high near 24 inches. The driest areas tend to be in the valley bottoms of southwest Montana in the rain shadow of the mountains. The wettest portions tend to be near the edge of the MLRA at the border with MLRA 43B. Frost free days also vary widely from less than 30 days in the Big Hole Valley to around 110 days in the warm valleys along the Yellowstone and Missouri Rivers. The plant communities of the MRLA are highly variable, but the dominant community is a cool-season grass and shrub-steppe community. Warm-season grasses have an extremely limited extent in this MLRA. Most subspecies of big sagebrush are present, to some degree, across the MLRA.

#### LRU notes

MLRA 44B has one LRU that covers the entire MLRA. The LRU has been broken into seven climate subsets based on a combination of Relative Effective Annual Precipitation (REAP) and frost free days. Each combination of REAP and frost free days results in a common plant community that is shared across the subset. Each subset is giving a letter designation of A through F for sites that do not receive additional water and Y for sites that receive additional water. LRU 01 Subset A has a REAP of nine to 14 inches (228.6-355.6mm) with a frost free days range of 70 to 110 days. This combination of REAP and frost free days results in a nearly treeless sagebrush steppe landscape. The soil moisture regime is Ustic, dry that borders on Aridic and has a Frigid soil temperature regime.

#### Classification relationships

Mueggler and Stewart. 1980. Grassland and Shrubland habitat types of Western Montana 1. *Stipa comata*/Bouteloua gracilis h.t. 2. *Agropyron spicatum*/Bouteloua gracilis h.t. Montana Natural Heritage Program Vegetation Classification 1. *Stipa comata* - Bouteloua gracilis Herbaceous Vegetation (STICOM – BOUGRA) Needle and thread/Blue grama Natural Heritage Conservation Rank-G5 / S5 Edition / Author- 99-11-16 / S.V. Cooper, EPA Ecoregions of Montana, Second Edition: Level I: Northwestern Forested Mountains Level II: Western Cordillera Level III: Middle Rockies & Northern Great Plains Level IV: Paradise Valley Townsend Basin Dry Intermontane Sagebrush Valleys Shield-Smith Valleys National Hierarchical Framework of Ecological Units: Domain: Dry Division: M330 – Temperate Steppe Division – Mountain Provinces Province: M332 –Middle Rocky Mountain Steppe – Coniferous Forest – Alpine Meadow Section: M332D – Belt Mountains Section M332E – Beaverhead Mountains Section Subsection: M332Ej – Southwest Montana Intermontane Basins and Valleys M332Dk – Central Montana Broad Valleys

#### Ecological site concept

The Shallow Droughty ecological site is an upland site formed from residuum of non-carbonatic geology. The site does not receive additional moisture from a water table or flooding. It has 32 percent or less clay in the upper four (4) inches of the mineral surface. It is a

shallow site and has a bedrock, lithic, or paralithic root-restrictive layer within 20 inches (50cm). The surface of the site has less than 15 percent stone. The Shallow Droughty is skeletal, with less than 35 percent rock fragments in the 10 to 20-inch depth. The site does not have a saline or saline-sodic influence and is not strongly or violently effervescent within four inches of the mineral surface. Calcium carbonates may increase with depth.

### Associated sites

EX044B01A038	<p><b>Droughty Steep (DrStp) 10-14" PZ Frigid</b></p> <p>Droughty Steep often resides lower on the landscape as the Shallow, Droughty site on slopes greater than 15 percent</p>
EX044B01A136	<p><b>Shallow Loamy (SwLo) 10-14" PZ Frigid</b></p> <p>The Shallow, Loamy and Shallow Droughty sites exist on the same landscape and are often neighboring sites</p>
EX044B01A036	<p><b>Droughty (Dr) 10-14" PZ Frigid</b></p> <p>This site exists in the same landscape as droughty</p>

### Similar sites

EX044B01A036	<p><b>Droughty (Dr) 10-14" PZ Frigid</b></p> <p>Droughty produces a similar plant community however does not have a root restrictive layer which allows for greater production and a slightly more resilient plant community</p>
EX044B01A136	<p><b>Shallow Loamy (SwLo) 10-14" PZ Frigid</b></p> <p>The Shallow, Loamy site shares a similar plant community and similar STMs concept</p>

Table 1. Dominant plant species

Tree	Not specified
Shrub	(1) <i>Artemisia tridentata</i> (2) <i>Tetradymia canescens</i>
Herbaceous	(1) <i>Pseudoroegneria spicata</i> (2) <i>Hesperostipa comata</i>

### Legacy ID

R044BA138MT

### Physiographic features

This ecological site occurs on nearly level to very steep hills, escarpments, and buttes. It often occurs in complexes with other ecological sites. This site can be found on all slopes and exposures, but it is most common on slopes lower than 25 percent. Aspect can have an impact on the composition and production of plant communities. The amount of rock outcrop tends to increase as slopes increase.

Table 2. Representative physiographic features

Geomorphic position, hills	(1) Crest
Landforms	(1) Intermontane basin > Escarpment (2) Intermontane basin > Butte (3) Intermontane basin > Ridge (4) Intermontane basin > Hill
Runoff class	Medium to high
Elevation	1,280 – 1,980 m
Slope	0 – 30 %
Aspect	W, NW, N, NE, E, SE, S, SW

**Table 3. Representative physiographic features (actual ranges)**

Runoff class	Not specified
Elevation	Not specified
Slope	0 – 50 %

### Climatic features

The Central Rocky Mountain Valleys MLRA has a continental climate. Fifty to sixty percent of the annual long-term average total precipitation falls between May and August. Snow on frozen ground makes up the majority of winter precipitation. Average precipitation for LRU 01 Subset A is 12 inches (305mm), and the frost-free period averages 78 days. Precipitation is highest in May and June. Some of Montana's driest areas are located in sheltered mountain valleys because of the rain-shadow effects on the leeward side of some ranges.

**Table 4 Representative climatic features**

Frost-free period (characteristic range)	70-110 days
Freeze-free period (characteristic range)	110-140 days
Precipitation total (characteristic range)	230-360 mm
Frost-free period (actual range)	70-110 days
Freeze-free period (actual range)	110-140 days

Precipitation total (actual range)	230-360 mm
Frost-free period (average)	80 days
Freeze-free period (average)	130 days
Precipitation total (average)	310 mm

- (1) DEER LODGE 3 W [USC00242275], Deer Lodge, MT
- (2) DILLION U OF MONTANA WESTERN [USC00242409], Dillon, MT
- (3) GLEN 2 E [USC00243570], Dillon, MT
- (4) ENNIS [USC00242793], Ennis, MT
- (5) BOULDER [USC00241008], Boulder, MT
- (6) GARDINER [USC00243378], Gardiner, MT
- (7) TOWNSEND [USC00248324], Townsend, MT
- (8) TRIDENT [USC00248363], Three Forks, MT
- (9) TWIN BRIDGES [USC00248430], Sheridan, MT
- (10) WHITE SULPHUR SPRNGS 2 [USC00248930], White Sulphur Springs, MT
- (11) DILLON AP [USW00024138], Dillon, MT
- (12) HELENA RGNL AP [USW00024144], Helena, MT

### Influencing water features

The site has a root-restrictive layer that may affect water infiltration; however, the site is considered water-limited due to its low available water holding capacity. Runoff is medium to high.

### Wetland description

This site is not associated with wetlands characteristics.

### Soil features

These soils are shallow (10 to 20 inches deep to root restrictive layer), moderate to moderately rapid permeability, and are well to somewhat excessively drained. These soils formed from residuum from mixed origins primarily from non-calcareous geology. Typically soil surface textures consist of loam, clay loam, and silt loam textures. Soils may have gravelly surface however will vary depending on its association with to a neighboring sites. The 10- to 20-inch section of the soil below the surface will have greater than 35 percent rock fragments. Common soils series in this ecological site includes Blackleaf and Gnojek. These soils may exist across multiple ecological sites due to natural variations in slope, texture, rock fragments, and pH.

Table 5. Representative soil features

Parent material	(1) Residuum – igneous, metamorphic and sedimentary rock
Surface texture	(1) Loam (2) Silt loam (3) Clay loam
Drainage class	Well drained to somewhat excessively drained

Permeability class	Moderate to moderately rapid
Depth to restrictive layer	30 – 50 cm
Soil depth	30 – 50 cm
Surface fragment cover <=3"	0 – 30 %
Surface fragment cover >3"	0 – 10 %
Available water capacity (0-50.8cm)	2.03 – 7.37 cm
Soil reaction (1:1 water) (0-50.8cm)	6.6 – 10
Subsurface fragment volume <=3" (25.4-50.8cm)	20 – 40 %
Subsurface fragment volume >3" (25.4-50.8cm)	0 – 20 %

## Ecological dynamics

The Shallow Droughty ecological site of MLRA 44B in the 9 to 14 inch climatic subset shares plant communities and state and transition model with Shallow Loamy and Shallow Droughty ecological sites in both MLRAs 44B and 43B in the 9 to 14 inch climatic subset. These sites are nearly indistinguishable.

The reference plant community is dominated by bluebunch wheatgrass (*Pseudoroegneria spicata*) and needle and thread (*Hesperostipa comata*). Subdominant species may include green needlegrass (*Nassella viridula*), gray horsebrush (*Tetradymia canescens*), Wyoming big sagebrush (*Artemisia tridentata* ssp. *wyomingensis*), and Indian ricegrass (*Achnatherum hymenoides*).

Natural fire was a major ecological driver of this entire ecological site. Fire tended to restrict tree and sagebrush growth to small patches and promote an herbaceous plant community. The natural fire return interval was highly variable, ranging up to 100 years; however, it was likely shorter than 35 years (Arno and Gruell 1983). Since 1910, there has been a significant increase in the suppression of fire in sagebrush and trees.

Wyoming big sagebrush steppe communities historically had low fuel loadings and were characterized by 10- to 70-year fire intervals that produced a mosaic of burned and unburned lands (Bunting et al., 1987). A shift to the dominance of shrubs may occur in response to improper grazing management, drought, or where big sagebrush occurs due to a lack of fire. Shrub encroachment by a variety of species, including broom snakeweed (*Gutierrezia sarothrae*), fringed sagewort (*Artemisia frigida*), Wyoming big sagebrush (*Artemisia tridentata* ssp. *wyomingensis*), rubber rabbitbrush (*Ericameria nauseosa*), green rabbitbrush (*Chrysothamnus viscidiflorus*), and plains prickly pear cactus (*Opuntia polyacantha*) occurs within this site as the mid-stature bunchgrasses decrease. Shrub dominance and grass loss can be associated with soil erosion and, ultimately, thinning of the native soil surface. Subsequent loss of soil could lead to a Degraded State. All states could also lead to the Invaded State when there is a lack of weed prevention and control measures.

Historical records indicate that, prior to the introduction of livestock (cattle and sheep) during the late 1800s, elk and bison grazed this

ecological site. Because of the nomadic nature and herd structure of bison, grazed areas received periodic high intensity, short duration grazing pressure. Forage for livestock was noted as minimal in areas recently grazed by bison (Lesica and Cooper 1997). Meriwether Lewis documented that he was met by 60 Shoshone warriors on horseback in August 1805, and the Corps of Discovery was later supplied with horses by the same band of Shoshone. This suggests that the areas near the modern-day Montana towns of Twin Bridges, Dillon, Grant, and Dell were grazed by an untold number of horses prior to the large introduction of cattle and sheep. Livestock grazing has occurred on most of this ecological site in southwestern Montana for more than 150 years. The gold boom in the 1860s brought the first herds of livestock overland from Texas, and homesteaders began settling the area. During this time, cattle were the primary domestic grazers in the area. In the 1890s, Montana sheep production began to increase and dominated the livestock industry until the 1930s. Since the 1930s, cattle production has dominated the livestock industry in the region (Wyckoff and Hansen 2001).

Some of the major invasive species that can occur on this site include (but are not limited to) spotted knapweed (*Centaurea stoebe*), leafy spurge (*Euphorbia esula*), cheatgrass (*Bromus tectorum*), field brome (*Bromus arevensis*), yellow toadflax (*Linaria vulgaris*), and dandelion (*Taraxicum spp.*). Invasive weeds are beginning to have a high impact on this ecological site, particularly cheatgrass invasion.

#### Plant Communities and Transitions

A state and transition model for this ecological site is depicted below. Thorough descriptions of each state, transition, plant community, and pathway follow the model. This model is based on available experimental research, field data, field observations, and interpretations by experts. It is likely to change as knowledge increases.

The plant communities within the same ecological site will differ across the MLRA due to the naturally occurring variability in weather, soils, and aspect. The biological processes on this site are complex; therefore, representative values are presented in a land management context. The species lists are representative and are not botanical descriptions of all species occurring, or potentially occurring, on this site. They are intended to cover the core species and the known range of conditions and responses.

Although there is considerable qualitative experience supporting the pathways and transitions within the state and transition model (STM), no quantitative information exists that specifically identifies threshold parameters between grassland types and invaded types in this ecological site.

## State and transition model

### Additional community tables

Table 6. Community 1.1 plant community composition

Group	Common Name	Symbol	Scientific Name	Annual Production (t)	Foliar Cover (%)
<b>Grass/Grasslike</b>					
1	<b>Deep Rooted Bunchgrasses</b>			314-650	
	bluebunch wheatgrass	PSSP6	<i>Pseudoroegneria spicata</i>	336-616	35-50
	green needlegrass	NAVI4	<i>Nassella viridula</i>	67-112	5-10
	Indian ricegrass	ACHY	<i>Achnatherum hymenoides</i>	22-45	1-5
	rough fescue	FECA4	<i>Festuca campestris</i>	0-11	0-1
2	<b>Shortgrasses</b>			67-112	
	Sandberg bluegrass	POSE	<i>Poa secunda</i>	22-45	5-10
	prairie Junegrass	KOMA	<i>Koeleria macrantha</i>	22-45	5-10
	blue grama	BOGR2	<i>Bouteloua gracilis</i>	0-22	0-5
	needleleaf sedge	CADU6	<i>Carex duriuscula</i>	6-22	1-3
	threadleaf sedge	CAFI	<i>Carex filifolia</i>	6-22	1-3
4	<b>Rhizomatous Grasses</b>			22-90	
	thickspike wheatgrass	ELLA3	<i>Elymus lanceolatus</i>	22-67	0-5
	western wheatgrass	PASM	<i>Pascopyrum smithii</i>	22-67	0-5
	plains reedgrass	CAMO	<i>Calamagrostis montanensis</i>	11-22	0-2
<b>Shrub/Vine</b>					
3	<b>Shrubs</b>			34-101	
	Wyoming big sagebrush	ARTRW8	<i>Artemisia tridentata ssp. wyomingensis</i>	67-90	5-15
	little sagebrush	ARAR8	<i>Artemisia arbuscula</i>	0-45	0-5
	yellow rabbitbrush	CHVI8	<i>Chrysothamnus viscidiflorus</i>	22-45	1-4
	spineless horsebrush	TECA2	<i>Tetradymia canescens</i>	22-45	1-3

	Woods' rose	ROWO	<i>Rosa woodsii</i>	11-34	0-3
	broom snakeweed	GUSA2	<i>Gutierrezia sarothrae</i>	0-22	0-2
	chokecherry	PRVI	<i>Prunus virginiana</i>	0-17	0-5
	black sagebrush	ARNO4	<i>Artemisia nova</i>	0-11	0-3
	currant	RIBES	<i>Ribes</i>	0-11	0-3
	plains pricklypear	OPPO	<i>Opuntia polyacantha</i>	0-1	0-1
<b>Forb</b>					
5	<b>Forbs</b>			11-45	
	arrowleaf balsamroot	BASA3	<i>Balsamorhiza sagittata</i>	6-22	0-5
	American vetch	VIAM	<i>Vicia americana</i>	6-17	0-3
	dotted blazing star	LIPU	<i>Liatris punctata</i>	6-17	0-3
	stemless mock goldenweed	STAC	<i>Stenotus acaulis</i>	0-11	0-3
	spiny phlox	PHHO	<i>Phlox hoodii</i>	0-6	0-3
	Bonneville shootingstar	DOCO	<i>Dodecatheon conjugens</i>	0-6	0-3
	common yarrow	ACMI2	<i>Achillea millefolium</i>	0-6	0-3
	desertparsley	LOMAT	<i>Lomatium</i>	0-6	0-3
	bastard toadflax	COUM	<i>Comandra umbellata</i>	0-6	0-3
	milkvetch	ASTRA	<i>Astragalus</i>	0-6	0-2
	Lava aster	IOAL	<i>Ionactis alpina</i>	0-6	0-1
<b>Tree</b>					
6	<b>Coniferous Trees/Tall Shrubs</b>			0-11	
	ponderosa pine	PIPO	<i>Pinus ponderosa</i>	0-11	0-1
	Douglas-fir	PSME	<i>Pseudotsuga menziesii</i>	0-11	0-1
	Rocky Mountain juniper	JUSC2	<i>Juniperus scopulorum</i>	0-11	0-1
	limber pine	PIFL2	<i>Pinus flexilis</i>	0-11	0-1

Table 7. Community 1.2 plant community composition

Group	Common Name	Symbol	Scientific Name	Annual Production ()	Foliar Cover (%)
-------	-------------	--------	-----------------	----------------------	------------------

Table 8. Community 2.1 plant community composition

Group	Common Name	Symbol	Scientific Name	Annual Production ()	Foliar Cover (%)
-------	-------------	--------	-----------------	----------------------	------------------

Table 9. Community 3.1 plant community composition

Group	Common Name	Symbol	Scientific Name	Annual Production ()	Foliar Cover (%)
-------	-------------	--------	-----------------	----------------------	------------------

Table 10. Community 4.1 plant community composition

Group	Common Name	Symbol	Scientific Name	Annual Production ()	Foliar Cover (%)
-------	-------------	--------	-----------------	----------------------	------------------

Table 11. Community 5.1 plant community composition

Group	Common Name	Symbol	Scientific Name	Annual Production ()	Foliar Cover (%)
-------	-------------	--------	-----------------	----------------------	------------------

## Animal community

This site provides a variety of wildlife habitats for an array of species. Prior to the settlement of this area, large herds of antelope, elk, and bison roamed. Though the bison have been replaced, mostly with domesticated livestock, elk and antelope still frequently utilize this largely intact landscape for winter habitat in areas adjacent to forests. Sites with large quantities of curl-leaf mountain mahogany are considered important winter range for mule deer, elk, and moose. In some areas, it may even be considered critical habitat for dwindling wild ungulate populations. Managed livestock grazing is suitable on this site due to the potential to produce an abundance of high quality forage. This is often a preferred site for grazing by livestock, and animals tend to congregate in these areas. In order to maintain the productivity of the site, grazing on adjoining sites with less production must be managed carefully to make sure utilization on this site is

not excessive. Management objectives should include maintenance or improvement of the native plant community. Careful management of the timing and duration of grazing is important. Shorter grazing periods and adequate deferment during the growing season are recommended for plant maintenance, health, and recovery. According to McLean et al., early-season defoliation of bluebunch wheatgrass can result in high mortality and reduced vigor in plants. They also suggest, based on prior studies, that the opportunity for regrowth is necessary before dormancy to reduce injury bluebunch. Since needle and thread normally matures earlier than bluebunch wheatgrass and produces a sharp awn, this species is usually avoided after seed set. Changing the grazing season will allow needle and thread to be used more efficiently. Continual non-prescribed grazing of this site will be injurious, will alter the plant composition and production over time, and will result in the transition to the Altered State. The transition to other states will depend on the duration of poorly managed grazing as well as other circumstances such as weather conditions and fire frequency. The Altered State can degrade further to the Degraded State or the Invaded State. Management should focus on grazing management strategies that will prevent further degradation, such as seasonal grazing deferment or winter grazing where feasible. Communities within this state are still stable and healthy under proper management. Forage quantity and/or quality may be substantially decreased from the Reference State. Grazing is possible in the Invaded State. Invasive species are generally less palatable than native grasses. Forage production is typically greatly reduced in this state. Due to the aggressive nature of invasive species, sites in the Invaded State face an increased risk of further degradation. Grazing has to be carefully managed to avoid further soil loss and degradation and possible livestock health issues. Prescribed grazing can be used to manage invasive species. In some instances, carefully targeted grazing (sometimes in combination with other treatments) can reduce or maintain the species composition of invasive species. In the Degraded State, grazing may be possible but is generally not economically or environmentally sustainable.

## Hydrological functions

The hydrologic cycle functions best in the Reference State (1) with good infiltration and deep percolation of rainfall; however, the cycle degrades as the vegetation community declines. Rapid rainfall infiltration, high soil organic matter, good soil structure, and good porosity accompany high bunchgrass canopy cover. High ground cover reduces raindrop impact on the soil surface, which keeps erosion and sedimentation transport low. Water leaving the site will have a minimal sediment load, which allows for high water quality in associated streams. High rates of infiltration will allow water to move below the rooting zone during periods of heavy rainfall. The Bluebunch Wheatgrass Community (1.1) should have no rills or gullies present, and drainage ways should be vegetated and stable. Water flow patterns, if present, will be barely observable. Plant pedestals are essentially nonexistent. Plant litter remains in place and is not moved by wind or water. Improper grazing management results in a community shift to the Mixed Bunchgrass Community (1.2). This plant community has a similar canopy cover, but bare ground may increase to more than 15 percent. Therefore, the hydrologic cycle is functioning at a level similar to the water cycle in the Bluebunch Wheatgrass Community (1.1). Compared to the Community (1.1), the infiltration rates are slightly reduced, and surface runoff is slightly higher. In the Degraded State (3) and the Invaded State (4), canopy and ground cover are greatly reduced compared to the Reference State (1), which impedes the hydrologic cycle. Infiltration will decrease and runoff will increase due to reduced ground cover, the presence of shallow-rooted species, rainfall splash, soil capping, reduced organic matter, and poor structure. Sparse ground cover and decreased infiltration can combine to increase the frequency and severity of flooding within a watershed. Soil erosion is accelerated, the quality of surface runoff is poor, and sedimentation increases. The hydrology of the Conifer Encroached State (5) is highly variable, but studies suggest that an increased tree canopy affects the interception of rainfall as well as the amount of available soil moisture for herbaceous vegetation. This can negatively affect infiltration and increase runoff.

## Recreational uses

This site is often utilized for photography, hiking, hunting, bird watching, and flower collecting.

## Inventory data references

Information presented was derived from a similar site's Range Site Description (Shallow 9 –14" P.Z., Northern Rocky Mountain Valleys, South, East of Continental Divide), NRCS clipping data, literature, field observations, and personal contacts with range-trained personnel (i.e., used professional opinion of agency specialists, observations of land managers, and outside scientists).

## References

. (Date accessed). **Fire Effects Information System**. <http://www.fs.fed.us/database/feis/>.

. 2021 (Date accessed). **USDA PLANTS Database**. <http://plants.usda.gov>.

Arno, S.F. and G.E. Gruell. 1982. Fire History at the Forest-Grassland Ecotone in Southwestern Montana. *Journal of Range Management* 36:332–336.

Barrett, H. 2007. *Western Juniper Management: A Field Guide*.

- Bestelmeyer, B., J.R. Brown, J.E. Herrick, D.A. Trujillo, and K.M. Havstad. 2004. Land Management in the American Southwest: a state-and-transition approach to ecosystem complexity. *Environmental Management* 34:38–51.
- Bestelmeyer, B. and J. Brown. 2005. State-and-Transition Models 101: A Fresh look at vegetation change.
- Blaisdell, J.P. 1958. Seasonal development and yield of native plants on the Upper Snake River Plains and their relation to certain climate factors.
- Blaisdell, J.P. and R.C. Holmgren. 1984. Managing Intermountain Rangelands--Salt-Desert Shrub Ranges. General Tech Report INT-163. USDA Forest Service, Intermountain Forest and Range Experiment Station, Ogden, UT. 52.
- Bunting, S.C., B.M. Kilgore, and C.L. Bushey. 1987. Guidelines for Prescribe burning sagebrush-grass rangelands in the Northern Great Basin. General Technical Report INT-231. USDA Forest Service Intermountain Research Station, Ogden, UT. 33.
- Colberg, T.J. and J.T. Romo. 2003. Clubmoss effects on plant water status and standing crop. *Journal of Range Management* 56:489–495.
- Daubenmire, R. 1970. Steppe vegetation of Washington.
- DiTomaso, J.M. 2000. Invasive weeds in Rangelands: Species, Impacts, and Management. *Weed Science* 48:255–265.
- Dormaar, J.F., B.W. Adams, and W.D. Willms. 1997. Impacts of rotational grazing on mixed prairie soils and vegetation. *Journal of Range Management* 50:647–651.
- Hobbs, J.R. and S.E. Humphries. 1995. An integrated approach to the ecology and management of plant invasions. *Conservation Biology* 9:761–770.
- Kuchler, A.W. 1964. Potential natural vegetation of the conterminous United States.
- Lacey, J.R., C.B. Marlow, and J.R. Lane. 1989. Influence of Spotted knapweed (*Centaurea maculosa*) on surface runoff and sediment yield.. *Weed Technology* 3:627–630.
- Lesica, P. and S.V. Cooper. 1997. Presettlement vegetation of Southern Beaverhead County, MT.
- Manske, L.L. 1980. Habitat, phenology, and growth of selected sandhills range plants.
- Masters, R. and R. Sheley. 2001. Principles and practices for managing rangeland invasive plants. *Journal of Range Management* 38:21–26.
- McCalla, G.R., W.H. Blackburn, and L.B. Merrill. 1984. Effects of Livestock Grazing on Infiltration Rates of the Edwards Plateau of Texas. *Journal of Range Management* 37:265–269.
- McLean, A. and S. Wikeem. 1985. Influence of season and intensity of defoliation on bluebunch wheatgrass survival and vigor in southern British Columbia. *Journal of Range Management* 38:21–26.

- Miller, R.F., T.J. Svejcar, and J.A. Rose. 2000. Impacts of western juniper on plant community composition and structure. *Journal of Range Management* 53:574–585.
- Moulton, G.E. and T.W. Dunlay. 1988. *The Journals of the Lewis and Clark Expedition*. Pages in University of Nebraska Press.
- Mueggler, W.F. and W.L. Stewart. 1980. *Grassland and Shrubland Habitat Types of Western Montana*.
- Pelant, M., P. Shaver, D.A. Pyke, and J.E. Herrick. 2005. Interpreting Indicators of Rangeland Health.
- Pellant, M. and L. Reichert. 1984. Management and Rehabilitation of a burned winterfat community in Southwestern Idaho. Proceedings-- Symposium on the biology of Atriplex and related Chenopods. 1983 May 2-6; Provo UT General Technical Report INT-172.. USDA Forest Service Intermountain Forest and Range Experiment Station. 281–285.
- Pitt, M.D. and B.M. Wikeem. 1990. Phenological patterns and adaptations in an Artemisia/Agropyron plant community. *Journal of Range Management* 43:350–357.
- Pokorny, M.L., R. Sheley, C.A. Zabinski, R. Engel, T.J. Svejcar, and J.J. Borkowski. 2005. Plant Functional Group Diversity as a Mechanism for Invasion Resistance.
- Ross, R.L., E.P. Murray, and J.G. Haigh. July 1973. **Soil and Vegetation of Near-pristine sites in Montana**.
- Schoeneberger, P.J. and D.A. Wysocki. 2017. **Geomorphic Description System, Version 5.0.**
- Smoliak, S., R.L. Ditterlin, J.D. Scheetz, L.K. Holzworth, J.R. Sims, L.E. Wiesner, D.E. Baldrige, and G.L. Tibke. 2006. *Montana Interagency Plant Materials Handbook*.
- Stavi, I. 2012. The potential use of biochar in reclaiming degraded rangelands. *Journal of Environmental Planning and Management* 55:1–9.
- Stringham, T.K., W.C. Kreuger, and P.L. Shaver. 2003. State and Transition Modeling: an ecological process approach. *Journal of Range Management* 56:106–113.
- Stringham, T.K. and W.C. Krueger. 2001. States, Transitions, and Thresholds: Further refinement fro rangeland applications.
- Sturm, J.J. 1954. *A study of a relict area in Northern Montana*. University of Wyoming, Laramie 37.
- Thurow, T.L., Blackburn W. H., and L.B. Merrill. 1986. Impacts of Livestock Grazing Systems on Watershed. Page in *Rangelands: A Resource Under Siege: Proceedings of the Second International Rangeland Congress*.
- Various NRCS Staff. 2013. *National Range and Pasture Handbook*.
- Walker, L.R. and S.D. Smith. 1997. Impacts of invasive plants on community and ecosystem properties. Pages 69–86 in *Assessment and management of plant invasions*. Springer, New York, NY.

Wambolt, C. and G. Payne. 1986. An 18-Year Comparison of Control Methods for Wyoming Big Sagebrush in Southwestern Montana. *Journal of Range Management* 39:314–319.

West, N.E. 1994. Effects of Fire on Salt-Desert shrub rangelands. *Proceedings--Ecology and Management of Annual Rangelands: 1992 May 18-22. Boise ID General Technical Report INT-GTR-313.. USDA Forest Service Intermountain Research Station.* 71–74.

Whitford, W.G., E.F. Aldon, D.W. Freckman, Y. Steinberger, and L.W. Parker. 1989. Effects of Organic Amendments on Soil Biota on a Degraded Rangeland. *Journal of Range Management* 41:56–60.

Wilson, A.M., G.A. Harris, and D.H. Gates. 1966. Cumulative Effects of Clipping on Yield of Bluebunch wheatgrass. *Journal of Range Management* 19:90–91.

## Contributors

Kirt Walstad  
Benjamin Moore  
Braden Pitcher  
Ken Scalzone  
Eve Wills

## Approval

Kirt Walstad, 2/11/2025

## Rangeland health reference sheet

**Interpreting Indicators of Rangeland Health** is a qualitative assessment protocol used to determine ecosystem condition based on benchmark characteristics described in the Reference Sheet. A suite of 17 (or more) indicators are typically considered in an assessment. The ecological site(s) representative of an assessment location must be known prior to applying the protocol and must be verified based on soils and climate. Current plant community cannot be used to identify the ecological site.

Author(s)/participant(s)	Grant Petersen
Contact for lead author	grant.petersen@usda.gov
Date	02/11/2025
Approved by	
Approval date	
Composition (Indicators 10 and 12) based on	Annual Production

## Indicators

1. **Number and extent of rills:** Rills will primarily be absent on gentle slopes however on the steepest of slopes of this site (greater than 30 percent) small, short rills (less than 2-3 feet) may be evident after high precipitation events.

---
2. **Presence of water flow patterns:** Water flow patterns are rare in the reference condition. If present, they are most likely to occur on steeper slopes (greater than 20 percent) and are inconspicuous, disconnected, and very short in length.

---
3. **Number and height of erosional pedestals or terracettes:** Pedestals are not evident in the reference condition.

---
4. **Bare ground from Ecological Site Description or other studies (rock, litter, lichen, moss, plant canopy are not bare ground):**

---
5. **Number of gullies and erosion associated with gullies:** Bare ground is between 20-30 percent.

---
6. **Extent of wind scoured, blowouts and/or depositional areas:**

---
7. **Amount of litter movement (describe size and distance expected to travel):** Wind scoured, or depositional areas are not evident in the reference condition.

---
8. **Soil surface (top few mm) resistance to erosion (stability values are averages - most sites will show a range of values):** Soil Surface Stable with Stability Ratings of 4-6 (both under canopy and bare). Biotic crusts and or root mats may be present.

---
9. **Soil surface structure and SOM content (include type of structure and A-horizon color and thickness):** Structure trends to weak, fine granular. The A horizon is approximately 3 inches thick with wet Munsell colors Value 5 or less, Chroma 3 or less. Dry colors tend to be quite light prior to wetting. Official Series Description (OSD) for characteristic range.

---
10. **Effect of community phase composition (relative proportion of different functional groups) and spatial distribution on infiltration and runoff:** Evenly distributed across the site, bunchgrasses improve infiltration while rhizomatous grass protects the surface from runoff forces. The Shallow droughty ecological site is well drained and has a moderately rapid infiltration rate. An even distribution of mid stature grasses, ~60-70% of site production, cool season rhizomatous grasses 10% of site production along with a mix of shortgrass (5-15%), forbs and shrubs (5-10%).

---
11. **Presence and thickness of compaction layer (usually none; describe soil profile features which may be mistaken for compaction on this site):** Not Present

---
12. **Functional/Structural Groups (list in order of descending dominance by above-ground annual-production or live foliar cover using symbols: >>, >, = to indicate much greater than, greater than, and equal to):**

**Dominant:** Mid-statured, cool season, perennial bunchgrasses

**Sub-dominant:** rhizomatous grass = perennial shortgrasses/grasslikes &gt; shrubs = forbs &gt;&gt; trees

**Other:**

**Additional:**

---

- 13. Amount of plant mortality and decadence (include which functional groups are expected to show mortality or decadence):**  
Mortality in herbaceous species is not evident. Species with bunch growth forms may have some natural mortality in centers.
- 

- 14. Average percent litter cover (%) and depth ( in):** Total litter cover ranges from 30-35 percent. Most litter is irregularly distributed on the soil surface and is not at a measurable depth.
- 

- 15. Expected annual annual-production (this is TOTAL above-ground annual-production, not just forage annual-production):**  
Average annual production is 800. Low: 550 High 1050 pounds per acre. Production varies based on effective precipitation and natural variability of soil properties for this ecological site.
- 

- 16. Potential invasive (including noxious) species (native and non-native). List species which BOTH characterize degraded states and have the potential to become a dominant or co-dominant species on the ecological site if their future establishment and growth is not actively controlled by management interventions. Species that become dominant for only one to several years (e.g., short-term response to drought or wildfire) are not invasive plants. Note that unlike other indicators, we are describing what is NOT expected in the reference state for the ecological site: Potential invasive (including noxious) species (native and non-native). Invasive species on this ecological site include (but not limited to) annual brome spp., spotted knapweed, yellow toadflax, leafy spurge, ventenata, crested wheatgrass, etc. Native species such as Rocky Mountain juniper, Ponderosa pine, broom snakeweed, rabbitbrush spp., blue grama, Sandberg's bluegrass, etc. when their populations are significant enough to affect ecological function, indicate site condition departure.**
- 

- 17. Perennial plant reproductive capability:** In the reference condition, all plants are vigorous enough for reproduction either by seed or rhizomes in order to balance natural mortality with species recruitment. Density of plants indicates that plants reproduce at level sufficient to fill available resource.
-