

Ecological site R052XN176MT

Shallow to Gravel (SwGr)

10-14" p.z.

Last updated: 4/14/2025
Accessed: 09/26/2026

General information

Provisional. A provisional ecological site description has undergone quality control and quality assurance review. It contains a working state and transition model and enough information to identify the ecological site.

MLRA notes

Major Land Resource Area (MLRA): 052X–Brown Glaciated Plains

The Brown Glaciated Plains, MLRA 52, is an expansive and agriculturally and ecologically significant area. It consists of around 14.5 million acres and stretches across 350 miles from east to west, encompassing portions of 15 counties in north-central Montana. This region represents the southwestern limit of the Laurentide Ice Sheet and is considered to be the driest and westernmost area within the vast network of glacially-derived prairie pothole landforms of the northern Great Plains. Elevation ranges from 2,000 feet (610 meters) to 4,600 feet (1,400 meters). Soils are primarily Mollisols but Entisols, Inceptisols, Alfisols and Vertisols are also common. Till from continental glaciation is the predominant parent material, but alluvium and bedrock are also common. Till deposits are typically less than 50 feet thick, and in some areas glacially deformed bedrock occurs at or near the soil surface (Soller, 2001). Underlying the till is sedimentary bedrock largely consisting of Cretaceous shale, sandstone and mudstone (Vuke et al. 2007). It is commonly exposed on hillslopes, particularly along drainage ways. Significant alluvial deposits occur along glacial outwash channels and major drainages, including portions of the Missouri, Teton, Marias, Milk, and Frenchman Rivers. Large glacial lakes, particularly in the western half of the MLRA, deposited clayey and silty lacustrine sediments (Fullerton et al. 2013). Much of the western portion of this MLRA was glaciated towards the end of the Wisconsin age, and the maximum glacial extent occurred approximately 20,000 years ago (Fullerton et al., 2004). The result is a geologically young landscape that is predominantly a level till plain interspersed with lake plains and dominated by soils in the Mollisol and Vertisol orders. These soils are very productive and generally are well-suited to dryland farming. Much of this area is aridic-ustic. Crop-fallow dryland wheat farming is the predominant land use. Areas of rangeland typically are on steep hillslopes along drainages. The rangeland, much of which is native mixed grass prairie, increases in abundance in the eastern half of the MLRA. The Wisconsin-age till in the north-central part of this area typically formed large disintegration moraines with steep slopes and numerous poorly-drained potholes. A large portion of Wisconsin-age till occurring on the type of the level terrain that would typically be optimal for farming has large amounts of less-suitable sodium-affected Natrustalfs. Significant portions of Blaine, Phillips, and Valley Counties were glaciated approximately 150,000 years ago during the Illinoian age. Due to erosion and dissection of the landscape, many of these areas have steeper slopes and more exposed bedrock than areas glaciated during the Wisconsin age (Fullerton and Colton, 1986). While much of the rangeland in the aridic-ustic portion of MLRA 52 is classified as belonging to the - "dry grassland" - climatic zone, sites in portions of southern MLRA 52 may belong to the -"dry shrubland" - climatic zone. The dry shrubland zone represents the northernmost extent of the big sagebrush (*Artemisia tridentata*) steppe on the Great Plains. Because similar soils occur in both southern and northern portions of the MLRA, it is currently hypothesized that climate is the primary driving factor affecting big sagebrush distribution in this area. However the precise factors are not fully understood at this time. Sizeable tracts of largely unbroken rangeland in the eastern half of the MLRA and adjacent southern Saskatchewan are home to the Northern Montana population of greater sage grouse (*Centrocercus urophasianus*), and large portions of this area are considered to be a Priority Area for Conservation (PAC) by the U.S. Fish and Wildlife Service (U.S. Fish and Wildlife Service, 2013). This population is unique among sage grouse populations because many individuals overwinter in the big sagebrush steppe (dry shrubland) in the southern portion of the MLRA and then migrate to the northern portion of the MLRA, which lacks big sagebrush (dry grassland), to live the rest of the year (Smith, 2013). Areas of the till plain near the Bearpaw and Highwood Mountains as well as the Sweetgrass Hills and Rocky Mountain foothills are at higher elevations, receive higher amounts of precipitation, and have a typic-ustic moisture regime. These areas have significantly more rangeland production than the drier aridic-ustic portions of the MLRA and have enough moisture to produce crops annually rather than just bi-annually, as in the drier areas. Ecological sites in this higher precipitation area are classified as the moist grassland climatic zone.

Classification relationships

NRCS Soil Geography Hierarchy • Land Resource Region: Northern Great Plains • Major Land Resource Area (MLRA) 052 Brown Glaciated Plains • Climate Zone: Dry Grassland National Hierarchical Framework of Ecological Units (Cleland et al. 1997, McNab et al. 2007) • Domain: Dry • Division: Temperate Steppe • Province: Great Plains-Palouse Dry Steppe Province 331 • Section: Northwestern Glaciated Plains 331D • Subsection: Montana Glaciated Plains 331Dh • Landtype association/Landtype phase: N/A National Vegetation Classification Standard (Federal Geographic Data Committee 2008) • Class: Mesomorphic Shrub and Herb Vegetation Class (2) •

Subclass: Temperate and Boreal Grassland and Shrubland Subclass (2.B) • Formation: Temperate Grassland, Meadow, and Shrubland Formation (2.B.2) • Division: Great Plains Grassland and Shrubland Division (2.b.2.Nb) • Macrogroup: Hesperostipa comata – Pascopyrum smithii – Festuca hallii Grassland Macrogroup (2.B.2.Nb.2) • Group: Pascopyrum smithii - Hesperostipa comata - Schizachyrium scoparium - Bouteloua spp. Mixedgrass Prairie Group (2.B.2.Nb.2.c) • Alliance: Pascopyrum smithii – Nassella viridula Northwestern Great Plains Herbaceous Alliance • Association: None identified EPA Ecoregions • Level 1: Great Plains (9) • Level 2: West-Central Semi-Arid Prairies (9.3) • Level 3: Northwestern Glaciated Plains (42) • Level 4: North Central Brown Glaciated Plains (42o) & Glaciated Northern Grasslands (42i)

Ecological site concept

The distinguishing characteristics of this site are depth to gravels being 10 to 20 inches from the soil surface and a clay content of less than 35 percent. Soils are derived from gravelly alluvium. This ecological site occurs on steep terraces and knolls. Slopes usually range from 0 to 15%.

Associated sites

R052XN162MT	Clayey (Cy) 10-14" p.z. soils >20 inches in depth, higher production, and no hardpan, different species composition
R052XN166MT	Overflow (Ov) 10-14" p.z. different position on landscape, site benefits from extra moisture, more production,different species, permanent water table > 42 inches
R052XN179MT	Shallow Clay (SwC) 10-14" p.z. soils 10-20" deep to bedrock' soils are clayey over clayey shale
R052XN178MT	Shallow (Sw) 10-14" p.z. soil depth less than or equal to 20 inches to a restrictive layer; less forage production; parent material variable
R052XN172MT	Dense Clay (DC) 10-14" p.z. has a hard restrictive layerinthesoilator near the surface, salt tolerant plants maybe present but are rarely dominant

Similar sites

R052XC216MT	Shallow to Gravel (SwGr) 10-14" p.z. Shift in plant communities and production, different LRU.
-------------	--

Table 1. Dominant plant species

Tree	Not specified
Shrub	(1) <i>Juniperus horizontalis</i> (2) <i>Yucca glauca</i>

Herbaceous	(1) <i>Pascopyrum smithii</i> (2) <i>Pseudoroegneria spicata</i>
------------	---

Physiographic features

This ecological site occurs on nearly level to steep terraces and knolls. Slopes usually range from 0 to 15%, can occur on slopes greater than 15%. Elevations normally vary from 2,200 to 3,500 feet.

Table 2. Representative physiographic features

Landforms	(1) Outwash terrace (2) Alluvial fan (3) Knoll
Runoff class	Low to high
Flooding frequency	None
Ponding frequency	None
Elevation	670 – 1,070 m
Slope	0 – 20 %
Aspect	W, NW, N, NE, E, SE, S, SW

Table 3. Representative physiographic features (actual ranges)

Runoff class	Not specified
Flooding frequency	Not specified
Ponding frequency	Not specified
Elevation	570 – 1,160 m
Slope	0 – 40 %

Climatic features

A semi-arid, temperate climate characterizes the Glaciated Plains. The predominance of cool season species has evolved to take advantage of the precipitation regime that peaks in late spring-early summer (June). Seventy-five percent of the annual precipitation usually falls as steady, soaking, frontal system rains. Summer rains usually come with thunderstorms. Precipitation is the most important

factor influencing production (Heitschmidt et al 2005). Severe drought occurs on average in two out of every ten years (Cooper, et al., 2001).

Table 4 Representative climatic features

Frost-free period (characteristic range)	70-140 days
Freeze-free period (characteristic range)	110-140 days
Precipitation total (characteristic range)	250-360 mm
Frost-free period (average)	90 days
Freeze-free period (average)	120 days
Precipitation total (average)	310 mm

- (1) CHESTER [USC00241692], Chester, MT
- (2) CHINOOK [USC00241722], Chinook, MT
- (3) HAVRE CITY CO AP [USW00094012], Havre, MT
- (4) MALTA 35 S [USC00245340], Zortman, MT
- (5) SHELBY [USC00247500], Shelby, MT

Influencing water features

This site is not influenced by water from streams.

Wetland description

This site is not influenced by water from wetlands.

Soil features

These soils are moderately deep to very deep. Depth to sand and gravel is typically 10-20 inches. The soils occupy uplands and side slopes of valleys. These are well drained soils that formed in gravelly alluvium. Permeability class is moderate to moderately slow. The surface textures are loam, sandy loam and fine sandy loam. Subsurface textures are usually clay loam and sandy clay loam. The upper 10-12 inches of the soil has less gravel than the next 8-10 inches. Soil ph varies from 6.6 to 9.0. This site is characterized by the following soil components: Beaverell.

Table 5. Representative soil features

Parent material	(1) Alluvium – igneous, metamorphic and sedimentary rock
Surface texture	(1) Gravelly loam (2) Gravelly sandy loam (3) Gravelly fine sandy loam

Family particle size	(1) Fine-loamy (2) Fine-loamy over sandy or sandy-skeletal
Depth to restrictive layer	50 – 150 cm
Soil depth	50 – 200 cm
Surface fragment cover <=3"	0 – 20 %
Surface fragment cover >3"	Not specified
Available water capacity (Depth not specified)	5.08 – 7.62 cm
Calcium carbonate equivalent (Depth not specified)	0 – 10 %
Electrical conductivity (Depth not specified)	Not specified
Sodium adsorption ratio (Depth not specified)	0 – 10
Soil reaction (1:1 water) (Depth not specified)	6.6 – 10
Subsurface fragment volume <=3" (Depth not specified)	20 %
Subsurface fragment volume >3" (Depth not specified)	10 – 20 %

Ecological dynamics

This site developed through time under the influence of climate, geological materials, fire, plants and animals. Research on upland ecological sites consistently shows that precipitation is the principal factor altering productivity (Heitschmidt et al. 2005). The same authors concluded that grazing reduces herbage standing crop, whereas its effects on aboveground net primary production vary with timing of grazing and precipitation events, along with the functional and structural composition of the plant community. Some ecologists believe that these lands may have burned on a natural interval of 10-12 years (Frost 1998). However, environmental characteristics of

this site limit herbage production and subsequent fuel accumulation. Therefore, in comparison to normal upland range sites, the role of natural fire is probably less significant in the development of this site.

The resultant historic climax plant community (HCPC) is the basis for plant community interpretations. The HCPC has been determined by evaluating rangeland relic areas, and other areas protected from excessive disturbance.

The HCPC is comprised of a mixture of cool and warm season grasses, forbs and shrubs. About 85% of the annual production is from grasses and sedges, most of which is produced during the cool season. Forbs and shrubs contribute 10% and 5%, respectively, to total annual production. Total vegetative production averages 1000 lbs/ac in normal years, 700 lbs/ac in "unfavorable" years, and 1200 lbs/ac in "favorable" years.

This site is resistant to disturbance when late-successional plants dominate the site. Departures from the HCPC generally result from management actions, drought, and/or a change in the natural fire regime. The site is considered fragile in the sense that vegetative vigor and composition will rapidly decline with continued adverse impacts. Plant communities that retain a high percentage of late successional species are resilient. With favorable precipitation and/or prescribed grazing treatments these plant communities can return to the HCPC. In contrast, significant succession is unusual within early-seral communities.

State and Transition Diagram

Successional pathways of Shallow to Gravel 10-14" p.z. ecological sites cannot be satisfactorily described using traditional theories of plant succession leading to a single climax community (Briske et al. 2005). As the HCPC regresses to an early seral state, it is theorized that a threshold is crossed somewhere within the mid-seral state. Plant communities occurring below this threshold are in a steady state. Succession back to the HCPC often does not occur within a reasonable length of time, or without a large input of energy.

Two plant communities and the successional pathways that commonly occur within the Reference State (State 1) are shown in the following diagram. The transition from State 1 to State 2, and two plant communities representative of State 2 are also illustrated. Ecological processes are discussed in the plant community descriptions that follow the diagram.

State and transition model

Additional community tables

Table 6. Community 1.1 plant community composition

Group	Common Name	Symbol	Scientific Name	Annual Production ()	Foliar Cover (%)
Grass/Grasslike					
1	Cool-season Bunchgrasses			616-1121	
	bluebunch wheatgrass	PSSP6	<i>Pseudoroegneria spicata</i>	448-785	–
	needle and thread	HECO26	<i>Hesperostipa comata</i>	168-336	–
1	Rhizomatous Wheatgrasses			112-280	
	western wheatgrass	PASM	<i>Pascopyrum smithii</i>	56-140	–
	tufted wheatgrass	ELMA7	<i>Elymus macrourus</i>	56-140	–
3	Warm-season Bunchgrasses			168-392	
	plains muhly	MUCU3	<i>Muhlenbergia cuspidata</i>	112-224	–
	little bluestem	SCSC	<i>Schizachyrium scoparium</i>	56-168	–
4	Miscellaneous Grasses			0-112	
	blue grama	BOGR2	<i>Bouteloua gracilis</i>	0-56	–
	sand dropseed	SPCR	<i>Sporobolus cryptandrus</i>	0-56	–
	plains reedgrass	CAMO	<i>Calamagrostis montanensis</i>	0-56	–
	threadleaf sedge	CAFI	<i>Carex filifolia</i>	0-56	–
	needleleaf sedge	CADU6	<i>Carex duriuscula</i>	0-56	–
	Sandberg bluegrass	POSE	<i>Poa secunda</i>	0-56	–
	prairie Junegrass	KOMA	<i>Koeleria macrantha</i>	0-56	–
	Grass, native	2GN	<i>Grass, native</i>	0-56	–
Forb					
5	Dominant Forb			56-112	
	American vetch	VIAM	<i>Vicia americana</i>	56-112	–
4	Clovers			112-224	

	purple prairie clover	DAPU5	<i>Dalea purpurea</i>	56-112	–
	white prairie clover	DACA7	<i>Dalea candida</i>	56-112	–
7	Miscellaneous Forbs			0-67	
	scarlet globemallow	SPCO	<i>Sphaeralcea coccinea</i>	0-56	–
	aster	ASTER	<i>Aster</i>	0-56	–
	scurfpea	PSORA2	<i>Psoraleidium</i>	0-56	–
	common yarrow	ACMI2	<i>Achillea millefolium</i>	0-56	–
	bastard toadflax	COUM	<i>Comandra umbellata</i>	0-56	–
	milkvetch	ASTRA	<i>Astragalus</i>	0-56	–
	spiny phlox	PHHO	<i>Phlox hoodii</i>	0-56	–
	onion	ALLIU	<i>Allium</i>	0-56	–
	Forb, native	2FN	<i>Forb, native</i>	0-56	–
	lesser spikemoss	SEDE2	<i>Selaginella densa</i>	0-1	–
Shrub/Vine					
8	Miscellaneous Shrubs			0-135	
	creeping juniper	JUHO2	<i>Juniperus horizontalis</i>	0-56	–
	soapweed yucca	YUGL	<i>Yucca glauca</i>	0-56	–
	silver sagebrush	ARCA13	<i>Artemisia cana</i>	0-56	–
	prairie sagewort	ARFR4	<i>Artemisia frigida</i>	0-56	–
	prairie rose	ROAR3	<i>Rosa arkansana</i>	0-56	–
	broom snakeweed	GUSA2	<i>Gutierrezia sarothrae</i>	0-56	–
	Shrub (>.5m)	2SHRUB	<i>Shrub (>.5m)</i>	0-56	–
	plains pricklypear	OPPO	<i>Opuntia polyacantha</i>	0-1	–
	brittle pricklypear	OPFR	<i>Opuntia fragilis</i>	0-1	–

Table 7. Community 1.2 plant community composition

Group	Common Name	Symbol	Scientific Name	Annual Production ()	Foliar Cover (%)
-------	-------------	--------	-----------------	----------------------	------------------

Table 8. Community 2.1 plant community composition

Group	Common Name	Symbol	Scientific Name	Annual Production ()	Foliar Cover (%)
-------	-------------	--------	-----------------	----------------------	------------------

Table 9. Community 2.2 plant community composition

Group	Common Name	Symbol	Scientific Name	Annual Production ()	Foliar Cover (%)
-------	-------------	--------	-----------------	----------------------	------------------

Animal community

Livestock Management The Shallow to Gravel 10-14" p.z. ecological site is fairly productive and is suited for livestock grazing. However, prescribed grazing management is needed. This site is often associated with slopes that may be susceptible to erosion. Species composition is favorable to livestock and is susceptible to heavy stocking and season long grazing. The cool season medium height grasses are generally selectively grazed giving the short grasses a competitive advantage. Grazing during early spring may also result in soil compaction. Any additional factor reducing infiltration and increasing runoff on this site is a management concern. Shorter grazing periods developed in conjunction with adequate periods of deferment to facilitate regrowth, replenish carbohydrate pools, and accumulate litter on the soil surface are recommended. This ecological site, as do most other sites in the northern mixed prairie, has a short grass component. The short grasses usually increase with grazing and decrease with protection or prescribed grazing. However, succession in direct response to a change in grazing pressure is not guaranteed in the Northern Great Plains. Sampling four-year old ungrazed exclosures and grazed areas with 35% utilization, Vogel and Van Dyne (1966) found essentially the same basal cover of grasses, sedges, forbs, litter and bare soil on protected and grazed sites. They concluded that four years was too short of a time for cover to change significantly. Hofmann and Ries (1989) observed similar results following a four-year study in North Dakota. Even after 41 years of exclosure, changes in species composition can be relatively small when the site is in the dry, low production portion of northern mixed prairie (Brand and Goetz, 1986). They concluded that site characteristics limited the development of potential vegetation with the exclusion of grazing, but the potential impacts of prescribed grazing on succession were not discussed. This ecological site is not as productive as the sites evaluated by Vogel and Van Dyne, Hofmann and Ries, or by Brand and Goetz. Therefore, range managers should

recognize the environmental limitations of this site. While a prescribed grazing system is always a good recommendation, it may not guarantee significant succession. Poisonous plants are not normally a problem on this site. Losses that do occur are usually result in livestock being forced to consume the poisonous plants because of an inadequate supply of desirable forage. Wildlife Interpretations The HCPC associated with this ecological site provides diverse and valuable wildlife habitat. This site often occurs as a mosaic with other ecological sites, thus creating "ecotones" that serve as a magnet to attract many species of wildlife. Antelope and mule deer prefer grazing this site because of its position in the landscape. However, its value for thermal and escape cover is limited by the lack of shrubs. This ecological site becomes less valuable for deer and antelope when plant diversity declines with regression. For example, the disappearance of the bluebunch wheatgrass and western wheatgrass shorten the length of the "green forage" season. The increase of blue grama, hoods phlox, etc. is associated with the loss of palatable forbs. These changes also adversely impact foraging opportunities for deer, antelope, upland birds, etc. Because of insufficient vegetative structural diversity, residual grass carry-over and litter cover, the value of Plant Communities in State 2 are greatly reduced for wildlife habitat.

Hydrological functions

Water is the main factor limiting vegetative production on this site. Soil components in this ecological site are normally classed into Hydrologic Group B. When cover conditions are good, these soils have a low runoff potential.

Recreational uses

This site provides hunting opportunities for upland game species. Outdoor enthusiasts may also appreciate the serenity and openness of this site.

Wood products

This site has no significant value for wood products.

Other information

This ecological site is not highly resistant to disturbances. Species diversity is adversely affected by season-long continuous grazing and by heavy stocking. Medium height grasses are replaced by short grasses. The number of structural/functional groups is reduced with regression from the HCPC. The amount of solar energy that is captured and converted to carbohydrates for plant growth is reduced in State 2. A reduction in total vegetative growth results in less potential vegetation that can be transformed into litter. Litter reductions result in less infiltration, and more runoff and soil erosion.

Inventory data references

Data Source Number of Records Sample Period State County SCS-Range-417 0 ECS-1 Modified Double Sampling 12 2002-2004 MT Phillips, Blaine, Valley, Roosevelt, Daniels Ross, R. L. and H. E. Hunter. 1976. Climax vegetation of Montana. USDA Soil Conservation Service. Bozeman, MT. USDA-SCS-MT 1981 Technical Range Site Description

Other references

Brand, M.D. and H. Goetz. 1986. Vegetation of exclosures in Southwestern North Dakota. J. Range Manage. 39:434-437.
Branson, F. A., and R. F. Miller. 1981. Effects of increased precipitation and grazing management on northeastern Montana rangelands. J. Range Manage. 34(1): 3-11.
Briske, D. D., S. D. Fuhlendorf, and F. E. Smeins, 2005. State-and-transition models, thresholds, and rangeland health: a synthesis of ecological concepts and perspectives. Rangeland Ecol. Manage 58:1-10.
Frost, C. C. 1998. Presettlement fire frequency regimes of the United States: a first approximation. Pages 70-81. in Teresa L. Pruden and Leonard A. Brennan (eds.). Fire in ecosystem management: shifting paradigm from suppression to prescription. Tall Timbers Fire Ecology Conference Proceedings. No. 20. Tall Timbers Research Station, Tallahassee, FL.
Heitschmidt, R. K., K. D. Klement, and M. R. Haferkamp. 2005. Interactive effects of drought and grazing on Northern Great Plains rangelands. Rangeland Ecol. Manage. 58:11-19.
Hofmann, L. and R.E. Ries. 1989. Animal performance and plant production from continuously grazed cool-season reclaimed and native pastures. J. Range Manage. 42:248-251.
U.S. Department of Interior and U.S. Department of Agriculture. 2000. Interpreting indicators of rangeland health. Tech. Ref. 1734-6.
Vogel, W.G. and G.M. Van Dyne. 1966. Vegetation responses to grazing management on a foothill sheep range. J. Range Manage. 19:80-85.

Contributors

Kirt Walstad

Approval

Kirt Walstad, 4/14/2025

Acknowledgments

Site Description Revisions The 2005 Shallow to Gravel 10-14" p.z. ecological site description replaces earlier dated versions of the Shallow to Gravel 10-14" p.z. description in Rangeland Resource Unit 52XN. This 2005 revision incorporates the State and Transition Model theory, additional data on site productivity, and an improved understanding of many rangeland health indicators. Site Description Approval This ecological site description is approved with the understanding that it is no more than another step in our continual effort to update the NRCS technical guide. In order to facilitate the process, NRCS field personnel are encouraged to forward existing information and/or new data that can be used to improve the utility of this site description. Please forward the information and data to the State Rangeland Management Specialist. Authors Date Approval Date Dr. John Lacey 02/28/2005 Loretta J. Metz 03/19/2005 Maxine Rasmussen, Area RMS, Glasgow, MT Jon Siddoway, Area RMS, Great Falls, MT Rick Bandy, Area RSS, Great Falls, MT Greg Snell, Area RSS, Glasgow, MT

Rangeland health reference sheet

Interpreting Indicators of Rangeland Health is a qualitative assessment protocol used to determine ecosystem condition based on benchmark characteristics described in the Reference Sheet. A suite of 17 (or more) indicators are typically considered in an assessment. The ecological site(s) representative of an assessment location must be known prior to applying the protocol and must be verified based on soils and climate. Current plant community cannot be used to identify the ecological site.

Author(s)/participant(s)	Loretta Metz, Maxine Rasmussen, Jon Siddoway
Contact for lead author	Area Rangeland Management Specialist, Glasgow Area Office, MT Reference site used? No
Date	05/04/2005
Approved by	
Approval date	
Composition (Indicators 10 and 12) based on	Annual Production

Indicators

- 1. Number and extent of rills:** Rills should not be present in HCPC. On slopes > 15%, careful examination will yield slight evidence of rills following a rainfall event in HCPC. On slopes > 15%, if in Plant community A, careful examination will yield slight evidence of rills regardless of precipitation event. On slopes > 15%, in HCPC and in plant community A, rill would be less than ½ inch deep, linear, but short in length
-

2. **Presence of water flow patterns:** Careful examination will yield slight evidence of water flow patterns following a rainfall event in HCPC on slopes > 15%. On slopes > 15%, if in Plant community A, careful examination will yield slight evidence of water flow patterns regardless of precipitation events.
-
3. **Number and height of erosional pedestals or terracettes:** Pedestals or terracettes would essentially be nonexistent in HCPC. On slopes > 8%, if in Plant community A, careful examination will yield occasional pedestals and terracettes approximately ¼ inch above the soil surface.
-
4. **Bare ground from Ecological Site Description or other studies (rock, litter, lichen, moss, plant canopy are not bare ground):** Up to 15% of the soil surface is bare ground in HCPC. In plant community A, 15-20% bare ground is possible
-
5. **Number of gullies and erosion associated with gullies:** None.
-
6. **Extent of wind scoured, blowouts and/or depositional areas:** None.
-
7. **Amount of litter movement (describe size and distance expected to travel):** Litter movement is not expected in HCPC. On slopes > 8%, in Plant community A, litter, both fine and coarse, movement is readily observable.
-
8. **Soil surface (top few mm) resistance to erosion (stability values are averages - most sites will show a range of values):** Stability class anticipated to be 1 to 4, depending on soil surface texture.
-
9. **Soil surface structure and SOM content (include type of structure and A-horizon color and thickness):** The surface layer is 0.5" to 4" thick and typically gravelly sandy loam surface textures. Surface color ranges brown and very dark brown. Soil organic matter ranges steadily between 1-3%.
-
10. **Effect of community phase composition (relative proportion of different functional groups) and spatial distribution on infiltration and runoff:** In HCPC, 75-85% plant canopy and 20% basal cover with small gaps between plants should reduce raindrop impact and slow overland flow, providing increased time for infiltration to occur. Healthy, deep rooted native grasses enhance infiltration and reduce runoff. Infiltration rate is moderate to moderately slow. If in plant community A, 70-80% plant canopy and 15% basal cover with small gaps between plants will still reduce raindrop impact and decrease overland flow.
-
11. **Presence and thickness of compaction layer (usually none; describe soil profile features which may be mistaken for compaction on this site):** No compaction layer or soil surface crusting should be evident in any of the State 1 plant communities. Unconsolidated sand or gravel layer begins at 10-20 inches.
-

12. **Functional/Structural Groups** (list in order of descending dominance by above-ground annual-production or live foliar cover using symbols: >>, >, = to indicate much greater than, greater than, and equal to):

Dominant:

Sub-dominant:

Other:

Additional: Tall and mid stature cool season bunch grasses = mid stature warm season bunch grasses > mid-stature cool season rhizomatous grasses > forbs > shrubs. Plant community A: Mid-stature cool season bunch grasses > mid-stature cool season rhizomatous grasses > short warm season rhizomatous grasses > forbs > shrubs.

13. **Amount of plant mortality and decadence** (include which functional groups are expected to show mortality or decadence): Plant mortality and decadence very low in HCPC and Plant community A. In periods of drought, shrubs would exhibit decadence in the state 1 reference communities.
-

14. **Average percent litter cover (%) and depth (in):**
-

15. **Expected annual annual-production** (this is TOTAL above-ground annual-production, not just forage annual-production): 700 - 1200 #/acre from Plant community A to HCPC.
-

16. **Potential invasive (including noxious) species** (native and non-native). List species which BOTH characterize degraded states and have the potential to become a dominant or co-dominant species on the ecological site if their future establishment and growth is not actively controlled by management interventions. Species that become dominant for only one to several years (e.g., short-term response to drought or wildfire) are not invasive plants. Note that unlike other indicators, we are describing what is NOT expected in the reference state for the ecological site: Needle and thread, threadleaf sedge, plains prickly pear, blue grama, Hood's phlox, hairy goldenaster, dense clubmoss and fringed sagewort
-

17. **Perennial plant reproductive capability:** All species are capable of reproducing in HCPC. In Plant community A, plant seedlings will be weighed in favor of marginal and undesirable species. Replacement of desirable species will be very few.
-