

# Ecological site FX053A99X150

## Subirrigated (Sb)

Last updated: 4/25/2025  
Accessed: 04/22/2026

---

### General information

**Provisional.** A provisional ecological site description has undergone quality control and quality assurance review. It contains a working state and transition model and enough information to identify the ecological site.

### MLRA notes

Major Land Resource Area (MLRA): 053A–Northern Dark Brown Glaciated Plains

The Northern Dark Brown Glaciated Plains, MLRA 53A, is a large, agriculturally and ecologically significant area. It consists of approximately 6.1 million acres and stretches 140 miles from east to west and 120 miles from north to south, encompassing portions of 8 counties in northeastern Montana and northwestern North Dakota. This region represents part of the southern edge of the Laurentide Ice Sheet during maximum glaciation. It is one of the driest and westernmost areas within the vast network of glacially derived prairie pothole landforms of the Northern Great Plains and falls roughly between the Missouri Coteau to the east and the Brown Glaciated Plains to the west. Elevation ranges from 1,800 feet (550 meters) to 3,300 feet (1,005 meters). Soils are primarily Mollisols, but Inceptisols and Entisols are also common. Till from continental glaciation is the predominant parent material, but alluvium and bedrock are also common. Till deposits are typically less than 50 feet thick (Soller, 2001). Underlying the till is sedimentary bedrock largely consisting of Cretaceous shale, sandstone, and mudstone (Vuke et al., 2007). The bedrock is commonly exposed on hillslopes, particularly along drainageways. Significant alluvial deposits occur in glacial outwash channels and along major drainages, including portions of the Missouri, Poplar, and Big Muddy Rivers. Large eolian deposits of sand occur in the vicinity of the ancestral Missouri River channel east of Medicine Lake (Fullerton et al., 2004). The northwestern portion of the MLRA contains a large unglaciated area containing paleoterraces and large deposits of sand and gravel known as the Flaxville gravel. Much of this MLRA was glaciated towards the end of the Wisconsin age, and the maximum glacial extent occurred approximately 20,000 years ago (Fullerton and Colton, 1986; Fullerton et al., 2004). Subsequent erosion from major stream and river systems has created numerous drainageways throughout much of the MLRA. The result is a geologically young landscape that is predominantly a dissected till plain interspersed with alluvial deposits and dominated by soils in the Mollisol and Inceptisol orders. Much of this area is typic ustic, making these soils very productive and generally well suited to production agriculture. Dryland farming is the predominant land use, and approximately 50 percent of the land area is used for cultivated crops. Winter, spring, and durum varieties of wheat are the major crops, with over 48 million bushels produced annually (USDA-NASS, 2017). Areas of rangeland typically are on steep hillslopes along drainages. The rangeland is mostly native mixedgrass prairie similar the *Stipa-Agropyron*, *Stipa-Bouteloua-Agropyron*, and *Stipa-Bouteloua* faciations (Coupland, 1950; 1961). Cool-season grasses dominate and include rhizomatous wheatgrasses, needle and thread, western porcupine grass, and green needlegrass. Woody species are generally rare; however, many of the steeper drainages support stands of trees and shrubs such as green ash and chokecherry. Seasonally ponded, prairie pothole wetlands may occur throughout the MLRA, but the greatest concentrations are in the east and northeast where receding glaciers stagnated and formed disintegration moraines with hummocky topography and numerous areas of poorly drained soils.

### Classification relationships

NRCS Soil Geography Hierarchy • Land Resource Region: Northern Great Plains • Major Land Resource Area (MLRA): 053A Northern Dark Brown Glaciated Plains National Hierarchical Framework of Ecological Units (Cleland et al., 1997; McNab et al., 2007) • Domain: Dry • Division: Temperate Steppe • Province: Great Plains-Palouse Dry Steppe Province 331 • Section: Glaciated Northern Grasslands Section 331L • Subsection: Glaciated Northern Grasslands Subsection 331La • Landtype association/Landtype phase: N/A National Vegetation Classification Standard (Federal Geographic Data Committee, 2008) • Class: Mesomorphic Shrub and Herb Vegetation Class (2) • Subclass: Shrub and Herb Wetland Subclass (2.C) • Formation: Temperate to Polar Freshwater Marsh, Wet Meadow and Shrubland Formation (2.C.4) • Division: Eastern North American Temperate and Boreal Freshwater Marsh, Wet Meadow and Shrubland Division (2.C.4.Nd) • Macrogroup: *Spartina pectinata* - *Typha* spp. - *Schoenoplectus* spp. Great Plains Marsh, Wet Meadow, Shrubland and Playa Macrogroup (2.C.4.Nd.5) • Group: *Spartina pectinata* - *Calamagrostis stricta* - *Carex* spp. Great Plains Wet Prairie, Wet Meadow and Seepage Fen Group (2.C.4.Nd.5.b) EPA Ecoregions • Level 1: Great Plains (9) • Level 2: West-Central Semi-Arid Prairies (9.3) • Level 3: Northwestern Glaciated Plains (42) • Level 4: Glaciated Dark Brown Prairie (42i) Glaciated Northern Grasslands (42j)

### Ecological site concept

Subirrigated is an ecological site of limited extent occurring on floodplains and stream terraces. The distinguishing characteristics of this site are that it receives additional moisture from groundwater, and a seasonal water table occurs 24 to 40 inches below the soil surface. Soils for this ecological site are typically very deep (more than 60 inches), somewhat poorly drained, and derived from alluvium. Characteristic vegetation is sedges (*Carex* spp.), slimstem reedgrass (*Calamagrostis stricta*), and prairie cordgrass (*Spartina pectinata*).

### Associated sites

<b>FX053A99X060</b>	<p><b>Overflow (Ov)</b></p> <p>This site is adjacent to the Subirrigated ecological site, typically on higher terraces where ground water is greater than 40 inches below the surface and the primary moisture source is surface water.</p>
<b>FX053A99X084</b>	<p><b>Slough (SI)</b></p> <p>This site is adjacent to the Subirrigated ecological site, typically in oxbows or channels where flooding is very frequent, the water table is shallow and persistent, and frequent ponding occurs.</p>
<b>FX053A99X713</b>	<p><b>Saline Lowland (SLL)</b></p> <p>This site is adjacent to the Subirrigated ecological site in similar landscape positions but in areas where salts have accumulated due to geology, hydrology, or soil properties.</p>
<b>FX053A99X061</b>	<p><b>Riparian Woodland (RW)</b></p> <p>This site is adjacent to the Subirrigated ecological site, typically on similar landscape positions, but where riparian woody plants are dominant.</p>

### Similar sites

<b>FX053A99X084</b>	<p><b>Slough (SI)</b></p> <p>This site differs from the Subirrigated ecological site in that depth to a water table is less than 24 inches and the site receives frequent long duration ponding. It is located in oxbows, old channels, or depressions on floodplains and is more productive. Vegetation is dominated by hydrophytes such as bulrush and cattail.</p>
<b>FX053A99X713</b>	<p><b>Saline Lowland (SLL)</b></p> <p>This site differs from the Subirrigated ecological site in that soils are saline, sodic, or saline-sodic (EC ? 4 or SAR ? 13). It supports more sodium-tolerant vegetation and is less productive.</p>
<b>FX053A99X061</b>	<p><b>Riparian Woodland (RW)</b></p> <p>This site differs from the Subirrigated ecological site in that it is dominated by riparian woody species. Shrubs and trees dominate the site in terms of cover and production.</p>

Table 1. Dominant plant species

Tree	Not specified
Shrub	Not specified

Herbaceous	(1) <i>Carex</i> (2) <i>Calamagrostis stricta</i>
------------	--

## Legacy ID

R053AY719MT

## Physiographic features

This ecological site occurs on alluvial fans and floodplain steps of perennial or intermittent streams, near springs or seeps, or other areas that have a seasonal water table 24 to 40 inches below the soil surface. Slopes typically range from 0 to 2 percent. This site occurs on all aspects.

Table 2. Representative physiographic features

Landforms	(1) River valley > Flood-plain step (2) River valley > Alluvial fan (3) Till plain > Drainageway
Flooding frequency	None to rare
Ponding frequency	None
Elevation	550 – 1,010 m
Slope	0 %
Water table depth	60 – 100 cm
Aspect	Aspect is not a significant factor

## Climatic features

The Northern Dark Brown Glaciated Plains is a semi-arid region with a temperate continental climate that is characterized by frigid winters and warm to hot summers (Coupland, 1958; Richardson and Hanson, 1977; Heidel et al., 2000). The majority of precipitation occurs as steady, soaking, frontal system rains in late spring to early summer. Summer rainfall comes mainly from convection thunderstorms that typically deliver scattered amounts of rain in intense bursts. These storms may be accompanied by damaging winds and large-diameter hail and result in flash flooding along low-order streams. Approximately 80 percent of the annual precipitation occurs during the growing season. June is the wettest month, followed by July and May (Richardson and Hanson, 1977; Heidel et al., 2000). Average annual precipitation ranges from 11 inches (280 mm) near Richey, Montana, to 15 inches (380 mm) in the Little Muddy drainage near Williston, North Dakota, but precipitation varies greatly from year to year. On average, severe drought and very wet years occur with the same frequency, which is 1 out of 10 years (Coupland, 1958; Heidel et al., 2000). Extreme climatic variations, especially droughts, have the greatest influence on species cover and production (Coupland, 1958, 1961; Biondini et al., 1998). The frost-free period for this ecological site ranges from 90 to 130 days, and the freeze-free period ranges from 115 to 155 days.

Table 3 Representative climatic features

Frost-free period (characteristic range)	90-130 days
--	-------------

Freeze-free period (characteristic range)	120-160 days
Precipitation total (characteristic range)	280-380 mm
Frost-free period (average)	110 days
Freeze-free period (average)	140 days
Precipitation total (average)	330 mm

- (1) BREDETTE [USC00241088], Poplar, MT
- (2) CULBERTSON [USC00242122], Culbertson, MT
- (3) OPHEIM 10 N [USC00246236], Opheim, MT
- (4) OPHEIM 12 SSE [USC00246238], Opheim, MT
- (5) PLENTYWOOD [USC00246586], Plentywood, MT
- (6) SCOBNEY 4 NW [USC00247425], Scobey, MT
- (7) SIDNEY [USC00247560], Sidney, MT
- (8) VIDA 6 NE [USC00248569], Vida, MT
- (9) WILLISTON SLOULIN INTL AP [USW00094014], Williston, ND

### Influencing water features

This is a floodplain or drainageway site that receives additional moisture from groundwater and occasionally stream overflow. When on floodplains, the site may be flooded for brief durations during major flood events. A seasonal groundwater table is present between 24 and 40 inches below the soil surface, particularly during spring.

### Wetland description

Palustrine Emergent

### Soil features

Soils for this ecological site are typically very deep (more than 60 inches), somewhat poorly drained, and derived from alluvium. All soils in this site concept are endosaturated, meaning that they receive additional moisture from groundwater. A seasonal water table is present at a depth of 24 to 40 inches below the soil surface. These soils have an aquic moisture regime and a frigid soil temperature regime (Soil Survey Staff, 2014).

Surface horizon textures in this site are typically loam, silt loam, or silty clay loam. The underlying horizons are typically comprised of stratified alluvial deposits, characterized by many thin layers of sediment deposited by past flood events. In the upper 20 inches, electrical conductivity is less than 4 and the sodium absorption ratio is less than 13. Calcium carbonate equivalent is typically less than 15 percent throughout the soil profile. Soil pH classes are neutral to slightly alkaline in the surface horizon and slightly alkaline to moderately alkaline in the subsurface horizons. Typically, the upper 20 inches of soil does not contain coarse fragments.

Table 4. Representative soil features

Parent material	(1) Alluvium – igneous, metamorphic and sedimentary rock
Surface texture	(1) Loam (2) Silt loam (3) Silty clay loam

Drainage class	Somewhat poorly drained
Soil depth	150 – 180 cm
Electrical conductivity (0-50.8cm)	Not specified
Sodium adsorption ratio (0-50.8cm)	0 – 10

### Ecological dynamics

The information in this ecological site description, including the state-and-transition model (STM), was developed based on historical data, current field data, professional experience, and a review of the scientific literature. As a result, all possible scenarios or plant species may not be included. Key indicator plant species, disturbances, and ecological processes are described to inform land management decisions.

The Subirrigated provisional ecological site in MLRA 53A consists of five vegetative states: The Historic Reference state (1), the Current Potential state (2), the Altered state (3), the Invaded state (4), and the Cropland state (5). Plant communities associated with the Subirrigated ecological site evolved under the combined influences of climate, fire, grazing, and hydrology. Extreme climatic variability results in frequent droughts, which can have great influence on the relative contribution of species cover and production (Coupland, 1958, 1961; Biondini et al., 1998).

The historic ecosystem experienced periodic lightning-caused fires with estimated fire return intervals of 6 to 25 years (Bragg, 1995). Historically, Native Americans also set periodic fires. The majority of lightning-caused fires occurred in July and August, whereas Native Americans typically set fires during spring and fall to correspond with the movement of bison (Higgins, 1986). Generally, fires were less frequent on the Subirrigated ecological site than on adjacent drier sites, however, early reports indicate that fires did occur in wetlands (Higgins, 1986). The Subirrigated ecological site is resilient to fire and the most significant effects of fire are most likely removing excess litter accumulations and triggering resprouting and reseeding of cattail and hardstem bulrush (Esser, 1995; Gucker, 2008).

Native grazers also shaped these plant communities. American bison (*Bison bison*) were the dominant historic grazer, but pronghorn (*Antilocapra americana*), elk (*Cervus canadensis*), and deer (*Odocoileus* spp.) were also common. Grasshoppers and periodic outbreaks of Rocky Mountain locusts (*Melanoplus spretus*) also played an important role in the ecology of these communities (Lockwood, 2004).

Hydrology is a crucial dynamic on this site. Depth and duration of the seasonal water table strongly influences species composition on this ecological site. Hydrologic alterations that modify the depth and the persistence of the seasonal water table may have a significant effect on species composition and production. In some cases, salinization may occur. On portions of this site the hydrology has been significantly altered by irrigation, major dams, and diversions. The implications of this alteration have not been fully studied and require further investigation.

Improper grazing of this site can result in a reduction in the cover of the palatable sedges and cool-season midgrasses (Hansen et al., 1995). Tall, warm-season rhizomatous grasses may sustain trampling damage. Improper grazing practices include any practices that do not allow sufficient opportunity for plants to physiologically recover from a grazing event or multiple grazing events within a given year, or that do not provide adequate cover to prevent soil erosion over time. These practices may include, but are not limited to, overstocking, continuous grazing, and/or inadequate seasonal rotation moves over multiple years. Over time species diversity, particularly of sedges, can be significantly reduced by improper grazing. Periods of drought can also reduce sedges and rhizomatous grasses. Further degradation of the site due to improper grazing can result in low vigor of rhizomatous grasses and dominance of unpalatable forbs and rushes. On some sites, sedges can be reduced to a single, unpalatable species (Hansen et al., 1995). Unpalatable forbs may also be common.

Most, if not all, extant examples of this site have some degree of invasion by non-native species. This site is highly susceptible to invasion by non-native species. Perennial grasses such as non-native bluegrasses (*Poa* spp.), smooth brome (*Bromus inermis*), and reed canarygrass (*Phalaris arundinacea*) are the most common invasive species. These species are widespread throughout the Northern Great Plains and can invade relatively undisturbed grasslands (DeKeyser et al., 2013; Grant et al., 2009; Toledo et al., 2014). In most cases native ecological function is relatively intact, but in some cases non-native grasses will displace native species and dominate the ecological functions of the site. Noxious weeds are also a major concern on this site. Leafy spurge (*Euphorbia esula*), Canada thistle

(*Cirsium arvense*), and Russian knapweed (*Acroptilon repens*), also known as hardheads, are common on this site and capable of displacing native species.

The Subirrigated ecological site is often considered prime farmland and large portions of this site have been converted to cropland, mostly for perennial hay. Common crop species include alfalfa (*Medicago sativa*), orchardgrass (*Dactylis glomerata*), creeping foxtail (*Alopecurus arundinaceus*), and grass/alfalfa mixes. Annual crops such as wheat and barley are occasionally planted as part of a rotation or when renovating hay fields. Sometimes irrigation is applied with flood irrigation being most common. Water is typically diverted from nearby streams and delivered to fields via canals. Extensive irrigation systems are in place on many parts of the major river drainages. Irrigated cropland is extremely valuable in the region, and once the site is converted it is unlikely to be taken out of production.

The state-and-transition model (STM) diagram and legend suggests possible pathways that plant communities on this site may follow as a result of a given set of ecological processes and management. The site may also support vegetative states not displayed in the STM diagram. Landowners and land managers should seek guidance from local professionals before prescribing a particular management or treatment scenario. Plant community responses vary across this MLRA due to variability in weather, soils, and aspect. The reference community phase may not necessarily be the management goal. The lists of plant species and species composition values are provisional and are not intended to cover the full range of conditions, species, and responses for the site. Species composition by dry weight is provided when available and is considered provisional based on the sources identified in the narratives associated with each community phase.

## State and transition model

### Additional community tables

Table 5. Community 1.1 plant community composition

Group	Common Name	Symbol	Scientific Name	Annual Production ()	Foliar Cover (%)
-------	-------------	--------	-----------------	----------------------	------------------

Table 6. Community 2.1 plant community composition

Group	Common Name	Symbol	Scientific Name	Annual Production ()	Foliar Cover (%)
-------	-------------	--------	-----------------	----------------------	------------------

Table 7. Community 2.2 plant community composition

Group	Common Name	Symbol	Scientific Name	Annual Production ()	Foliar Cover (%)
-------	-------------	--------	-----------------	----------------------	------------------

Table 8. Community 3.1 plant community composition

Group	Common Name	Symbol	Scientific Name	Annual Production ()	Foliar Cover (%)
-------	-------------	--------	-----------------	----------------------	------------------

Table 9. Community 4.1 plant community composition

Group	Common Name	Symbol	Scientific Name	Annual Production ()	Foliar Cover (%)
-------	-------------	--------	-----------------	----------------------	------------------

Table 10. Community 5.1 plant community composition

Group	Common Name	Symbol	Scientific Name	Annual Production ()	Foliar Cover (%)
-------	-------------	--------	-----------------	----------------------	------------------

## Inventory data references

Only one field plot was available for this site, but data was collected from this plot for six consecutive years. These data, in conjunction with a review of the scientific literature and professional experience, were used to approximate the plant communities for this provisional ecological site. Information for the state-and-transition model was obtained from the same sources. All community phases are considered provisional based on these plots and the sources identified in this ecological site description.

## Other references

Anderson, R.C. 2006. Evolution and origin of the central grassland of North America: Climate, fire, and mammalian grazers. *Journal of Torrey Botanical Society* 133:626-647.

Biondini, M.E., and L. Manske. 1996. Grazing frequency and ecosystem processes in a northern mixed prairie, USA. *Ecological Applications* 6:239-256.

Biondini, M.E., B.D. Patton, and P.E. Nyren. 1998. Grazing intensity and ecosystem processes in a northern mixed-grass prairie, USA. *Ecological Applications* 8:469-479.

Bragg, T.B. 1995. The physical environment of the Great Plains grasslands. In: A. Joern and K.H. Keeler (eds.) *The Changing Prairie*,

Oxford University Press, Oxford, pp. 49–81.

Cleland, D.T., et al. 1997. National hierarchical framework of ecological units. In: M.S. Boyce and A. Haney (eds.) *Ecosystem Management Applications for Sustainable Forest and Wildlife Resources*, Yale University Press, New Haven, CT.

Cooper, S.V. and W.M. Jones. 2003. Site descriptions of high-quality wetlands derived from existing literature sources. Report to the Montana Department of Environmental Quality. Montana Natural Heritage Program, Helena.

Coupland, R.T. 1950. Ecology of the mixed prairie of Canada. *Ecological Monographs* 20:271-315.

Coupland, R.T. 1958. The effects of fluctuations in weather upon the grasslands of the Great Plains. *Botanical Review* 24:273-317.

Coupland, R.T. 1961. A reconsideration of grassland classification in the Northern Great Plains of North America. *Journal of Ecology* 49:135-167.

Coupland, R.T., and R.E. Johnson. 1965. Rooting characteristics of native grassland species in Saskatchewan. *Journal of Ecology* 53:475-507.

Cowardin, L.M., V. Carter, F.C. Golet, and E.T. LaRoe. 1979. Classification of wetlands and deepwater habitats of the United States. US Fish and Wildlife Service FWS/OBS, 79(31), 131.

Crowe, E. and G. Kudray. 2003. Wetland Assessment of the Whitewater Watershed. Report to U.S. Bureau of Land Management, Malta Field Office. Montana Natural Heritage Program, Helena.

DeKeyser, E.S., M. Meehan, G. Clambey, and K. Krabbenhoft. 2013. Cool season invasive grasses in northern Great Plains natural areas. *Natural Areas Journal* 33:81-90.

DeKeyser, S., G. Clambey, K. Krabbenhoft, and J. Ostendorf. 2009. Are changes in species composition on central North Dakota rangelands due to non-use management? *Rangelands* 31:16-19.

Derner, J.D., and R.H. Hart. 2007. Grazing-induced modifications to peak standing crop in northern mixed-grass prairie. *Rangeland Ecology and Management* 60:270-276.

Dix, R.L. 1960. The effects of burning on the mulch structure and species composition of grasslands in western North Dakota. *Ecology* 41:49-56.

Federal Geographic Data Committee. 2008. The national vegetation classification standard, version 2. FGDC Vegetation Subcommittee, FGDC-STD-005-2008 (Version 2), p. 126.

Fullerton, D.S., and R.B. Colton. 1986. Stratigraphy and correlation of the glacial deposits on the Montana Plains. U.S. Geological Survey.

Fullerton, D.S., R.B. Colton, C.A. Bush, and A.W. Straub. 2004. Map showing spatial and temporal relations of mountain and continental glaciations on the northern plains, primarily in northern Montana and northwestern North Dakota. U.S. Geologic Survey pamphlet accompanying Scientific Investigations Map 2843.

Fullerton, D.S., R.B. Colton, and C.A. Bush. 2013. Quaternary geologic map of the Shelby 1° x 2° quadrangle, Montana: U.S. Geological Survey Open-File Report 2012–1170, scale 1:250,000.

Galatowitsch, S.M. and A.G. Van der Valk. 1996. The vegetation of restored and natural prairie wetlands. *Ecological Applications*. 6:1 pp.102-112.

Gilbert, M.C., P.M. Whited, E.J. Clairain Jr., and R.D. Smith. 2006. A regional guidebook for applying the hydrogeomorphic approach to assessing wetland functions of prairie potholes. U.S. Army Corps of Engineers Final Report, Washington, DC.

Grant, T.A., B. Flanders-Wanner, T.L. Shaffer, R.K. Murphy, and G.A. Knutsen. 2009. An emerging crisis across northern prairie refuges: Prevalence of invasive plants and a plan for adaptive management. *Ecological Restoration* 27:58-65.

Hansen, P.L., et al. 1995. Classification and management of Montana's riparian and wetland sites. University of Montana, Montana Forest and Conservation Experiment Station, Miscellaneous Publication No. 54.

Heidel, B., S.V. Cooper, and C. Jean. 2000. Plant species of special concern and plant associations of Sheridan County, Montana. Report to U.S. Fish and Wildlife Service. Montana Natural Heritage Program, Helena.

Heitschmidt, R.K., and L.T. Vermeire. 2005. An ecological and economic risk avoidance drought management decision support system.

In: J.A. Milne (ed.) *Pastoral Systems in Marginal Environments*, XXth International Grasslands Congress, July 2005, p. 178.

Herrick, J.E., J.W. Van Zee, K.M. Havstad, L.M. Burkett, and W.G. Whitford. 2009. *Monitoring manual for grassland, shrubland and savanna ecosystems*. U.S. Department of Agriculture, Agricultural Research Service, Jornada Experimental Range, Las Cruces, NM.

Higgins, K.F. 1986. *Interpretation and compendium of historical fire accounts in the Northern Great Plains*. U.S. Fish and Wildlife Service Resource Publication 161.

Jones, W.M. 2004. *Using vegetation to assess wetland condition: a multimetric approach for temporarily and seasonally flooded depression wetlands and herbaceous-dominated intermittent and ephemeral riverine wetlands in the northwestern glaciated plains ecoregion, Montana*. Report to the Montana Department of Environmental Quality and the U.S. Environmental Protection Agency. Montana Natural Heritage Program, Helena.

Knopf, F.L. 1996. *Prairie legacies—birds*. In: F.B. Samson and F.L. Knopf (eds.) *Prairie Conservation: Preserving North America's Most Endangered Ecosystem*, Island Press, Washington, DC, pp. 135-148.

Knopf, F.L., and F.B. Samson. 1997. *Conservation of grassland vertebrates*. In: F.B. Samson and F.L. Knopf (eds.) *Ecology and Conservation of Great Plains Vertebrates: Ecological Studies 125*, Springer-Verlag, New York, NY, pp. 273-289.

Laycock, W.A. 1991. *Stable states and thresholds of range condition on North American rangelands*. *Journal of Range Management* 44:427-433.

Lesica, P. and P. Husby. 2006. *Field Guide to Montana's Wetland Vascular Plants*. Montana Wetlands Trust. Helena.

Lockwood, J.A. 2004. *Locust: The devastating rise and mysterious disappearance of the insect that shaped the American frontier*. Basic Books, New York, NY.

McNab, W.H., et al. 2007. *Description of ecological subregions: Sections of the conterminous United States [CD-ROM]*. USDA Forest Service, General Technical Report WO-76B.

Montana State College. 1949. *Similar vegetative rangeland types in Montana*. Montana State College, Agricultural Experiment Station.

Murphy, R.K., and T.A. Grant. 2005. *Land management history and floristics in mixed-grass prairie, North Dakota, USA*. *Natural Areas Journal* 25:351-358.

Nesser, J.A., G.L. Ford, C.L. Maynard, and D.S. Page-Dumroese. 1997. *Ecological units of the Northern Region: Subsections*. USDA Forest Service, Intermountain Research Station, General Technical Report INT-GTR-369.

Richardson, R.E., and L.T. Hanson. 1977. *Soil survey of Sheridan County, Montana*. USDA Soil Conservation Service, Bozeman, MT.

Rowe, J.S. 1969. *Lightning fires in Saskatchewan grassland*. *Canadian Field Naturalist* 83:317-327.

Salo, E.D., et al. 2004. *Grazing intensity effects on vegetation, livestock and non-game birds in North Dakota mixed-grass prairie*. *Proceedings of the 19th North American Prairie Conference*, Madison, WI.

Schoeneberger, P.J., D.A. Wysocki, E.C. Benham, and Soil Survey Staff. 2012. *Field book for describing and sampling soils*. Version 3.0. USDA Natural Resources Conservation Service, National Soil Survey Center, Lincoln, NE.

Soil Survey Staff. 2014. *Keys to soil taxonomy*, 12th edition. USDA Natural Resources Conservation Service.

Soller, D.R. 2001. *Map showing the thickness and character of Quaternary sediments in the glaciated United States east of the Rocky Mountains*. U.S. Geological Survey Miscellaneous Investigations Series I-1970-E, scale 1:3,500,000.

Toledo, D., M. Sanderson, K. Spaeth, J. Hendrickson, and J. Printz. 2014. *Extent of Kentucky bluegrass and its effect on native plant species diversity and ecosystem services in the Northern Great Plains of the United States*. *Invasive Plant Science and Management* 7:543-552.

U.S. Department of Agriculture, National Agricultural Statistics Service. 2017. *Montana Annual Bulletin, Volume LIV, Issue 1095-7278*. [https://www.nass.usda.gov/Statistics\\_by\\_State/Montana/Publications/Annual\\_Statistical\\_Bulletin/2017/Montana\\_Annual\\_Bulletin\\_2017.pdf](https://www.nass.usda.gov/Statistics_by_State/Montana/Publications/Annual_Statistical_Bulletin/2017/Montana_Annual_Bulletin_2017.pdf) (Accessed 14 February 2017).

U.S. Department of Agriculture, Natural Resources Conservation Service. *Glossary of landform and geologic terms*. National Soil Survey Handbook, Title 430-VI, Part 629.02c. [http://www.nrcs.usda.gov/wps/portal/nrcs/detail/soils/ref/?cid=nrcs142p2\\_054242](http://www.nrcs.usda.gov/wps/portal/nrcs/detail/soils/ref/?cid=nrcs142p2_054242) (Accessed 13 April 2016).

Vance, L., S. Owen, and J. Horton. 2013. Literature review: Hydrology-ecology relationships in Montana prairie wetlands and intermittent/ephemeral streams. Report to the Cadmus Group and the U.S. Environmental Protection Agency. Montana Natural Heritage Program, Helena.

Vuke, S.M., K.W. Porter, J.D. Lonn, and D.A. Lopez. 2007. Geologic Map of Montana - information booklet: Montana Bureau of Mines and Geology Geologic Map 62-D.

Wilson, S.D., and J.M. Shay. 1990. Competition, fire, and nutrients in a mixed-grass prairie. Ecology 71:1959-1967.

## Contributors

Scott Brady  
Stuart Veith

## Approval

Kirt Walstad, 4/25/2025

## Acknowledgments

A number of USDA-NRCS staff supported this project. Staff contributions are as follows: Soil Concepts, Soils Information, and Field Descriptions Charlie French, USDA-NRCS (retired) Steve Sieler, USDA-NRCS NASIS Reports, Data Dumps, and Soil Sorts Bill Drummond, USDA-NRCS (retired) Pete Weikle, USDA-NRCS Peer Review Kirt Walstad, USDA-NRCS Mark Hayek, USDA-NRCS Kami Kilwine, USDA-NRCS Robert Mitchell, USDA-NRCS Quality Control Kirt Walstad, USDA-NRCS Quality Assurance Stacey Clark, USDA-NRCS

## Rangeland health reference sheet

**Interpreting Indicators of Rangeland Health** is a qualitative assessment protocol used to determine ecosystem condition based on benchmark characteristics described in the Reference Sheet. A suite of 17 (or more) indicators are typically considered in an assessment. The ecological site(s) representative of an assessment location must be known prior to applying the protocol and must be verified based on soils and climate. Current plant community cannot be used to identify the ecological site.

Author(s)/participant(s)	
Contact for lead author	
Date	04/25/2025
Approved by	
Approval date	
Composition (Indicators 10 and 12) based on	Annual Production

## Indicators

### 1. Number and extent of rills:

---

2. Presence of water flow patterns:

---

3. Number and height of erosional pedestals or terracettes:

---

4. Bare ground from Ecological Site Description or other studies (rock, litter, lichen, moss, plant canopy are not bare ground):

---

5. Number of gullies and erosion associated with gullies:

---

6. Extent of wind scoured, blowouts and/or depositional areas:

---

7. Amount of litter movement (describe size and distance expected to travel):

---

8. Soil surface (top few mm) resistance to erosion (stability values are averages - most sites will show a range of values):

---

9. Soil surface structure and SOM content (include type of structure and A-horizon color and thickness):

---

10. Effect of community phase composition (relative proportion of different functional groups) and spatial distribution on infiltration and runoff:

---

11. Presence and thickness of compaction layer (usually none; describe soil profile features which may be mistaken for compaction on this site):

---

12. Functional/Structural Groups (list in order of descending dominance by above-ground annual-production or live foliar cover using symbols: >>, >, = to indicate much greater than, greater than, and equal to):

Dominant:

Sub-dominant:

Other:

Additional:

---

13. Amount of plant mortality and decadence (include which functional groups are expected to show mortality or decadence):

---

14. Average percent litter cover (%) and depth ( in):

---

15. Expected annual annual-production (this is TOTAL above-ground annual-production, not just forage annual-production):

---

16. Potential invasive (including noxious) species (native and non-native). List species which BOTH characterize degraded states and have the potential to become a dominant or co-dominant species on the ecological site if their future establishment and growth is not actively controlled by management interventions. Species that become dominant for only one to several years (e.g., short-term response to drought or wildfire) are not invasive plants. Note that unlike other indicators, we are describing what is NOT expected in the reference state for the ecological site:

---

17. Perennial plant reproductive capability:

---