

Ecological site R062XB024SD

Shallow Loamy - High Central

Last updated: 8/19/2024
Accessed: 07/22/2026

General information

Provisional. A provisional ecological site description has undergone quality control and quality assurance review. It contains a working state and transition model and enough information to identify the ecological site.

MLRA notes

Major Land Resource Area (MLRA): 062X–Black Hills

The Black Hills (MLRA 62) is a unique, low lying mountain range situated in the midst of a mixed short and mid-grass prairie. It is a true Island in the Plains, as it has geophysical and biological attributes that are unlike the surrounding area. The Black Hills have strong floristic ties to four of the North American biomes: Cordilleran (Rocky Mountain) Forest, Northern Coniferous Forest, Eastern Deciduous Forest, and Grasslands. MLRA 62 is approximately 3,040 square miles in size; 74 percent is located in South Dakota, and 26 percent is in Wyoming. The towns of Lead, Deadwood, Hill City, and Custer, South Dakota, are in this area. U.S. Highways 16 and 385 cross the MLRA. The Black Hills National Forest, Custer State Park, Mt. Rushmore National Monument, Jewel Cave National Monument, and Wind Cave National Park are located in this MLRA. This area forms the core of the Black Hills and the Bear Lodge Mountains where the elevation ranges between 3,600 to 6,565 feet, however, Black Elk Peak (Harney Peak) rises to 7,242 feet. Slopes range from moderately sloping on some of the high plateaus to very steeply sloping along drainageways and on peaks and ridges. Narrow valleys generally are gently sloping to strongly sloping. The Black Hills uplift is the product of the Laramide mountain-building episodes that produced most of the ranges in the Rocky Mountains. Uplift began near the end of the Cretaceous period, 65 million years ago and ended by 35 million years ago (Froiland 1999). The core of the Black Hills is a plutonic mass of granite with steeply dipping metamorphic rocks, primarily slate and schist, which directly surrounds the granite core. A plateau of Mississippian limestone surrounds the igneous and metamorphic rock core. The Madison limestone is broken around the outer edges of the uplifted area. The Permian Minnekahta limestone forms the outermost boundary of the area. Many other tilted sandstone, shale, and limestone units are exposed like a bathtub ring inside the steeply dipping Madison limestone. The dominant soil orders in this MLRA are Alfisols (forest soils) and Mollisols (grassland soils). The soils in the area have a frigid or cryic soil temperature regime, a udic or ustic soil moisture regime, and mixed, micaceous, or smectitic mineralogy. They are shallow to very deep, generally well drained, and loamy in texture. The Black Hills MLRA supports open to dense forest vegetation. Ponderosa pine is the dominant species across the Black Hills. White spruce grows at the higher elevations and along the major drainageways. Bur oak is found intermixed with pine in the northern and eastern fringes of the Black Hills, and Rocky Mountain Juniper is most common in the southern portion of the Black Hills. Aspen is of minor extent throughout the Black Hills area. Roughleaf ricegrass, slender wheatgrass, bearded wheatgrass, poverty oatgrass, Richardson's needlegrass, and mountain ricegrass are the most common native grasses under open forest stands. The most common native shrubs are common snowberry, bearberry, common juniper, Oregon grape, and Saskatoon serviceberry. MLRA 62 land ownership is approximately 47 percent private and 53 percent federal. Rangeland and forestland are split almost equally between private and federal ownership (47 percent each). Minor areas of land are privately owned cropland and urban development. The forestland in this area is used mainly for timber production, recreation, and grazing. The major resource concerns are soil erosion and surface compaction caused by logging, mining, wildfires, grazing, and urban expansion. The quality of ground and surface water is another concern, especially in the northern part of the Black Hills. The primary cause is contamination from mine waste and septic systems in areas of rural development and urban expansion (USDA-NRCS, 2006: Ag Handbook 296).

LRU notes

For development of ecological sites, MLRA 62 is divided into three Land Resource Units (LRU's) or physiographic zones (A, B, C, and Y). Each LRU has a set of ecological sites that represents these zones. The LRU is identified in the Ecological Site ID: R062XY000SD; "062X" identifies the MLRA, the next letter "Y" identifies the LRU. Note: The organization of Ecological Site ID's will likely change in the future. LRU-A includes the northern Black Hills and Bear Lodge Mountains (22-30" PZ); LRU-B includes the high elevation central core of the Black Hills (25-35" PZ); and LRU-C includes the southern portion of the Black Hills (17-21" PZ). The Forest ecological sites are representative of sites in the Black Hills, Bear Lodge Mountains (MLRA-62), and the surrounding Dakota Hogback (MLRA-61). These sites are separated by elevation, soil temperature regimes, and slope. The Low Mountain area includes all of the Black Hills, Bear Lodge Mountains, and Dakota Hogback below 6,200 feet in elevation (LRU's A and C). The soils in this area have a frigid soil temperature regime. The High Mountain area includes all of the Black Hills above 6,200 feet elevation (LRU-B). The soils in this area have a cryic soil temperature regime.

Classification relationships

USDA Land Resource Region G—Western Great Plains Range and Irrigated Region: Major Land Resource Area (MLRA) 62—Black Hills
US Environmental Protection Agency (EPA) Level IV Ecoregions of the Conterminous United States: Black Hills Plateau—17b Black Hills
Core Highlands—17c USDA Forest Service Ecological Subregions: Sections and Subsections of Conterminous United States: Black Hills
Coniferous Forest Province—M334: Black Hills Section—334A Black Hills Limestone Plateau-Core Highlands Subsection—M334Ab

Ecological site concept

This site was previously referred to as a High Country Shallow range site. The site is located on upland landscapes within the central portion of the Black Hills, in Land Resource Unit (LRU)-B. The soils are shallow, 10 to 20 inches in depth, with a restrictive limestone bedrock layer which impedes water movement and rooting depth. The surface layer is loamy and ranges from 3 to 7 inches in depth. The soils typically are calcareous to the surface. The slopes can range from 2 to 30 percent. The site does not receive additional water from runoff or overflow. Vegetation in the Reference State consists primarily of cool-season grass species, with warm-season species as a minor component. The dominant grasses include bearded wheatgrass, western wheatgrass, Columbia needlegrass, prairie Junegrass, and needleandthread or porcupinegrass. Plains muhly and prairie dropseed are the dominant warm-season grasses. Forbs are common and diverse but never dominant, and shrubs such as shrubby cinquefoil and common juniper are often present in the plant community. Ponderosa pine or Black Hills spruce can be present in Reference condition, but are sparsely scattered across the site and typically do not exceed 1 percent canopy cover. This site is very susceptible to pine encroachment.

Associated sites

R062XB010SD	Loamy - High Central The Loamy 62B site has deep soils and located on shallower slopes above the Shallow Loamy site.
R062XY043SD	Valley Loam The Valley Loam ecological site can be adjacent to shallow loamy site in a lower positions on the landscape.

Similar sites

R062XB010SD	Loamy - High Central The Loamy 62B site will have a similar plant community but with more production, fewer forbs and shrubs. The Loamy site appears to be more susceptible to conifer encroachment than the Shallow Loamy in this LRU.
-------------	---

Table 1. Dominant plant species

Tree	Not specified
Shrub	(1) <i>Dasiphora fruticosa</i> ssp. <i>floribunda</i> (2) <i>Juniperus communis</i>
Herbaceous	(1) <i>Elymus trachycaulus</i> subsp. <i>subsecundus</i> (2) <i>Achnatherum nelsonii</i>

Physiographic features

This site occurs on gently sloping to very steep summits, shoulders, and backslopes in the Black Hills.

Table 2. Representative physiographic features

Landforms	(1) Ridge (2) Mountain slope (3) Structural bench
Runoff class	Low to very high
Flooding frequency	None
Ponding frequency	None
Elevation	1,890 – 2,130 m
Slope	0 – 60 %
Water table depth	200 cm
Aspect	Aspect is not a significant factor

Climatic features

MLRA 62 is in a microclimate caused by the influence of increased elevation which leads to increased precipitation, moderate air temperature, and lower wind velocities compared to the surrounding Great Plains. In general, the Black Hills climate is a continental type, cold in the winter and hot in the summer. Annual precipitation in MLRA 62 typically increases with elevation, and decreases from west to east and north to south. The average annual precipitation range for LRU-B (Central High Country) is 25 to 35 inches. Most of the rainfall occurs as frontal storms early in the growing season, in May and June. Some high-intensity, convective thunderstorms occur in July and August. Precipitation in the winter occurs mostly as snow. The annual average snowfall ranges from 23 inches at the lower elevations in the south, to 54 inches at the higher elevations in the central part of MLRA 62. The average annual temperature ranges from 36 to 48 degrees F. January is the coldest month, with an average temperature of 22°F in the central part and 25°F in the southern part of MLRA 62. July is the warmest month, with an average daily temperature of 67°F in the central part and 73°F in the southern part of this MLRA. The frost-free period ranges from 129 to 168 days. It is shortest at higher elevations and in the northwestern part of the MLRA. Hourly winds are estimated to average about 11 miles per hour (mph) annually. Growth of cool-season plants begins in April, slowing or ceasing growth by mid-August. Warm-season plants begin growth in May, and continue to mid-September. Regrowth of cool-season plants may occur in September and October, depending upon soil moisture availability.

* Deerfield 3 SE (USC00392231) climate station is not located within LRU-B, but is adjacent to it in LRU-A. The mean annual precipitation recorded at this station is less than what LRU-B actually receives.

Table 3 Representative climatic features

Frost-free period (characteristic range)	20 days
Freeze-free period (characteristic range)	50 days
Precipitation total (characteristic range)	530 mm

Frost-free period (actual range)	20 days
Freeze-free period (actual range)	50 days
Precipitation total (actual range)	530 mm
Frost-free period (average)	20 days
Freeze-free period (average)	50 days
Precipitation total (average)	530 mm

- (1) DEERFIELD 3 SE [USC00392231], Hill City, SD

Influencing water features

This site does not receive additional water from runoff or overflow.

Soil features

The soils on this site are shallow and well to somewhat excessively drained. The surface layer ranges from about 3 to 7 inches thick. Encroachment of ponderosa pine and/or Black Hills spruce is not common, but may occur in some areas. If present, the mineral soil surface could be covered with 1 to 2 inches of pine needles and duff. Most soils on this site are calcareous to the surface. The slopes range from 2 to 30 percent; slopes may exceed 30 percent in localized situations.

This site typically shows slight to no evidence of rills, wind-scoured areas, or pedestalled plants. If present, water flow paths could include long, continuous rills, or they could be broken, irregular in appearance, or discontinuous. The soil surface is stable and intact under normal conditions. There is a restrictive layer of bedrock (typically limestone, less commonly sandstone) at about 10 to 20 inches in depth. This material usually is fractured at intervals of about 6 to more than 20 inches apart. Although fractured, this bedrock impedes water movement and root penetration. These soils mainly are susceptible to water erosion. The hazard of water erosion increases on slopes greater than about 10 percent, and is more common on soils with few rock fragments. Loss of 50 percent or more of the surface layer of the soils on this site can result in a shift in species composition and/or production.

The commonly-occurring soils for this site is Rockerville, cool. It is a taxadjunct to the Rockerville series because it has cooler soil temperatures than are typical.

Access Web Soil Survey (<http://websoilsurvey.nrcs.usda.gov/app/>) for specific local soils information.

Figure 7. General Soil Profile for SwLy-B

Table 4. Representative soil features

Parent material	(1) Residuum – limestone and siltstone
Surface texture	(1) Gravelly silt loam (2) Silt loam

Family particle size	(1) Loamy
Drainage class	Well drained to somewhat excessively drained
Permeability class	Moderate to moderately rapid
Soil depth	30 – 50 cm
Surface fragment cover <=3"	0 – 20 %
Surface fragment cover >3"	0 – 10 %
Available water capacity (0-101.6cm)	2.54 – 7.62 cm
Calcium carbonate equivalent (0-101.6cm)	0 – 20 %
Electrical conductivity (0-101.6cm)	Not specified
Sodium adsorption ratio (0-101.6cm)	Not specified
Soil reaction (1:1 water) (0-101.6cm)	7.4 – 8.4
Subsurface fragment volume <=3" (Depth not specified)	30 – 60 %
Subsurface fragment volume >3" (Depth not specified)	0 – 30 %

Ecological dynamics

Black Hills vegetation types consist of a mixture of forest and grasslands resulting from the varied topography, geology, soils, climate and natural disturbances. Frequent fires, periodic drought, and episodic infestations of mountain pine beetles all contribute to the maintenance of large open grasslands scattered throughout the Black Hills. Ponderosa pine is the dominant tree species in the Black

Hills. It is a fire-adapted species that coexists with frequent, low-intensity fires that consume small seedlings, prune lower branches from larger trees, and reduce fuel loads on the forest floor.

This site developed under Black Hills climatic conditions with short-term weather variations, light to severe grazing by bison, elk, and small mammals, sporadic natural or man-caused wildfire (often of light intensities), and other biotic and abiotic factors that typically influence soil/site development. The natural fire regime maintained this site as a grassland and the plant communities were free of non-native cool-season grasses.

Fire, or the lack of fire, grazing and potential conifer encroachment shaped this site and adjacent ecological sites. Between 1388 and 1900, fire intervals in the Black Hills ranged from 16 to 20 years (Brown and Hull-Sieg, 1996). In the absence of fire, ponderosa pine and/or Black Hills spruce can encroach on this site, but it appears to be uncommon in this LRU. The reason is not fully understood, but competition with cool-season grasses which dominate the site could be hindering the establishment of pine seedlings.

Changes will occur in the plant communities due to short-term weather variations, impacts of native and/or non-native plant and animal species, and management actions. Severe disturbances, such as periods of well-below average precipitation, severe defoliation, or no fire and no use can also cause significant shifts in plant communities and species composition.

The historic native vegetation on this site consisted of mixed prairie grasses occurring on ridges and uplands. The vegetation predominately is cool-season grasses mixed with warm-season grasses, sedges, shrubs and a diverse forb community. Bearded wheatgrass, western wheatgrass, Columbia needlegrass, needleandthread, porcupinegrass, and prairie Junegrass are the dominant cool-season grasses. Plains muhly and prairie dropseed are the dominant warm-season grasses. Subdominant cool-season species include Richardson needlegrass and threadleaf sedge. Kentucky bluegrass and timothy can be present, but typically do not dominate the plant community. Forbs make up a significant proportion of the plant community. Common forbs include stemless hymenoxys, Indian paintbrush, dotted gayfeather, cudweed sagewort, stiff goldenrod, prairie coneflower, hairy goldaster, hood phlox, and pussytoes. Shrubs include shrubby cinquefoil, common juniper, fringed sagewort, and skunkbush sumac. Ponderosa pine or Black Hills spruce can be present, but are sparsely scattered across the site. Rocks and rock fragments are very noticeable, and bare ground can be present but is not common.

The following diagram illustrates the common plant community phases that can occur on the site and the transition pathways between communities. These are the most common plant community phases based on current knowledge and experience, and changes will be made as more data is collected. Narratives following the diagram contain more detail pertaining to the ecological processes.

State and transition model

Figure 8. Shallow Loamy - 062XB024SD - LRU-B

Figure 9. Shallow Loamy - 062XBSD - LRU-B

Additional community tables

Table 5. Community 1.1 plant community composition

Group	Common Name	Symbol	Scientific Name	Annual Production ()	Foliar Cover (%)
Grass/Grasslike					
1	Cool-Season Bunchgrass			740-1603	
	slender wheatgrass	ELTR7	<i>Elymus trachycaulus</i>	247-493	–
	slender wheatgrass	ELTRS	<i>Elymus trachycaulus ssp. subsecundus</i>	247-493	–
	Columbia needlegrass	ACNE9	<i>Achnatherum nelsonii</i>	123-370	–
	Richardson's needlegrass	ACRI8	<i>Achnatherum richardsonii</i>	123-370	–
	Rocky Mountain fescue	FESA	<i>Festuca saximontana</i>	25-247	–
	green needlegrass	NAVI4	<i>Nassella viridula</i>	25-247	–
	mountain brome	BRMA4	<i>Bromus marginatus</i>	0-123	–
	needle and thread	HECOC8	<i>Hesperostipa comata ssp. comata</i>	0-123	–
	nodding brome	BRAN	<i>Bromus anomalus</i>	0-123	–
	prairie Junegrass	KOMA	<i>Koeleria macrantha</i>	0-123	–
	Pumpelly's brome	BRINP	<i>Bromus inermis ssp. pumpellianus</i>	0-123	–
	timber oatgrass	DAIN	<i>Danthonia intermedia</i>	0-123	–
2	Cool-Season Rhizomatous Grass			247-493	

	western wheatgrass	PASM	<i>Pascopyrum smithii</i>	247-493	–
3	Tall and Mid- Warm-Season Bunchgrasses			123-247	
	prairie dropseed	SPHE	<i>Sporobolus heterolepis</i>	123-247	–
	composite dropseed	SPCOC2	<i>Sporobolus compositus</i> var. <i>compositus</i>	0-123	–
4	Mid- Warm-Season Rhizomatous Grasses			0-247	
	plains muhly	MUCU3	<i>Muhlenbergia cuspidata</i>	0-247	–
5	Other Native Grasses			0-123	
	Grass, perennial	2GP	<i>Grass, perennial</i>	0-123	
6	Grass-Likes			123-247	
	Hood's sedge	CAHO5	<i>Carex hoodii</i>	49-123	–
	needleleaf sedge	CADU6	<i>Carex duriuscula</i>	0-123	–
	Richardson's sedge	CARI	<i>Carex richardsonii</i>	49-123	–
	sedge	CAREX	<i>Carex</i>	25-123	–
	Grass-like (not a true grass)	2GL	<i>Grass-like (not a true grass)</i>	0-123	–
7	Non-Native Cool-Season Grasses			0	
Forb					
8	Forbs			123-370	
	goldenrod	SOLID	<i>Solidago</i>	25-247	–
	white sagebrush	ARLU	<i>Artemisia ludoviciana</i>	0-123	–
	western yarrow	ACMIO	<i>Achillea millefolium</i> var. <i>occidentalis</i>	25-123	–
	white prairie aster	SYFA	<i>Symphyotrichum falcatum</i>	25-123	–
	dotted blazing star	LIPU	<i>Liatris punctata</i>	25-123	–
	old man's whiskers	GETR	<i>Geum triflorum</i>	25-123	–
	scurfpea	PSORA2	<i>Psoraleidium</i>	25-123	–
	blacksamson echinacea	ECAN2	<i>Echinacea angustifolia</i>	0-49	–
	fleabane	ERIGE2	<i>Erigeron</i>	0-49	–
	wild bergamot	MOFI	<i>Monarda fistulosa</i>	0-49	–
	Forb, perennial	2FP	<i>Forb, perennial</i>	0-49	–
	hairy false goldenaster	HEVI4	<i>Heterotheca villosa</i>	0-49	–
	mariposa lily	CALOC	<i>Calochortus</i>	0-49	–
	beardtongue	PENST	<i>Penstemon</i>	0-49	–
	prairie clover	DALEA	<i>Dalea</i>	0-49	–
	American vetch	VIAM	<i>Vicia americana</i>	0-49	–
	blanketflower	GAAR	<i>Gaillardia aristata</i>	0-49	–
	Lewis flax	LILE3	<i>Linum lewisii</i>	0-49	–
	bluebell bellflower	CARO2	<i>Campanula rotundifolia</i>	0-49	–
	cinquefoil	POTEN	<i>Potentilla</i>	0-49	–
	deathcamas	ZIGAD	<i>Zigadenus</i>	0-25	–
	spiny phlox	PHHO	<i>Phlox hoodii</i>	0-25	–
	common starlily	LEMO4	<i>Leucocrinum montanum</i>	0-25	–
Shrub/Vine					
9	Shrubs			49-247	
	common juniper	JUCO6	<i>Juniperus communis</i>	49-247	–
	prairie sagewort	ARFR4	<i>Artemisia frigida</i>	25-123	–
	field sagewort	ARCA12	<i>Artemisia campestris</i>	0-123	–
	shrubby cinquefoil	DAFRF	<i>Dasiphora fruticosa</i> ssp. <i>floribunda</i>	25-123	–

	Shrub (>.5m)	2SHRUB	<i>Shrub (>.5m)</i>	0-49	–
Tree					
10	Trees			0-25	
	ponderosa pine	PIPO	<i>Pinus ponderosa</i>	0-25	–
	white spruce	PIGL	<i>Picea glauca</i>	0-25	–

Table 6. Community 1.2 plant community composition

Group	Common Name	Symbol	Scientific Name	Annual Production ()	Foliar Cover (%)
-------	-------------	--------	-----------------	----------------------	------------------

Table 7. Community 2.1 plant community composition

Group	Common Name	Symbol	Scientific Name	Annual Production ()	Foliar Cover (%)
-------	-------------	--------	-----------------	----------------------	------------------

Other information

Revision Notes: This PROVISIONAL ecological site concept has been QCd to ensure that the site meets the NESH standards for a provisional ecological site that provides basic compiled information in one location. This site should not be considered an Approved ESD, as it is only the foundational site concepts and requires further data collection—specifically high-intensity data characterizations and full 232 soil descriptions—and further site investigations and final STM reviews before it can be used as an Approved ESD meeting NESH standards. Site Development and Testing Plan: Future work, as described in a Project Plan, to validate the information in this Provisional Ecological Site Description is needed. This will include field activities to collect low, medium and high intensity sampling, soil correlations, and analysis of that data. Annual field reviews should be done by soil scientists and vegetation specialists. A final field review, peer review, quality control, and quality assurance reviews of the ESD will be needed to produce the final document.

Inventory data references

Information presented here has been derived from NRCS clipping data and other inventory data. Field observations from range-trained personnel also were used. Those involved in developing this site include Lakhdar Benkobi, ESI/SRIC, NRCS; Stan Boltz, Range Management Specialist, NRCS; Dan Brady, Soil Scientist, NRCS; Mitch Faulkner, Range Management Specialist, NRCS; Roger Gates, Associate Professor/Rangeland Management Specialist, West River Ag Center; Rick Peterson, Ecological Site Specialist, NRCS; Matthew, Scott, Botanist - USFS Hell Canyon District Ranger; L. Michael Stirling, Range Management Specialist, NRCS; and Jim Westerman, Soil Scientist, NRCS. All inventory information and data records are compiled within the Rapid City, SD USDA-NRCS Shared “S” network drive.

Other references

Brown, P. M. and C. Hull-Sieg. 1996. Fire history in interior ponderosa pine communities of the Black Hills, South Dakota, USA, Int. J. Wildland Fire 6(3): 97-105.

High Plains Regional Climate Center, University of Nebraska. (<http://www.hprcc.unl.edu/>) USDA, NRCS.

National Water and Climate Center. (<http://wcc.nrcs.usda.gov>) USDA, NRCS. National Range and Pasture Handbook. September, 1997.

USDS, NRCS. 2006. Land Resource Regions and Major Land Resource Areas of the United States, the Caribbean, and the Pacific Basin. U.S. Department of Agriculture Handbook 296.

USDA, NRCS. National Soil Information System, Information Technology Center. (<http://nasis.nrcs.usda.gov>)

USDA, NRCS. 2001. The PLANTS Database, Version 3.1 (<http://plants.usda.gov>). National Plant Data Center.

USDA, NRCS. National Ecological Site Handbook, 1st Ed. January, 2014.

Contributors

Westerman James
Peterson Rick L.

Approval

Suzanne Mayne-Kinney, 8/19/2024

Acknowledgments

MLRA 62 sites were written to the Provisional Level by Rick L. Peterson, ESS, Rapid City, SSO in FY20. The ESDs were reviewed for quality control by Emily Helms, John Hartung, Mitch Faulkner, and Ryan Murray. All ecological sites were then reviewed and approved at the Provisional Level by David Kraft, Regional ESS, Salina, KS in September 2020.

Rangeland health reference sheet

Interpreting Indicators of Rangeland Health is a qualitative assessment protocol used to determine ecosystem condition based on benchmark characteristics described in the Reference Sheet. A suite of 17 (or more) indicators are typically considered in an assessment. The ecological site(s) representative of an assessment location must be known prior to applying the protocol and must be verified based on soils and climate. Current plant community cannot be used to identify the ecological site.

Author(s)/participant(s)	
Contact for lead author	
Date	07/22/2026
Approved by	
Approval date	
Composition (Indicators 10 and 12) based on	Annual Production

Indicators

1. Number and extent of rills:

2. Presence of water flow patterns:

3. Number and height of erosional pedestals or terracettes:

4. Bare ground from Ecological Site Description or other studies (rock, litter, lichen, moss, plant canopy are not bare ground):

5. Number of gullies and erosion associated with gullies:

6. Extent of wind scoured, blowouts and/or depositional areas:

7. Amount of litter movement (describe size and distance expected to travel):

8. Soil surface (top few mm) resistance to erosion (stability values are averages - most sites will show a range of values):

9. Soil surface structure and SOM content (include type of structure and A-horizon color and thickness):

10. Effect of community phase composition (relative proportion of different functional groups) and spatial distribution on infiltration and runoff:

11. Presence and thickness of compaction layer (usually none; describe soil profile features which may be mistaken for compaction on this site):

12. Functional/Structural Groups (list in order of descending dominance by above-ground annual-production or live foliar cover using symbols: >>, >, = to indicate much greater than, greater than, and equal to):

Dominant:

Sub-dominant:

Other:

Additional:

13. Amount of plant mortality and decadence (include which functional groups are expected to show mortality or decadence):

14. Average percent litter cover (%) and depth (in):

15. Expected annual annual-production (this is TOTAL above-ground annual-production, not just forage annual-production):

16. Potential invasive (including noxious) species (native and non-native). List species which BOTH characterize degraded states and have the potential to become a dominant or co-dominant species on the ecological site if their future establishment and growth is not actively controlled by management interventions. Species that become dominant for only one to several years (e.g., short-term response to drought or wildfire) are not invasive plants. Note that unlike other indicators, we are describing what is NOT expected in the reference state for the ecological site:

17. Perennial plant reproductive capability:
