

Ecological site R062XC024SD

Shallow Loamy - South

Last updated: 7/31/2024
Accessed: 07/12/2026

General information

Provisional. A provisional ecological site description has undergone quality control and quality assurance review. It contains a working state and transition model and enough information to identify the ecological site.

MLRA notes

Major Land Resource Area (MLRA): 062X–Black Hills

The Black Hills (MLRA 62) is a unique, low lying mountain range situated in the midst of a mixed short and mid-grass prairie. It is a true Island in the Plains, as it has geophysical and biological attributes that are unlike the surrounding area. The Black Hills have strong floristic ties to four of the North American biomes: Cordilleran (Rocky Mountain) Forest, Northern Coniferous Forest, Eastern Deciduous Forest, and Grasslands. MLRA 62 is approximately 3,040 square miles in size; 74 percent is located in South Dakota, and 26 percent is in Wyoming. The towns of Lead, Deadwood, Hill City, and Custer, South Dakota, are in this area. U.S. Highways 16 and 385 cross the MLRA. The Black Hills National Forest, Custer State Park, Mt. Rushmore National Monument, Jewel Cave National Monument, and Wind Cave National Park are located in this MLRA. This area forms the core of the Black Hills and the Bear Lodge Mountains where the elevation ranges between 3,600 to 6,565 feet, however, Black Elk Peak (Harney Peak) rises to 7,242 feet. Slopes range from moderately sloping on some of the high plateaus to very steeply sloping along drainageways and on peaks and ridges. Narrow valleys generally are gently sloping to strongly sloping. The Black Hills uplift is the product of the Laramide mountain-building episodes that produced most of the ranges in the Rocky Mountains. Uplift began near the end of the Cretaceous period, 65 million years ago and ended by 35 million years ago (Froiland 1999). The core of the Black Hills is a plutonic mass of granite with steeply dipping metamorphic rocks, primarily slate and schist, which directly surrounds the granite core. A plateau of Mississippian limestone surrounds the igneous and metamorphic rock core. The Madison limestone is broken around the outer edges of the uplifted area. The Permian Minnekahta limestone forms the outermost boundary of the area. Many other tilted sandstone, shale, and limestone units are exposed like a bathtub ring inside the steeply dipping Madison limestone. The dominant soil orders in this MLRA are Alfisols (forest soils) and Mollisols (grassland soils). The soils in the area have a frigid or cryic soil temperature regime, a udic or ustic soil moisture regime, and mixed, micaceous, or smectitic mineralogy. They are shallow to very deep, generally well drained, and loamy in texture. The Black Hills MLRA supports open to dense forest vegetation. Ponderosa pine is the dominant species across the Black Hills. White spruce grows at the higher elevations and along the major drainageways. Bur oak is found intermixed with pine in the northern and eastern fringes of the Black Hills, and Rocky Mountain juniper is most common in the southern portion of the Black Hills. Aspen is of minor extent throughout the Black Hills area. Roughleaf ricegrass, slender wheatgrass, bearded wheatgrass, poverty oatgrass, Richardson's needlegrass, and mountain ricegrass are the most common native grasses under open forest stands. The most common native shrubs are common snowberry, bearberry, common juniper, Oregon grape, and Saskatoon serviceberry. MLRA 62 land ownership is approximately 47 percent private and 53 percent federal. Rangeland and forestland are split almost equally between private and federal ownership (47 percent each). Minor areas of land are privately owned cropland and urban development. The forestland in this area is used mainly for timber production, recreation, and grazing. The major resource concerns are soil erosion and surface compaction caused by logging, mining, wildfires, grazing, and urban expansion. The quality of ground and surface water is another concern, especially in the northern part of the Black Hills. The primary cause is contamination from mine waste and septic systems in areas of rural development and urban expansion (USDA-NRCS, 2006: Ag Handbook 296).

LRU notes

For development of ecological sites, MLRA 62 is divided into three Land Resource Units (LRU's) or physiographic zones (A, B, C, and Y). Each LRU has a set of ecological sites that represents these zones. The LRU is identified in the Ecological Site ID: R062XY000SD; "062X" identifies the MLRA, the next letter "Y" identifies the LRU. Note: The organization of Ecological Site ID's will likely change in the future. LRU-A includes the northern Black Hills and Bear Lodge Mountains (22-30" PZ); LRU-B includes the high elevation central core of the Black Hills (25-35" PZ); and LRU-C includes the southern portion of the Black Hills (17-21" PZ). The Forest ecological sites are representative of sites in the Black Hills, Bear Lodge Mountains (MLRA-62), and the surrounding Dakota Hogback (MLRA-61). These sites are separated by elevation, soil temperature regimes, and slope. The Low Mountain area includes all of the Black Hills, Bear Lodge Mountains, and Dakota Hogback below 6,200 feet in elevation (LRU's A and C). The soils in this area have a frigid soil temperature regime. The High Mountain area includes all of the Black Hills above 6,200 feet elevation (LRU-B). The soils in this area have a cryic soil temperature regime.

Classification relationships

USDA Land Resource Region G—Western Great Plains Range and Irrigated Region: Major Land Resource Area (MLRA) 62—Black Hills
US Environmental Protection Agency (EPA) Level IV Ecoregions of the Conterminous United States: Black Hills Plateau—17b Black Hills
Core Highlands—17c USDA Forest Service Ecological Subregions: Sections and Subsections of Conterminous United States: Black Hills
Coniferous Forest Province—M334: Black Hills Section—334A Black Hills Limestone Plateau-Core Highlands Subsection—M334Ab

Ecological site concept

This site is located on upland landscapes in the southern portion of the Black Hills (LRU-C). Soils on this site are shallow, between 10 and 20 inches in depth, with a loamy surface layer ranging from 2 to 6 inches in depth. Most soils on this site are calcareous to the surface. The soils are well to excessively drained with a restrictive bedrock layer of sandstone or limestone, which impedes water movement and rooting depth. Slope can range from 2 to 60 percent. The site does not receive additional water from runoff or overflow. Vegetation in reference consist primarily of warm season grass species with cool season species being sub-dominant. The dominant warm season grasses include little bluestem, big bluestem, and sideoats grama. Cool season grasses include western wheatgrass and needleandthread. Forbs are common and diverse but never dominant, shrubs such as western snowberry, skunkbush sumac, prairie rose and green sagewort are often present in the plant community. Ponderosa pine can be scattered throughout the site but will not exceed 2 percent canopy cover. This site is very susceptible to pine encroachment.

Associated sites

R062XC010SD	Loamy - South The loamy ecological site can be adjacent to shallow loamy and often at lower positions on the landscape.
R062XY029SD	Stony Hills Stony Hills ecological site can be adjacent to shallow loamy and often higher on the landscape.

Similar sites

R062XY029SD	Stony Hills Stony Hills more big bluestem and lead plant, higher potential for ponderosa pine, higher production.
-------------	--

Table 1. Dominant plant species

Tree	Not specified
Shrub	Not specified
Herbaceous	(1) <i>Schizachyrium scoparium</i> (2) <i>Bouteloua curtipendula</i>

Physiographic features

MLRA 62 forms the core of the Black Hills which consists mainly of rounded domes and peaks. The shallow loamy site occurs mostly on gently sloping to very steep summits, shoulders, and backslopes. Slopes range from moderately sloping on some of the high plateaus to very steep along drainages and ridges.

Figure 1. Block Diagram of Black Hills

Table 2. Representative physiographic features

Landforms	(1) Ridge (2) Mountain slope (3) Structural bench
Runoff class	Low to very high
Flooding frequency	None
Ponding frequency	None
Elevation	1,160 – 1,680 m
Slope	0 – 60 %
Water table depth	200 cm
Aspect	SE, S

Climatic features

MLRA 62 is in a microclimate caused by the influence of increased elevation which leads to increased precipitation, moderate air temperature, and lower wind velocities as compared to the surrounding Great Plains. In general, the Black Hills climate is a continental type, cold in the winter and hot in the summer. Annual precipitation in MLRA 62 typically increases with elevation and decreases from west to east and north to south. The average annual precipitation range for LRU-C (South) is 17 to 21 inches. Most of the rainfall occurs as frontal storms early in the growing season, May and June. Some high-intensity, convective thunderstorms occur in July and August. Precipitation in the winter occurs mostly as snow. The annual average snowfall ranges from 23 inches at the lower elevations in the south to 54 inches at the higher elevations in the central part of MLRA 62. Average annual temperature ranges from 36 to 48 degrees F. January is the coldest month, with an average temperature of 22 °F in the central part and 25 °F in the southern part of MLRA 62. July is the warmest month, with an average daily temperature of 67 °F in the central part and 73 °F in the southern part of this MLRA. The frost free period ranges from 129 to 168 days. It is shortest at higher elevations and in the northwestern part of the MLRA. Hourly winds are estimated to average about 11 miles per hour (mph) annually. Growth of cool-season plants begins in April, slowing or ceasing growth by Mid-August. Warm-season plants begin growth in May, and continue to Mid-September. Regrowth of cool-season plants may occur in September and October depending on soil moisture availability.

Table 3 Representative climatic features

Frost-free period (characteristic range)	90-110 days
Freeze-free period (characteristic range)	120-130 days
Precipitation total (characteristic range)	480-510 mm
Frost-free period (actual range)	80-110 days

Freeze-free period (actual range)	120-130 days
Precipitation total (actual range)	460-510 mm
Frost-free period (average)	100 days
Freeze-free period (average)	120 days
Precipitation total (average)	480 mm

- (1) CUSTER [USC00392087], Custer, SD
- (2) HOT SPRINGS [USC00394007], Hot Springs, SD
- (3) WIND CAVE [USC00399347], Buffalo Gap, SD

Influencing water features

This site is not influenced by streams or wetlands

Soil features

The soils on this site are shallow and well to somewhat excessively drained. The surface layer ranges from 2 to 6 inches thick. Encroachment of ponderosa pine is common in many areas, thereby causing the mineral soil surface to be covered with 1 to 2 inches of pine needles and duff. Surface textures are variable and are listed below. Most soils on this site are calcareous to the surface. There are a few places where the soil parent material is derived from noncalcareous sources (primarily sandstone or schist), and the soils are noncalcareous. Slopes range from 2 to 60 percent.

This site typically show slight to no evidence of rills, wind scoured areas, or pedestalled plants. If present, water flow paths could include long, continuous, shallow gullies, or they could be broken, irregular in appearance or discontinuous. The soil surface is stable and intact. There is a restrictive layer of bedrock (typically sandstone or limestone) at about 10 to 20 inches in depth which impedes water movement and root penetration. These soils are mainly susceptible to water erosion. The hazard of water erosion increases on slopes greater than about 10 percent, and is more common on soils with few rock fragments. Loss of 50 percent or more of the surface layer of the soils on this site can result in a shift in species composition and/or production.

The commonly-occurring soils for this site include Corpening, and Rockerville. Corpening occurs in the North and South LRUs. Rockerville occurs throughout the MLRA.

Figure 8. Corpening - MLRA 62 - Shallow Loamy

Table 4. Representative soil features

Parent material	(1) Residuum – limestone and sandstone
Surface texture	(1) Channery loam (2) Channery silt loam (3) Loam
Family particle size	(1) Loamy

Drainage class	Well drained to somewhat excessively drained
Permeability class	Moderate to moderately rapid
Soil depth	30 – 50 cm
Surface fragment cover <=3"	0 – 20 %
Surface fragment cover >3"	0 – 10 %
Available water capacity (0-101.6cm)	2.54 – 7.62 cm
Calcium carbonate equivalent (0-101.6cm)	0 – 30 %
Electrical conductivity (0-101.6cm)	Not specified
Sodium adsorption ratio (0-101.6cm)	Not specified
Soil reaction (1:1 water) (0-101.6cm)	5.6 – 8.4
Subsurface fragment volume <=3" (Depth not specified)	0 – 80 %
Subsurface fragment volume >3" (Depth not specified)	0 – 30 %

Ecological dynamics

The vegetation of the shallow loamy ecological site is part of the Black Hills ecosystem vegetation consisting of a mixture of forests and grasslands resulting from the interaction of a varied topography, geology, climate and natural disturbance. Frequent fires, periodic droughts, and episodic mountain pine beetle infestations are also thought to have contributed to maintain this park-like forest structure of large prairie areas scattered throughout the Black Hills.

Ponderosa pine is a fire adapted species as it has evolved to coexist and flourish with fire; particularly frequent, low-intensity fires that consume mostly small seedlings, prune lower branches from large trees, and reduce woody fuels debris on the forest floor. The result is a

mosaic of conditions ranging from open grassland areas to groups of young seedlings to clumps and groups of large older trees. When large crown fires did occur, they probably did not completely consume all trees within the landscape, but left large healthy trees as sources of seeds to eventually recolonize the burnt areas (Shepperd and Battaglia, 2002). Between 1388 and 1900, fire intervals in the Black Hills ranged from 16 to 20 years (Brown and Hull-Sieg, 1996). The last large-scale recorded fire in the Black Hills was in 2000 (due to arson) but the previous fire was in 1900.

With the advent of fire suppression over the past 120 years, forests density has increased and grassland areas decreased across the whole landscape. Historical records indicated the presence of large areas covered by tall grasses on the outer edges and slopes of the limestone plateau but did not mention ponderosa pine encroachment. McIntosh (1949) conducted botanical field work in the Black hills during the summers of 1924 to 1930 and noted that pines were invading the grassland. Gardner and Thompson (1972) also mentioned that pine encroachment was most evident on areas dominated by warm season grasses. Today, ponderosa pine encroachment is still quite evident on the shallow loamy ecological site where trees seem to germinate and survive well on the stony and gravelly soils in the absence of fire.

The historic native vegetation on this site consisted of mixed prairie grasses occurring on ridges and uplands. The vegetation was predominately warm season grasses mixed with some cool-season grasses portraying a mixed grass prairie characteristic. Little bluestem is the dominant species. Big bluestem and sideoats grama are in association with little bluestem. Subdominant cool season grasses include needle-and-thread and western wheatgrass with prairieJunegrass. Blue grama, hairy grama, and threadleaf sedge are present in the understory. Kentucky bluegrass can be present but does not dominate. Forbs make up a significant proportion in the mixture. Common forbs include cudweed sagewort, scurfpeas, coneflowers, gayfeather, asters along with hood phlox and pussytoes in the understory. Shrubs include fringed sagewort, broom snakeweed, green sagewort, prairie rose, and a few yucca plants scattered on the landscape. A few ponderosa pine and Rocky Mountain juniper trees are often dispersed across the landscape. Bare ground, rocks, and rock fragments are very noticeable across the landscape.

This site developed under Northern Great Plains historic conditions. This consisted of light grazing by ungulates and small mammals, sporadic to routine natural or man-caused fires (of light to moderate intensities) and drought. Changes will occur in the plant communities due to short-term weather variations, impacts of native and/or exotic plants and animals, and management actions. Heavy continuous grazing (stocking levels well above carrying capacity for extended portions of the growing season) without adequate recovery periods cause a departure from the Reference State. Big bluestem becomes reduced or absent from the plant community allowing less palatable, more grazing resistant plants to increase. With continued grazing pressure, non-native plant species such as Kentucky bluegrass, cheat grass, Japanese brome, and smooth brome may begin to invade the site. Continued grazing pressure with stocking rates well above the carrying capacity for the entire growing season could result in an increase in little bluestem (mostly on drier areas with slope). Short-statured grasses and grass-like plants may begin to dominate the site. Shrubs such as soapweed yucca, fringed sagewort, and western snowberry will also increase. If these conditions continue, fine fuels needed for fire will be reduced allowing conifer and shrub encroachment and maintain their dominance unless fire is returned to the site or conifers are removed mechanically.

State and transition model

Figure 9. Shallow Loamy 062XC024SD - LRU-C

Figure 10. Shallow Loamy 062XC024SD - LRU-C

Additional community tables

Table 5. Community 1.1 plant community composition

Group	Common Name	Symbol	Scientific Name	Annual Production ()	Foliar Cover (%)
Grass/Grasslike					
1	Tall and Mid- Warm-Season Grasses			504-1110	
	little bluestem	SCSC	<i>Schizachyrium scoparium</i>	202-706	–
	sideoats grama	BOCU	<i>Bouteloua curtipendula</i>	202-303	–
	prairie dropseed	SPHE	<i>Sporobolus heterolepis</i>	0-101	–
	composite dropseed	SPCOC2	<i>Sporobolus compositus var. compositus</i>	20-101	–
	plains muhly	MUCU3	<i>Muhlenbergia cuspidata</i>	0-101	–
	big bluestem	ANGE	<i>Andropogon gerardii</i>	40-101	–
	marsh muhly	MURA	<i>Muhlenbergia racemosa</i>	0-40	–
2	Cool-Season Bunchgrass			101-303	
	needle and thread	HECOC8	<i>Hesperostipa comata ssp. comata</i>	101-202	–
	prairie Junegrass	KOMA	<i>Koeleria macrantha</i>	20-101	–
	Richardson's needlegrass	ACRI8	<i>Achnatherum richardsonii</i>	0-101	–

	slender wheatgrass	ELTR7	<i>Elymus trachycaulus</i>	40-101	—
	timber oatgrass	DAIN	<i>Danthonia intermedia</i>	0-101	—
	slender wheatgrass	ELTRS	<i>Elymus trachycaulus ssp. subsecundus</i>	40-101	—
	Columbia needlegrass	ACNE9	<i>Achnatherum nelsonii</i>	0-101	—
	green needlegrass	NAVI4	<i>Nassella viridula</i>	20-101	—
	mountain brome	BRMA4	<i>Bromus marginatus</i>	0-101	—
	bluebunch wheatgrass	PSSP6	<i>Pseudoroegneria spicata</i>	0-20	—
3	Cool-Season Rhizomatous Grass			40-202	
	western wheatgrass	PASM	<i>Pascopyrum smithii</i>	40-202	—
	thickspike wheatgrass	ELLAL	<i>Elymus lanceolatus ssp. lanceolatus</i>	0-101	—
4	Short Warm-Season Grasses			40-202	
	blue grama	BOGR2	<i>Bouteloua gracilis</i>	20-202	—
	hairy grama	BOHI2	<i>Bouteloua hirsuta</i>	20-101	—
	threeawn	ARIST	<i>Aristida</i>	0-101	—
5	Other Native Grasses			0-101	
	Grass, perennial	2GP	<i>Grass, perennial</i>	0-101	—
6	Grass-Likes			40-202	
	needleleaf sedge	CADU6	<i>Carex duriuscula</i>	20-101	—
	threadleaf sedge	CAFI	<i>Carex filifolia</i>	20-101	—
	Grass-like (not a true grass)	2GL	<i>Grass-like (not a true grass)</i>	0-101	—
	Richardson's sedge	CARI	<i>Carex richardsonii</i>	0-40	—
	sedge	CAREX	<i>Carex</i>	20-40	—
7	Non-Native Cool-Season Grasses			0	
Forb					
8	Forbs			101-303	
	goldenrod	SOLID	<i>Solidago</i>	20-101	—
	blacksamson echinacea	ECAN2	<i>Echinacea angustifolia</i>	20-101	—
	scurfpea	PSORA2	<i>Psoraleidium</i>	20-101	—
	western yarrow	ACMIO	<i>Achillea millefolium var. occidentalis</i>	20-40	—
	white prairie aster	SYFA	<i>Symphyotrichum falcatum</i>	20-40	—
	wild bergamot	MOFI	<i>Monarda fistulosa</i>	0-40	—
	Forb, perennial	2FP	<i>Forb, perennial</i>	0-40	—
	hairy false goldenaster	HEVI4	<i>Heterotheca villosa</i>	0-40	—
	dotted blazing star	LIPU	<i>Liatris punctata</i>	20-40	—
	fleabane	ERIGE2	<i>Erigeron</i>	0-40	—
	mariposa lily	CALOC	<i>Calochortus</i>	0-40	—
	beardtongue	PENST	<i>Penstemon</i>	0-40	—
	prairie clover	DALEA	<i>Dalea</i>	20-40	—
	upright prairie coneflower	RACO3	<i>Ratibida columnifera</i>	20-40	—
	old man's whiskers	GETR	<i>Geum triflorum</i>	0-40	—
	American vetch	VIAM	<i>Vicia americana</i>	0-40	—
	blanketflower	GAAR	<i>Gaillardia aristata</i>	0-40	—
	Lewis flax	LILE3	<i>Linum lewisii</i>	0-40	—
	bluebell bellflower	CARO2	<i>Campanula rotundifolia</i>	0-40	—
	cinquefoil	POTEN	<i>Potentilla</i>	0-40	—
	white sagebrush	ARLU	<i>Artemisia ludoviciana</i>	20-40	—

	deathcamas	ZIGAD	<i>Zigadenus</i>	0-20	–
	downy paintedcup	CASE5	<i>Castilleja sessiliflora</i>	0-20	–
	spiny phlox	PHHO	<i>Phlox hoodii</i>	0-20	–
	common starlily	LEMO4	<i>Leucocrinum montanum</i>	0-20	–
Shrub/Vine					
9	Shrubs			20-202	
	creeping juniper	JUHO2	<i>Juniperus horizontalis</i>	20-101	–
	prairie sagewort	ARFR4	<i>Artemisia frigida</i>	20-101	–
	field sagewort	ARCA12	<i>Artemisia campestris</i>	20-40	–
	skunkbush sumac	RHTR	<i>Rhus trilobata</i>	0-40	–
	smooth sumac	RHGL	<i>Rhus glabra</i>	0-40	–
	soapweed yucca	YUGL	<i>Yucca glauca</i>	0-40	–
	Shrub (>.5m)	2SHRUB	<i>Shrub (>.5m)</i>	0-40	–
	common juniper	JUCO6	<i>Juniperus communis</i>	0-40	–
	plains pricklypear	OPPO	<i>Opuntia polyacantha</i>	0-20	–
Tree					
10	Trees			0-20	
	ponderosa pine	PIPO	<i>Pinus ponderosa</i>	0-20	–
	Rocky Mountain juniper	JUSC2	<i>Juniperus scopulorum</i>	0-20	–

Table 6. Community 1.2 plant community composition

Group	Common Name	Symbol	Scientific Name	Annual Production ()	Foliar Cover (%)
-------	-------------	--------	-----------------	----------------------	------------------

Table 7. Community 2.1 plant community composition

Group	Common Name	Symbol	Scientific Name	Annual Production ()	Foliar Cover (%)
-------	-------------	--------	-----------------	----------------------	------------------

Table 8. Community 3.1 plant community composition

Group	Common Name	Symbol	Scientific Name	Annual Production ()	Foliar Cover (%)
-------	-------------	--------	-----------------	----------------------	------------------

Recreational uses

In this area, there is wondrous scenery in any season and it almost has something for almost everyone for outdoor recreation activities. Despite that they may cause some ecological disturbances to the sites, visitors provide a good income to the small communities across the area. Opportunities for hiking, mountain biking, snowmobiling, and horse riding exist through a network of trails covering several hundreds of miles. The presence of two adjacent parks in the area (Custer State Park and Wind Cave National Park) brings in a large flow of resident and non-resident tourists which visit the area for its natural beauty and for the viewing of many wildlife species particularly buffalo, bighorn sheep, and mountain goats. Fishing and hunting attract a great number of sports men and women for trout fishing, turkey hunting, and big game hunting of elk, deer, and antelope. Wild cats such as mountain lion are hunted during the fall season. All of these recreation activities combined with the artistry business attracts a great number of visitors and brings in a substantial income to the small towns across the areas.

Wood products

No appreciable wood products are available in this site.

Other information

Revision Notes: "This PROVISIONAL ecological site concept has been QC'd and QA'd to ensure that the site meets the NESH standards for a provisional ecological site that provides basic compiled information in one location. This site should not be considered an Approved ESD, as it is only the foundational site concepts and requires further data collection—specifically high-intensity data characterizations and full 232 soil descriptions—and further site investigations and final STM reviews before it can be used as an Approved ESD meeting NESH standards." Site Development and Testing Plan: Future work, as described in a Project Plan, to validate the information in this Provisional Ecological Site Description is needed. This will include field activities to collect low, medium and high intensity sampling, soil correlations, and analysis of that data. Annual field reviews should be done by soil scientists and vegetation specialists. A final field

review, peer review, quality control, and quality assurance reviews of the ESD will be needed to produce the final document.

Inventory data references

Information presented here has been derived from NRCS clipping data and other inventory data. Field observations from range-trained personnel were also used. Those involved in developing this site include: Lakhdar Benkobi, ESI/SRIC, NRCS; Stan Boltz, Range Management Specialist, NRCS; Dan Brady, Soil Scientist, NRCS; Mitch Faulkner, Range Management Specialist, NRCS; Roger Gates, Associate Professor/Rangeland Management Specialist, West River Ag Center; Rick Peterson, Ecological Site Specialist, NRCS; Matthew, Scott, Botanist - USFS Hell Canyon District Ranger; L. Michael Stirling, Range Management Specialist, NRCS; Jim Westerman, Soil Scientist, NRCS. All inventory information and data records are compiled within the Rapid City, SD USDA-NRCS Shared "S" network drive.

Other references

Brown, P. M. and C. Hull-Sieg. 1996. Fire History in Interior Ponderosa Pine Communities of the Black Hills, South Dakota, USA, *Int. J. Wildland Fire* 6(3): 97-105.

Gartner F. R. and W. W. Thompson. 1972. Fire in the Black Hills forest-grass ecotone, South Dakota Agricultural Experiment Station, Journal Series No 1115.

High Plains Regional Climate Center, University of Nebraska, 830728 Chase Hall, Lincoln, NE 68583-0728. (<http://www.hprcc.unl.edu/>)

McIntosh, A. C. 1949. A botanical survey of the Black Hills of South Dakota. *Black Hills Engineer*. 28 (4): 3-75.

Shepperd, W. D. and M. A. Battaglia. 2002. Ecology, silviculture, and management of Black Hills ponderosa pine. Gen. Tech. Rep. RMRS-GTR-97. Fort Collins, CO: US Department of Agriculture, Forest Service, Rocky Mountain Research Station. 112 p.

USDA, NRCS. National Water and Climate Center,

Toledo, D., Sanderson, M., Spaeth, K., Hendrickson, J., Printz, J. 2014. Extent of Kentucky Bluegrass and Its Effect on Native Plant Species Diversity and Ecosystem Services in the Northern Great Plains of the United States. *Invasive Plant Science and Management*. 7(4):543-522. Weed Science Society of America.

101 SW Main, Suite 1600, Portland, OR 97204-3224. (<http://wcc.nrcs.usda.gov>)

USDA, NRCS. National Range and Pasture Handbook, September 1997

USDA, NRCS. National Soil Information System, Information Technology Center, 2150 Centre Avenue, Building A, Fort Collins, CO 80526. (<http://nasis.nrcs.usda.gov>)

USDA, NRCS. 2001. The PLANTS Database, Version 3.1 (<http://plants.usda.gov>). National Plant Data Center, Baton Rouge, LA 70874-4490 USA.

USDA, NRCS. National Ecological Site Handbook, 1st Ed, January 2014

Wrage, K. J. 1994. The effects of ponderosa pine on soil moisture, precipitation, and understory vegetation in the Black Hills of South Dakota. 158 p. Thesis.

Contributors

Westerman Jim
Benkobi Lakhdar
Faulkner Mitch
Peterson Rick L.

Approval

Suzanne Mayne-Kinney, 7/31/2024

Acknowledgments

All ecological sites were then reviewed and approved at the Provisional Level by David Kraft, Regional ESS, Salina, KS.

Rangeland health reference sheet

Interpreting Indicators of Rangeland Health is a qualitative assessment protocol used to determine ecosystem condition based on benchmark characteristics described in the Reference Sheet. A suite of 17 (or more) indicators are typically considered in an assessment. The ecological site(s) representative of an assessment location must be known prior to applying the protocol and must be verified based on soils and climate. Current plant community cannot be used to identify the ecological site.

Author(s)/participant(s)	
Contact for lead author	
Date	07/12/2026
Approved by	
Approval date	
Composition (Indicators 10 and 12) based on	Annual Production

Indicators

1. Number and extent of rills:

2. Presence of water flow patterns:

3. Number and height of erosional pedestals or terracettes:

4. Bare ground from Ecological Site Description or other studies (rock, litter, lichen, moss, plant canopy are not bare ground):

5. Number of gullies and erosion associated with gullies:

6. Extent of wind scoured, blowouts and/or depositional areas:

7. Amount of litter movement (describe size and distance expected to travel):

8. Soil surface (top few mm) resistance to erosion (stability values are averages - most sites will show a range of values):

9. Soil surface structure and SOM content (include type of structure and A-horizon color and thickness):

10. Effect of community phase composition (relative proportion of different functional groups) and spatial distribution on infiltration and runoff:

11. Presence and thickness of compaction layer (usually none; describe soil profile features which may be mistaken for compaction on this site):

12. Functional/Structural Groups (list in order of descending dominance by above-ground annual-production or live foliar cover using symbols: >>, >, = to indicate much greater than, greater than, and equal to):

Dominant:

Sub-dominant:

Other:

Additional:

13. Amount of plant mortality and decadence (include which functional groups are expected to show mortality or decadence):

14. Average percent litter cover (%) and depth (in):

15. Expected annual annual-production (this is TOTAL above-ground annual-production, not just forage annual-production):

16. Potential invasive (including noxious) species (native and non-native). List species which BOTH characterize degraded states and have the potential to become a dominant or co-dominant species on the ecological site if their future establishment and growth is not actively controlled by management interventions. Species that become dominant for only one to several years (e.g., short-term response to drought or wildfire) are not invasive plants. Note that unlike other indicators, we are describing what is NOT expected in the reference state for the ecological site:

17. Perennial plant reproductive capability:
