

Ecological site R067BY038CO

Wet Meadow

Last updated: 12/05/2024
Accessed: 09/26/2026

General information

Provisional. A provisional ecological site description has undergone quality control and quality assurance review. It contains a working state and transition model and enough information to identify the ecological site.

MLRA notes

Major Land Resource Area (MLRA): 067B—Central High Plains, Southern Part

MLRA 67B occurs in eastern Colorado and consists of rolling plains and river valleys. Some canyonlands occur in the southeast portion. The major rivers are the South Platte and Arkansas which flow from the Rocky Mountains to Nebraska and Kansas. Other rivers in the MLRA include the Cache la Poudre and Republican and associated tributaries. This MLRA is traversed by Interstate 25, 70 and 76; and U.S. Highways 50 and 287. Major land uses include 54 percent rangeland, 35 percent cropland, and 2 percent pasture and hayland. Urban, developed open space, and miscellaneous land occupy approximately 9 percent. Major Cities in this area include Fort Collins, Greeley, Sterling, and Denver. Other cities include Limon, Cheyenne Wells, and Springfield. Land ownership is mostly private. Federal lands include Pawnee and Comanche National Grasslands (U.S. Forest Service), Sand Creek Massacre National Historic Site (National Park Service), and Rocky Mountain Arsenal National Wildlife Refuge (U.S. Fish & Wildlife Service). State Parks include Cherry Creek and Chatfield Reservoirs, and Barr and Jackson Lakes. This region is periodically affected by severe drought, including the historic “Dust Bowl” of the 1930s. Dust storms may form during drought years in windy periods. Elevations range from 3,400 to 6,000 feet. The Average annual precipitation ranges from 14 to 17 inches per year and ranges from 13 inches to over 18 inches, depending upon location. Precipitation occurs mostly during the growing season, often during rapidly developing thunderstorms. Mean annual air temperature (MAAT) is 48 to 52 degrees Fahrenheit. Summer temperatures may exceed 100 degrees Fahrenheit. Winter temperatures may be sub-zero, and snowfall varies from 20 to 40 inches per year. Snow cover frequently melts between snow events.

LRU notes

Land Resource Unit (LRU) A is the northeast portion of MLRA 67B, to an extent of approximately 9 million acres. Most of the LRU is rangeland, and includes the Pawnee National Grassland. Dryland winter wheat/fallow rotations (that may include dryland corn, sunflowers, and sorghum) are grown in most counties. Irrigated cropland is utilized in the South Platte Valley. Small acreage and urban ownership are more concentrated on the Front Range. This LRU is found in portions of Adams, Arapahoe, Elbert, Kit Carson, Larimer, Lincoln, Logan, Washington, and Weld counties. Other counties include Boulder, Cheyenne, Denver, Jefferson, and Yuma. The soil moisture regime is aridic ustic. The mean annual air temperature (MAAT) is 50 degrees Fahrenheit. LRU B is in the southeast portion of MLRA 67B (2.6 million acres) and includes portions of Baca, Bent, Cheyenne, Kiowa, Las Animas, and Prowers counties. Most of the LRU remains in rangeland and includes the Comanche National Grassland. On the farmed land, a system of dryland winter wheat/fallow rotations (that may include dryland corn, sunflowers, and sorghum) is implemented. Irrigated cropland is found in the Arkansas Valley. The soil moisture regime is aridic ustic and the MAAT is 52 degrees Fahrenheit. LRU C occurs in portions of Morgan and Weld counties (approximately 1.2 million acres). Most of LRU C is in rangeland. On the farmed land, a system of dryland winter wheat/fallow rotations (that may include dryland corn, sunflowers, and sorghum) is implemented. The soil moisture regime is ustic aridic and the MAAT is 48 degrees Fahrenheit.

Classification relationships

MLRA 67B is in the Colorado Piedmont and Raton Sections of the Great Plains Province (USDA, 2006). The MLRA is further defined by Land Resource Units (LRUs) A, B, and C. Features such as climate, geology, landforms, and key vegetation further refine these concepts and are described in other sections of the Ecological Site Description (ESD). NOTE: To date, these LRUs are DRAFT. Relationship to Other Hierarchical Classifications: NRCS Classification Hierarchy: Physiographic Division, Physiographic Province, Physiographic Section, Land Resource Region, Major Land Resource Area, Land Resource Unit (Fenneman, 1946). USFS Classification Hierarchy: Domain, Division, Province, Section, Subsection, Land Type Association: Land Type, Land Type Phase (Cleland et al, 1997). REVISION NOTES: The Wet Meadow Ecological Site was developed by an earlier version (2004, revised 2007). This earlier version was based on input from Natural Resources Conservation Service (formerly Soil Conservation Service) and historical information obtained from the Wet Meadow Range Site descriptions (1975). This ESD meets the Provisional requirements of the National Ecological Site Handbook (NESH). This ESD will continue refinement towards an Approved status according to the NESH.

Ecological site concept

This site is a run-on site with the soil surface within four feet of the water table, and will often have redoximorphic features in the upper portion of the soil profile. There are no visible salts on the surface or within the soil profile and it does not have sandy subsoil textures.

Associated sites

R067BY073CO	Riparian This ecological site is commonly adjacent.
R067BY029CO	Sandy Meadow This ecological site is commonly adjacent.
R067BY031CO	Sandy Bottomland This ecological site is commonly adjacent.
R067BY035CO	Salt Meadow This ecological site is commonly adjacent.

Similar sites

R067BY035CO	Salt Meadow The Salt Meadow Ecological Site has visible salts on the surface or in the soil profile.
--------------------	--

Table 1. Dominant plant species

Tree	Not specified
Shrub	(1) <i>Amorpha fruticosa</i>
Herbaceous	(1) <i>Andropogon gerardii</i> (2) <i>Sorghastrum nutans</i>

Physiographic features

This site occurs on floodplains of the river valleys and drainageways on the plains. There is an influential water table associated with this site.

Table 2. Representative physiographic features

Landforms	(1) Flood plain (2) Drainageway
Runoff class	Negligible to low

Flooding duration	Very brief (4 to 48 hours) to brief (2 to 7 days)
Flooding frequency	Occasional to frequent
Ponding frequency	None
Elevation	1,100 – 1,710 m
Slope	0 %
Ponding depth	0 cm
Water table depth	20 – 60 cm
Aspect	Aspect is not a significant factor

Climatic features

Average annual precipitation across the MLRA extent is 14 to 17 inches, and ranges from 13 to over 18 inches, depending on location. Precipitation increases from north to south. Mean Annual Air Temperature (MAAT) is 50 degrees Fahrenheit in the northern part and increases to 52 degrees Fahrenheit in the southern part. Portions of Morgan and Weld counties are cooler and drier, the MAAT is 48 degrees Fahrenheit, and average precipitation is 13 to 14 inches per year.

Two-thirds of the annual precipitation occurs during the growing season from mid-April to late September. Snowfall averages 30 inches per year, area-wide, but varies by location from 20 to 40 inches per year. Winds are estimated to average 9 miles per hour annually. Daytime winds are generally stronger than at night, and occasional strong storms may bring periods of high winds with gusts to more than 90 mph. High-intensity afternoon thunderstorms may arise. The average length of the freeze-free period (28 degrees Fahrenheit) is 155 days from April 30th to October 3rd. The average frost-free period (32 degrees Fahrenheit) is 136 days from May 11th to September 24th. July is the hottest month, and December and January are the coldest months. Summer temperatures average 90 degrees Fahrenheit and occasionally exceed 100 degrees Fahrenheit. Summer humidity is low and evaporation is high. Winters are characterized with frequent northerly winds, producing severe cold with temperatures occasionally dropping to -30 degrees Fahrenheit or lower. Blizzard conditions may form quickly. For detailed information, visit the Western Regional Climate Center website:

Western Regional Climate Center Historical Data Western U.S. Climate summaries, NOAA Coop Stations Colorado <http://www.wrcc.dri.edu/summary/Climsmco.html>.

Table 3 Representative climatic features

Frost-free period (characteristic range)	120-130 days
Freeze-free period (characteristic range)	130-150 days
Precipitation total (characteristic range)	360-430 mm
Frost-free period (actual range)	100-130 days
Freeze-free period (actual range)	130-160 days

Precipitation total (actual range)	360-430 mm
Frost-free period (average)	120 days
Freeze-free period (average)	140 days
Precipitation total (average)	380 mm

- (1) BRIGGSDALE [USC00050945], Briggsdale, CO
- (2) BYERS 5 ENE [USC00051179], Byers, CO
- (3) GREELEY UNC [USC00053553], Greeley, CO
- (4) NUNN [USC00056023], Nunn, CO
- (5) BRIGHTON 3 SE [USC00050950], Brighton, CO
- (6) CHEYENNE WELLS [USC00051564], Cheyenne Wells, CO
- (7) FLAGLER 1S [USC00052932], Flagler, CO
- (8) KIT CARSON [USC00054603], Kit Carson, CO
- (9) FT MORGAN [USC00053038], Fort Morgan, CO
- (10) SPRINGFIELD 7 WSW [USC00057866], Springfield, CO
- (11) LIMON WSMO [USW00093010], Limon, CO

Influencing water features

There is a seasonal or perennial water table that influences the kinds and amounts of vegetation on this site. The variability in water table provides a mosaic of drier and wetter areas on a meadow. This provides a diversity of non-hydrophytic and hydrophytic vegetation on the site. The water table in some areas is artificially induced, caused by seepage from nearby irrigation ditches, canals, and reservoirs.

Note: This is a general overview for the site concept, and is not a wetland determination.

Soil features

The soils on this site are very deep, very poorly to somewhat poorly drained soils that formed from alluvium from mixed sources. They typically have a moderate to moderately rapid permeability class, but may range to moderately slow. The soil moisture regime is typically aquic. The soil temperature regime is mesic.

The surface layer of the soils in this site are typically sandy loam or loam, but may be variable in some places. The surface layer ranges from 3 to 18 inches thick. The subsoil is typically stratified, with textures ranging from sandy loam, loamy sand, sand, sandy clay loam, clay loam, or silt loam. Rock fragments range from 0 to 40 percent in the underlying material. Soils in this site are typically leached of carbonates at the surface; some soils may have carbonates at the surface and throughout the profile. These soils are typically not susceptible to erosion by water and wind due to the wetness of the soil profile by the seasonal water table. However, these areas may be prone to wind erosion if these areas are drained and the surface is not protected by vegetation.

Major soil series correlated to this ecological site include: Edgewater, Las Animas, Seldom.

Other soil series that have been correlated to this site, but may eventually be re-correlated include: Aquepts, Bijou wet*, Fluvaquentic Haplaquolls, Haplaquolls, Kitcarson, and Wet Alluvial Land.

*Bijou is typically found on terraces or uplands and does not have a water table.

The attributes listed below represent 0-40 inches in depth or to the first restrictive layer.

Note: Revisions to soil surveys are on-going. For the most recent updates, visit the Web Soil Survey, the official site for soils information: <http://websoilsurvey.nrcs.usda.gov/app/WebSoilSurvey.aspx>.

The attributes listed below represent 0-40 inches in depth or to the first restrictive layer.

Table 4. Representative soil features

Parent material	(1) Alluvium
Surface texture	(1) Sandy loam (2) Loam
Drainage class	Very poorly drained to somewhat poorly drained
Permeability class	Moderately slow to moderately rapid
Soil depth	200 cm
Surface fragment cover <=3"	Not specified
Surface fragment cover >3"	Not specified
Available water capacity (0-101.6cm)	5.08 – 20.32 cm
Calcium carbonate equivalent (0-101.6cm)	0 – 20 %
Electrical conductivity (0-101.6cm)	0 – 10 mmhos/cm
Sodium adsorption ratio (0-101.6cm)	0 – 10
Soil reaction (1:1 water) (0-101.6cm)	6.6 – 8.4
Subsurface fragment volume <=3" (Depth not specified)	0 – 40 %
Subsurface fragment volume >3" (Depth not specified)	Not specified

Ecological dynamics

The Wet Meadow Ecological Site is characterized by three states: Reference, Cool-Season Dominant, and Increased Bare Ground. The Reference State is characterized by warm-season tall bunchgrass (big bluestem, switchgrass, Indiangrass), warm-season tall rhizomatous grass (prairie cordgrass), and cool-season grasslike (Nebraska sedge). The Cool-season Dominant State is characterized by cool-season grasslikes (mountain rush, Baltic rush), and bunchgrass (foxtail barley). The Increased Bare Ground State is characterized by early successional cool-season grass (foxtail barley), introduced cool-season rhizomatous grass (Kentucky bluegrass), annual grasses, and annual forbs.

The site is subirrigated (i.e. high water table) throughout the growing season. The availability of water has a major influence on the vegetation that will persist on this site.

Today, this is an important site for livestock grazing, especially beef cattle. The management of livestock grazing, coupled with annual climatic variations, has a major influence on the plant communities of this site. Continuous, heavy grazing without adequate recovery periods following each grazing occurrence causes prairie cordgrass, Indiangrass, big bluestem, switchgrass, Nebraska sedge, and false indigo bush to decrease in frequency and production. Kentucky bluegrass, mountain rush, Baltic rush, scouring rush, and other various grass-likes increase forming a cool-season dominated plant community. Heavy, continuous grazing or excessive defoliation ultimately results in a plant community dominated by foxtail barley and annuals. Excessive litter, decadence, and plant mortality result from the lack of fire or non-use. Extended periods of non-use, lack of fire, or heavy, long term continuous grazing leads to increased bare areas.

The degree of grazing has a significant impact on the ecological dynamics of the site. This region was historically occupied by large grazing animals, such as bison, elk, pronghorn, and mule deer. Grazing by these large herbivores, along with climatic and seasonal weather fluctuations, had a major influence on the ecological dynamics of the site. Deer and pronghorn are widely distributed throughout the MLRA. Secondary influences of herbivory by species such as prairie dogs and other small rodents, insects, and root-feeding organisms continues to impact the vegetation.

Historically, grazing patterns by herds of large ungulates were driven by water distribution, precipitation events, drought events, and fire. It is believed that grazing periods would have been shorter, followed by longer recovery periods. These large migrating herds impacted the ecological processes of nutrient and hydrologic cycles, by urination, trampling (incorporation of litter into the soil surface), and breaking of surface crust, (which increases water infiltration).

Today, livestock grazing, especially beef cattle has been a major influence on the ecological dynamics of the site. Grazing management, coupled with the effects of annual climatic variations, largely dictates the plant communities for the site.

Recurrent drought has historically impacted the vegetation of this region. Changes in species composition vary depending upon the duration and severity of the drought cycle and prior grazing management. Drought events since 2002 have significantly increased mortality of blue grama and buffalograss in some locales.

This site developed with occasional fire as part of the ecological processes. Historic fire frequency (pre-industrial) is estimated at 10 to14 years (Guyette, 2012), randomly distributed, and started by lightning at various times throughout the growing season. Early human inhabitants also were likely to start fires for various reasons (deliberate or accidental). It is believed that fires were set as a management tool for attracting herds of large migratory herbivores (Stewart, 2002). The impact of fire over the past 100 years has been relatively insignificant due to the human control of wildfires and the lack of acceptance of prescribed fire as a management tool.

Mechanical treatment consisting of contour pitting, furrowing, terracing, chiseling, and disking has been practiced in the past. It was theorized that the use of this high-input technology would improve production and plant composition on rangeland. These high-cost practices have shown to have no significant long-term benefits on production or plant composition and have only resulted in a permanently rough ground surface. Prescribed grazing that mimics the historic grazing of herds of migratory herbivores, as described earlier, has been shown to result in desired improvements based on management goals for this ecological site.

Eastern Colorado was strongly affected by extended drought conditions in the “Dust Bowl” period of the 1930’s, with recurrent drought cycles in the 1950s and 1970s. Extreme to exceptional drought conditions have re-visited the area from 2002 to 2012, with brief interludes of near normal to normal precipitation years. Long-term effects of these latest drought events have yet to be determined. Growth of native cool-season plants begins about April 1 and continues to mid-June. Native warm-season plants begin growth about May 1 and continue to about August 15. Regrowth of cool-season plants occurs in September in most years, depending on the availability of moisture.

State and transition model

Additional community tables

Table 5. Community 1.1 plant community composition

Group	Common Name	Symbol	Scientific Name	Annual Production ()	Foliar Cover (%)
Grass/Grasslike					

1				3811-4259	
	big bluestem	ANGE	<i>Andropogon gerardii</i>	673-1121	–
	Indiangrass	SONU2	<i>Sorghastrum nutans</i>	673-1121	–
	prairie cordgrass	SPPE	<i>Spartina pectinata</i>	673-1121	–
	switchgrass	PAVI2	<i>Panicum virgatum</i>	673-897	–
	Nebraska sedge	CANE2	<i>Carex nebrascensis</i>	224-448	–
	western wheatgrass	PASM	<i>Pascopyrum smithii</i>	224-448	–
	Canada wildrye	ELCA4	<i>Elymus canadensis</i>	90-224	–
	Grass-like, perennial	2GLP	<i>Grass-like, perennial</i>	90-224	–
	Grass, perennial	2GP	<i>Grass, perennial</i>	45-224	–
	slender wheatgrass	ELTR7	<i>Elymus trachycaulus</i>	45-135	–
	bulrush	SCIRP	<i>Scirpus</i>	45-90	–
	little bluestem	SCSC	<i>Schizachyrium scoparium</i>	45-90	–
	mountain rush	JUARL	<i>Juncus arcticus ssp. littoralis</i>	45-90	–
	vine mesquite	PAOB	<i>Panicum obtusum</i>	0-45	–
	smooth horsetail	EQLA	<i>Equisetum laevigatum</i>	0-45	–
	American sloughgrass	BESY	<i>Beckmannia syzigachne</i>	0-45	–
Forb					
2	Forbs			224-448	
	Forb, perennial	2FP	<i>Forb, perennial</i>	90-224	–
	showy prairie gentian	EUEXR	<i>Eustoma exaltatum ssp. russellianum</i>	45-224	–
	Colorado butterfly plant	GANEC	<i>Gaura neomexicana ssp. coloradensis</i>	45-90	–
	American licorice	GLLE3	<i>Glycyrrhiza lepidota</i>	45-90	–
	Maximilian sunflower	HEMA2	<i>Helianthus maximiliani</i>	0-45	–
	Colorado butterfly plant	OECOC	<i>Oenothera coloradensis ssp. coloradensis</i>	0-45	–
	Pennsylvania smartweed	POPE2	<i>Polygonum pensylvanicum</i>	0-45	–
	giant goldenrod	SOGI	<i>Solidago gigantea</i>	0-45	–
	white heath aster	SYER	<i>Symphyotrichum ericoides</i>	0-45	–
	swamp milkweed	ASIN	<i>Asclepias incarnata</i>	0-45	–
	threelobe beggarticks	BITR	<i>Bidens tripartita</i>	0-45	–
	false boneset	BREU	<i>Brickellia eupatorioides</i>	0-45	–
Shrub/Vine					
3				45-224	
	false indigo bush	AMFR	<i>Amorpha fruticosa</i>	45-90	–
	Woods' rose	ROWO	<i>Rosa woodsii</i>	0-45	–
	western snowberry	SYOC	<i>Symphoricarpos occidentalis</i>	0-45	–
	Shrub (>.5m)	2SHRUB	<i>Shrub (>.5m)</i>	0-45	–

Table 6. Community 1.2 plant community composition

Group	Common Name	Symbol	Scientific Name	Annual Production ()	Foliar Cover (%)
-------	-------------	--------	-----------------	----------------------	------------------

Table 7. Community 1.3 plant community composition

Group	Common Name	Symbol	Scientific Name	Annual Production ()	Foliar Cover (%)
-------	-------------	--------	-----------------	----------------------	------------------

Table 8. Community 2.1 plant community composition

Group	Common Name	Symbol	Scientific Name	Annual Production ()	Foliar Cover (%)
-------	-------------	--------	-----------------	----------------------	------------------

Table 9. Community 3.1 plant community composition

Group	Common Name	Symbol	Scientific Name	Annual Production ()	Foliar Cover (%)
-------	-------------	--------	-----------------	----------------------	------------------

Animal community

WILDLIFE INTERPRETATIONS: Wet Meadow sites support a unique suite of wildlife species due to their association with river and stream systems. Riparian corridors generally represent areas of increased biodiversity compared to adjacent upland sites, and these sites often occur in those riparian systems. Wet meadows provide habitat components for white-tailed and mule deer, wild turkey, and bobwhite quail. These areas can also have increased diversity of herpetile species. 1.1 Reference Plant Community The Reference Plant Community provides important habitat components for white-tailed and mule deer, including foraging, bedding, and fawning areas. Wild turkey hens use shrubs for nesting and brood-rearing, and all turkeys use these sites incidentally the rest of the year. The taller grasses, especially switchgrass and prairie cordgrass, provide important nesting habitats for northern bobwhite quail, and snowberry provides the highest quality loafing and escape habitat for this species on this site. Plains and common garter snakes may be found in this plant community, especially if low areas with seasonal water are present. 1.2 Community The 1.2 Community has reduced tall grasses and shrubs which degrades the overall quality of the site for wildlife. White-tailed and mule deer may move through this community and feed to some extent, but with reduced cover the value for bedding and fawning is also reduced. Northern bobwhite quail use of this community is reduced as well as the tallgrass and shrub components declines. Quail will be absent from sites where these components are entirely absent. 1.3 Community The 1.3 community has greatly reduced value for wildlife due to the loss of tall grasses and the reduced abundance of shrubs. White-tailed and mule deer may move through this community and feed to some extent, but with reduced cover the value for bedding and fawning is also reduced. Northern bobwhite quail use of this community is reduced as the tallgrass and shrub components decline, and because the litter layer is too thick and dense for quail to navigate. 2.1 Community The 2.1 Community on this site represents a lower amount of wildlife diversity. White-tailed and mule deer may move through this community but spend little time feeding or bedding. Wild turkey may use the edges of these meadows in the spring for breeding displays but nesting and brood-rearing value has been lost. 3.1 Community Due to the greatly reduced plant diversity of this site and impairment to nutrient cycling and water infiltration processes, the wildlife community that uses this site is also greatly reduced. **GRAZING INTERPRETATIONS:** The following table lists suggested initial stocking rates for an animal unit (1000-pound beef cow) under continuous grazing (yearlong grazing or growing-season-long grazing) based on normal growing conditions. However, continuous grazing is not recommended. These estimates should only be used as preliminary guidelines in the initial stages of the conservation planning process. Often, the existing plant composition does not entirely match any particular plant community described in this ecological site description. Therefore, field inventories are always recommended to document plant composition, total production, and palatable forage production. Carrying capacity estimates that reflect on-site conditions should be calculated using field inventories. If the following production estimates are used, they should be adjusted based on animal kind or class and on the specific palatability of the forage plants in the various plant community descriptions. Under a properly stocked, properly applied, prescribed grazing management system that provides adequate recovery periods following each grazing event, improved harvest efficiencies eventually result in increased carrying capacity. See USDA-NRCS Colorado Prescribed Grazing Standard and Specification Guide (528). The stocking rate calculations are based on the total annual forage production in a normal year multiplied by 25 percent harvest efficiency divided by 912.5 pounds of ingested air-dry vegetation for an animal unit per month (AUM). Reference PC - (4,000) (1.10) 1.2 PC - (2,500) (0.68) 2.1 PC - (1,300) (0.36) Grazing by domestic livestock is one of the major income-producing industries in the area. Rangelands in this area provide yearlong forage under prescribed grazing for cattle, sheep, horses, and other herbivores. An on-site inventory is required prior to developing a grazing plan.

Hydrological functions

Water is the principal factor limiting forage production on this site. This site is dominated by soils in hydrologic group A and B. Infiltration is moderate and runoff potential for this site varies from moderately poor to poor depending on ground cover. Areas where ground cover is less than 50 percent have the greatest potential to have reduced infiltration and higher runoff (refer to NRCS Section 4, National Engineering Handbook (USDA–NRCS, 1972–2012) for runoff quantities and hydrologic curves).

Recreational uses

This site provides hunting, hiking, photography, bird watching, and other opportunities. The wide varieties of plants that bloom from spring until fall have an aesthetic value that appeals to visitors.

Wood products

No appreciable wood products are present on the site.

Other products

Site Development and Testing Plan General Data (MLRA and Revision Notes, Hierarchical Classification, Ecological Site Concept, Physiographic, Climate, and Water Features, and Soils Data): Updated. All "Required" items complete to Provisional level. Community Phase Data (Ecological Dynamics, STM, Transition & Recovery Pathways, Reference Plant Community, Species Composition List, Annual Production Table): Updated. All "Required" items complete to Provisional level. NOTE: Annual Production Table is from the "Previously Approved" ESD 2004. The Species Composition List is also from the 2004 version, with minor edits. These will need review

for future updates at Approved level. Each Alternative State/Community: Complete to Provisional level Supporting Information (Site Interpretations, Assoc. & Similar Sites, Inventory Data References, Agency/State Correlation, References): Updated. All "Required" items complete to Provisional level. Livestock Interpretations updated to reflect Total Annual Production revisions in each plant community. Wildlife interpretations, general narrative, and individual plant communities updated to the Provisional level. Hydrology, Recreational Uses, Wood Products, Other Products, Plant Preferences table, and Rangeland Health Reference Sheet carried over from previously "Approved" ESD 2004. Reference Sheet The Reference Sheet was previously approved in 2007. It will be updated at the next "Approved" level. "Future work, as described in a project plan, to validate the information in this provisional ecological site description is needed. This will include field activities to collect low and medium intensity sampling, soil correlations, and analysis of that data. Annual field reviews should be done by soil scientists and vegetation specialists. A final field review, peer review, quality control, and quality assurance reviews of the ESD will be needed to produce the final document." (NI 430_306 ESI and ESD, April, 2015).

Other information

Relationship to Other Hierarchical Classifications: NRCS Classification Hierarchy: Physiographic Divisions of the United States (Fenneman, 1946): Physiographic DivisionPhysiographic ProvincePhysiographic SectionLand Resource RegionMajor Land Resource Area (MLRA)Land Resource Unit (LRU). USFS Classification Hierarchy: National Hierarchical Framework of Ecological Units (Cleland et al, 181-200): DomainDivisionProvinceSectionSubsectionLandtype AssociationLandtypeLandtype Phase.

Inventory data references

NRI: references to Natural Resource Inventory data Information presented here has been derived from data collection on private and federal lands using: • Double Sampling (clipped 2 of 5 plots)* • Rangeland Health (Pellant et al., 2005) • Soil Stability (Pellant et al., 2005) • Line Point Intercept : Foliar canopy, basal cover (Forb, Graminoid, Shrub, subshrub, Lichen, Moss, Rock fragments, bare ground, % Litter) (Herrick et al., 2005) • Soil pedon descriptions collected on site (Schoeneberger et al., 2012) *NRCS double-sampling method, CO NRCS Similarity Index Worksheet 528(1). Additional reconnaissance data collection using numerous ocular estimates and other inventory data; NRCS clipping data for USDA program support; Field observations from experienced range trained personnel. Specific data information is contained in individual landowner/user case files and other files located in county NRCS field offices. Those involved in developing the 2004 site description include: Harvey Sprock, Rangeland Management Specialist, CO-NRCS; Ben Berlinger, Rangeland Management Specialist, CO-NRCS; James Borchert, Soil Scientist, CO-NRCS; Terri Skadeland, Biologist, CO-NRCS.

References

- Guyette, R.P., M.C. Stambaugh, D.C. Dey, and R. Muzika. 2012. **Predicting Fire Frequency with Chemistry and Climate**. Ecosystems 15:322–335.
- Stewart, O.C., H.T. Lewis, and M.K. Anderson. 2002. **Forgotten Fires: Native Americans and the Transient Wilderness**. University of Oklahoma Press, Norman, OK. 351p.

Other references

Data collection for this ecological site was done in conjunction with the progressive soil surveys within the 67B Central High Plains (Southern Part) of Colorado. It has been mapped and correlated with soils in the following soil surveys: Adams County, Arapahoe County, Baca County, Bent County, Boulder County, Cheyenne County, El Paso County Area, Elbert County, Eastern Part, Kiowa County, Kit Carson County, Larimer County Area, Las Animas County Area, Lincoln County, Logan County, Morgan County, Prowers County, Washington County, Weld County, Northern Part, and Weld County, Southern Part.

30 Year Climatic and Hydrologic Normals (1981-2010) Reports. National Water and climate Center: Portland, OR. August 2015

ACIS-USDA Field Office Climate Data (WETS), period of record 1971-2000 <http://agacis.rcc-acis.org> (powered by WRCC) Accessed March 2016

Andrews, R. and R. Righter. 1992. **Colorado Birds**. Denver Museum of Natural History, Denver, CO. 442

Armstrong, D.M. 1972. **Distribution of mammals in Colorado**. Univ. Kansas Museum Natural History Monograph #3. 415.

Butler, LD., J.B. Cropper, R.H. Johnson, A.J. Norman, G.L. Peacock, P.L. Shaver, and K.E. Spaeth. 1997, revised 2003. **National Range and Pasture Handbook**. National Cartography and Geospatial Center's Technical Publishing Team: Fort Worth, TX. <http://www.glti.nrcs.usda.gov/technical/publications/nrph.html> Accessed August 2015

Clark, J., E. Grimm, J. Donovan, S. Fritz, D. Engrstom, and J. Almendinger. 2002. Drought cycles and landscape responses to past Aridity on prairies of the Northern Great Plains, USA. *Ecology*, 83(3), 595-601.

Cleland, D., P. Avers, W.H. McNab, M. Jensen, R. Bailey, T. King, and W. Russell. 1997. National Hierarchical Framework of Ecological Units, published in *Ecosystem Management: Applications for Sustainable Forest and Wildlife Resources*, Yale University Press

Cooperative climatological data summaries. NOAA. Western Regional Climate Center: Reno, NV. Web. <http://www.wrcc.dri.edu/climatedata/climsum> Accessed August 2015

Egan, Timothy. 2006. *The Worst Hard Time*. Houghton Mifflin Harcourt Publishing Company: New York, NY.

Fitzgerald, J.P., C.A. Meaney, and D.M. Armstrong. 1994. *Mammals of Colorado*. Denver Museum of Natural History, Denver, CO. 467.
Hammerson, G.A. 1986. *Amphibians and reptiles in Colorado*. CO Div. Wild. Publication Code DOW-M-I-3-86. 131.

Herrick, Jeffrey E., J.W. Van Zee, K.M. Haystad, L.M. Burkett, and W.G. Witford. 2005. *Monitoring Manual for Grassland, Shrubland, and Savanna Ecosystems, Volume II*. U.S. Dept. of Agriculture, Agricultural Research Service. Jornada Experimental Range, Las Cruces, N.M.

Kingery, H., Ed. (1998) *Colorado Breeding Birds Atlas*. Dist. CO Wildlife Heritage Foundation: Denver, CO. 636.

National Water & Climate Center. USDA-NRCS. USDA Pacific Northwest Climate Hub: Portland, OR. <http://www.wcc.nrcs.usda.gov/> Accessed March 2016

National Weather Service Co-op Program. 2010. Colorado Climate Center. Colorado State Univ. Web. <http://climate.atmos.colostate.edu/dataaccess.php> March 2016

Pellant, M., P. Shaver, D.A. Pyke, J.E. Herrick. (2005) *Interpreting Indicators of Rangeland Health, Version 4*. BLM National Business Center Printed Materials Distribution Service: Denver, CO.

PLANTS Database. 2015. USDA-NRCS. Web. <http://plants.usda.gov/java/> Accessed August 2015. February 2016

PRISM Climate Data. 2015. Prism Climate Group. Oregon State Univ. Corvallis, OR. <http://www.prism.oregonstate.edu/> Accessed August 2015.

Rennicke, J. 1990. *Colorado Wildlife*. Falcon Press, Helena and Billings, MT and CO Div. Wildlife, Denver CO. 138.

Schoeneberger, P.J., D.A. Wysockie, E.C. Benham, and Soil Survey Staff. 2012. *Field book for describing and sampling soils, Version 3.0*. Natural Resources Conservation Service, National Soil Survey Center: Lincoln, NE.

The Denver Posse of Westerners. 1999. *The Cherokee Trail: Bent's Old Fort to Fort Bridger*. The Denver Posse of Westerners, Inc. Johnson Printing: Boulder, CO

U.S. Dept. of Agriculture, Agricultural Research Service. September 1991. *Changes in Vegetation and Land Use I eastern Colorado, A Photographic study, 1904-1986*.

U.S. Dept. of Agriculture, Natural Resources Conservation Service. 2006. *Land Resource Regions and Major Land Resource areas of the United States, the Caribbean, and the Pacific Basin*. US Department of Agriculture Handbook 296.

U.S. Dept. of Agriculture, Natural Resources Conservation Service. National Geospatial Center of Excellence. *Colorado annual Precipitation Map from 1981-2010, Annual Average Precipitation by State*

U.S. Dept. of Agriculture, Natural Resources Conservation Service. 2009. Part 630, Hydrology, National Engineering Handbook

U.S. Dept. of Agriculture, Natural Resources Conservation Service. 1972-2012. National Engineering Handbook Hydrology Chapters. <http://www.nrcs.usda.gov/wps/portal/nrcs/detailfull/national/water/?&cid=stelprdb1043063> Accessed August 2015.

U.S. Dept. of Agriculture, Natural Resources Conservation Service. National Soil Survey Handbook title 430-VI. http://www.nrcs.usda.gov/wps/portal/nrcs/detail/soils/ref/?cid=nrcs142p2_054242 Accessed July 2015

U.S. Dept. of Agriculture, Soil Survey Division Staff. 1993. *Soil Survey Manual*.

U.S. Dept. of Agriculture. 1973. *Soil Survey of Baca County, Colorado*.

U.S. Dept. of Agriculture. 1970. *Soil Survey of Bent County, Colorado*.

U.S. Dept. of Agriculture. 1968. *Soil Survey of Crowley County, Colorado*.

U.S. Dept. of Agriculture. 1981 *Soil Survey of El Paso County Area, Colorado*.

U.S. Dept. of Agriculture. 1995. Soil Survey of Fremont County Area, Colorado.

U.S. Dept. of Agriculture. 1983. Soil Survey of Huerfano County Area, Colorado.

U.S. Dept. of Agriculture. 1981. Soil Survey of Kiowa County, Colorado.

Western Regional Climate Center. 2022. Climate of Colorado, climate of the eastern plains. https://wrcc.dri.edu/Climate/narrative_co.php (accessed 9 August 2022).

Additional Literature:

Clark, J., E. Grimm, J. Donovan, S. Fritz, D. Engrstrom, and J. Almendinger. 2002. Drought cycles and landscape responses to past Aridity on prairies of the Northern Great Plains, USA. *Ecology*, 83(3), 595-601.

Collins, S. and S. Barber. (1985). Effects of disturbance on diversity in mixed-grass prairie. *Vegetation*, 64, 87-94.

Egan, Timothy. 2006. *The Worst Hard Time*. Houghton Mifflin Harcourt Publishing Company: New York, NY.

Hart, R. and J. Hart. 1997. Rangelands of the Great Plains before European Settlement. *Rangelands*, 19(1), 4-11.

Hart, R. 2001. Plant biodiversity on shortgrass steppe after 55 years of zero, light, moderate, or heavy cattle grazing. *Plant Ecology*, 155, 111-118.

Heitschmidt, Rodney K., J.W. Stuth, (edited by). 1991. *Grazing Management, an Ecological Perspective*. Timberland Press, Portland, OR.

Jackson, D. 1966. *The Journals of Zebulon Montgomery Pike with letters & related documents*. Univ. of Oklahoma Press, First edition: Norman, OK.

Mack, Richard N., and J.N. Thompson. 1982. Evolution in Steppe with Few Large, Hooved Mammals. *The American Naturalist*. 119, No. 6, 757-773.

Reyes-Fox, M., Stelzer H., Trlica M.J., McMaster, G.S., Andales, A.A., LeCain, D.R., and Morgan J.A. 2014. Elevated CO₂ further lengthens growing season under warming conditions. *Nature*, April 23 2014. Available online. <http://www.nature.com/nature/journal/v510/n7504/full/nature13207.html>, accessed March 2017.

Stahl, David W., E.R. Cook, M.K. Cleaveland, M.D. Therrell, D.M. Meko, H.D. Grissino-Mayer, E. Watson, and B.H. Luckman. Tree-ring data document 16th century megadrought over North America. 2000. *Eos*, 81(12), 121-125.

The Denver Posse of Westerners. 1999. *The Cherokee Trail: Bent's Old Fort to Fort Bridger*. The Denver Posse of Westerners, Inc. Johnson Printing: Boulder, CO.

U.S. Dept. of Agriculture. 2004. *Vascular plant species of the Comanche National Grasslands in southeastern Colorado*. US Forest Service. Rocky Mountain Research Station. Fort Collins, CO.

Zelikova, Tamara Jane, D.M. Blumenthal, D.G. Williams, L. Souza, D.R. LeCain, J.Morgan. 2014. Long-term Exposure to Elevated CO₂ Enhances Plant Community Stability by Suppressing Dominant Plant Species in a Mixed-Grass Prairie. *Ecology*, 2014 issue. Available online. www.pnas.org/cgi/doi/10.1073/pnas.1414659111.

Contributors

Kimberly Diller, Ecological Site Specialist, NRCS MLRA, Pueblo SSO
Andy Steinert, MLRA 67B Soil Survey Leader, NRCS MLRA Fort Morgan SSO
Ben Berlinger, Rangeland Management Specialist, Retired NRCS La Junta, CO
Doug Whisenhunt, Ecological Site Specialist, NRCS MLRA, Pueblo SSO

Approval

Kirt Walstad, 12/05/2024

Acknowledgments

Program Support: Rachel Murph, NRCS State Rangeland Management Specialist-QC, Denver, CO David Kraft, NRCS MLRA Ecological Site Specialist-QA, Emporia, KS Josh Saunders, Rangeland Management Specialist-QC, NRCS Fort Morgan, CO Patty Knupp, Biologist,

Area 3, NRCS Pueblo, CO Noe Marymor, Biologist, Area 2, NRCS Greeley, CO Richard Mullaney, Resource Conservationist, Retired., NRCS, Akron, CO Chad Remley, Regional Director, N. Great Plains Soil Survey, Salina, KS B.J. Shoup, State Soil Scientist, Denver Eugene Backhaus, State Resource Conservationist, Denver Carla Green Adams, Editor, NRCS, Denver, CO Partners/Contributors: Rob Alexander, Agricultural Resources, Boulder Parks & Open Space, Boulder, CO David Augustine, Research Ecologist, Agricultural Research Service, Fort Collins, CO John Fusaro, Rangeland Management Specialist, NRCS, Fort Collins, CO Jeff Goats, Resource Soil Scientist, NRCS, Pueblo, CO Clark Harshbarger, Resource Soil Scientist, NRCS, Greeley, CO Mike Moore, Soil Scientist, NRCS MLRA Fort Morgan SSO Tom Nadgwick, Rangeland Management Specialist, NRCS, Akron CO Dan Nosal, Rangeland Management Specialist, NRCS, Franktown, CO Steve Olson, Botanist, USFS, Pueblo, CO Randy Reichert, Rangeland Specialist, retired, USFS, Nunn, CO Don Schoderbeck, Range Specialist, CSU Extension, Sterling CO Terri Schultz, The Nature Conservancy, Ft. Collins, CO Chris Tecklenburg, Ecological Site Specialist, Hutchison, KS

Rangeland health reference sheet

Interpreting Indicators of Rangeland Health is a qualitative assessment protocol used to determine ecosystem condition based on benchmark characteristics described in the Reference Sheet. A suite of 17 (or more) indicators are typically considered in an assessment. The ecological site(s) representative of an assessment location must be known prior to applying the protocol and must be verified based on soils and climate. Current plant community cannot be used to identify the ecological site.

Author(s)/participant(s)	Harvey Sprock, Ben Berlinger, Daniel Nosal
Contact for lead author	Harvey Sprock, Area Rangeland Management Specialist, Greeley, CO
Date	11/17/2004
Approved by	
Approval date	
Composition (Indicators 10 and 12) based on	Annual Production

Indicators

1. Number and extent of rills: None

2. Presence of water flow patterns: None

3. Number and height of erosional pedestals or terracettes: None

4. Bare ground from Ecological Site Description or other studies (rock, litter, lichen, moss, plant canopy are not bare ground):
None

5. Number of gullies and erosion associated with gullies: None

6. Extent of wind scoured, blowouts and/or depositional areas: None

7. Amount of litter movement (describe size and distance expected to travel): Typically slight, however during major flooding events this site slows water flow and captures litter and sediment.

8. Soil surface (top few mm) resistance to erosion (stability values are averages - most sites will show a range of values):
Stability class rating is anticipated to be 6 at soil surface.

9. Soil surface structure and SOM content (include type of structure and A-horizon color and thickness): SOM ranges from 3 to 5 percent. Soils are very deep, poorly drained with a water table depth from 5 to 36 inches. Color of the A-horizon is dark brown to black at 0 to 6 inches in depth. Surface structure is weak sub-angular blocky with strong surface aggregates.

10. Effect of community phase composition (relative proportion of different functional groups) and spatial distribution on infiltration and runoff: Raindrop impact is reduced by the diverse grass, forb, shrub functional/structural groups and root structure. This slows overland flow and provides increased time for infiltration to occur. Extended drought, wildfire or both may reduce basal density, canopy cover, and litter amounts (primarily from tall, warm-season bunch and rhizomatous grasses), resulting in decreased infiltration and increased runoff on steep slopes following intense rainfall events.

11. Presence and thickness of compaction layer (usually none; describe soil profile features which may be mistaken for compaction on this site): None

12. Functional/Structural Groups (list in order of descending dominance by above-ground annual-production or live foliar cover using symbols: >>, >, = to indicate much greater than, greater than, and equal to):

Dominant: Warm-season tall grasses >>>

Sub-dominant: Cool-season grasslikes >> cool-season mid rhizomatous >> forbs >>

Other: Cool-season mid grasses = shrubs

Additional:

13. Amount of plant mortality and decadence (include which functional groups are expected to show mortality or decadence):
None to slight.
-
14. Average percent litter cover (%) and depth (in):
-
15. Expected annual annual-production (this is TOTAL above-ground annual-production, not just forage annual-production):
3500 lbs./ac. low precipitation years; 4000 lbs./ac. average; 5000 lbs./ac. high years. Extended drought may reduce annual production by
750 – 1000 lbs./ac.
-
16. Potential invasive (including noxious) species (native and non-native). List species which BOTH characterize degraded
states and have the potential to become a dominant or co-dominant species on the ecological site if their future
establishment and growth is not actively controlled by management interventions. Species that become dominant for only
one to several years (e.g., short-term response to drought or wildfire) are not invasive plants. Note that unlike other
indicators, we are describing what is NOT expected in the reference state for the ecological site: Invasive plants should not
occur in reference plant community.
-
17. Perennial plant reproductive capability: The only limitations are weather-related, wildfire, natural disease, and insects that
temporarily reduce reproductive capability.
-