

Ecological site R070BC025NM Shallow

Accessed: 09/17/2026

General information

Provisional. A provisional ecological site description has undergone quality control and quality assurance review. It contains a working state and transition model and enough information to identify the ecological site.

Table 1. Dominant plant species

Tree	Not specified
Shrub	Not specified
Herbaceous	Not specified

Physiographic features

This site occurs on knolls, ridges, hillslopes alluvial fans and escarpments. Slopes range fro 0 to 25 percent and average about 7 percent. Direction of slope varies and is usually not significant. Elevations range from 2,842 to 4,500 feet.

Table 2. Representative physiographic features

Landforms	(1) Hill (2) Ridge (3) Fan piedmont
Flooding frequency	None
Ponding frequency	None
Elevation	870 – 1,370 m
Slope	0 – 30 %
Aspect	Aspect is not a significant factor

Climatic features

The average annual precipitation ranges from 8 to 13 inches. Variations of 5 inches, more or less, are common. Over 80 percent of the precipitation falls from April through October. Most of the summer precipitation comes in the form of high intensity – short duration

thunderstorms.

Temperatures are characterized by distinct seasonal changes and large annual and diurnal temperature changes. The average annual temperature is 61 degrees with extremes of 25 degrees below zero in the winter to 112 degrees in the summer.

The average frost-free season is 180 to 220 days. The last killing frost is late March or early April, and the first killing frost is in late October or early November.

Temperature and rainfall both favor warm season perennial plant growth. In years of abundant spring moisture, annual forbs and cool season grasses can make up an important component of this site. Because of the shallow soil depth, the vegetation on this site can take advantage of moisture almost anytime it falls. Strong winds that blow from the west and southwest blow from January through June, which accelerates soil drying at a critical time for cool season plant growth.

Climate data was obtained from <http://www.wrcc.sage.dri.edu/summary/climsmnm.html> web site using 50% probability for freeze-free and frost-free seasons using 28.5 degrees F and 32.5 degrees F respectively.

Table 3 Representative climatic features

Frost-free period (average)	220 days
Freeze-free period (average)	240 days
Precipitation total (average)	330 mm

Influencing water features

This site is not influenced from water from wetlands or streams.

Soil features

The soils of this site are shallow to very shallow. Soils are derived from mixed calcareous eolian deposits derived from sedimentary rock. Surface layers are very cobbly loam, very gravelly loam, gravelly loam, cobbly loam, gravelly fine sandy loam or gravelly sandy loam.

There is an indurated caliche layer or limestone bedrock that occurs within 20 inches and averages less than 10 inches. Limestone or caliche layer may be the restrictive layer.

Minimum and maximum values listed below represent the characteristic soils for this site.

Characteristic soils:

Lozier
Potter
Tencee
Upton
Ector
Kimbrough

Table 4. Representative soil features

Surface texture	(1) Gravelly loam (2) Extremely gravelly loam (3) Extremely cobbly loam
Family particle size	(1) Loamy

Drainage class	Well drained
Permeability class	Very slow to moderately slow
Soil depth	10 – 50 cm
Surface fragment cover <=3"	20 – 40 %
Available water capacity (0-101.6cm)	2.54 cm
Calcium carbonate equivalent (0-101.6cm)	20 – 60 %
Electrical conductivity (0-101.6cm)	Not specified
Sodium adsorption ratio (0-101.6cm)	Not specified
Soil reaction (1:1 water) (0-101.6cm)	7.4 – 8.4
Subsurface fragment volume <=3" (Depth not specified)	10 – 40 %
Subsurface fragment volume >3" (Depth not specified)	Not specified

Ecological dynamics

Overview:

The Shallow site is associated with and Limestone Hills, Loamy, and Shallow Sandy sites. When associated with Limestone Hills, the Shallow site occurs on the summits, foot slopes and toeslopes of hills. Loamy sites often occur as areas between low elongated hills with rounded crests (Shallow site). When the Shallow Sandy site and Shallow site occur in association, the Shallow Sandy soils occupy the tops of low ridges and the Shallow site soils occur on the steeper sideslopes of the ridge. The historic plant community of the Shallow site has the aspect of a grassland/shrub mix, dominated by grasses, but with shrubs common throughout the site. Black grama is the dominant grass species; creosotebush, mesquite, and catclaw mimosa are common shrubs. Overgrazing and or extended drought can reduce grass cover, effect a change in grass species dominance, and may result in a shrub-dominated state. 1

State and transition model

Additional community tables

Table 5. Community 1.1 plant community composition

Group	Common Name	Symbol	Scientific Name	Annual Production ()	Foliar Cover (%)
Grass/Grasslike					
1				118-177	
	black grama	BOER4	<i>Bouteloua eriopoda</i>	118-177	–
2				89-118	
	sideoats grama	BOCU	<i>Bouteloua curtipendula</i>	89-118	–
3				89-118	
	blue grama	BOGR2	<i>Bouteloua gracilis</i>	89-118	–
	hairy grama	BOHI2	<i>Bouteloua hirsuta</i>	89-118	–
4				29-59	
	bush muhly	MUPO2	<i>Muhlenbergia porteri</i>	29-59	–
5				18-29	
	cane bluestem	BOBA3	<i>Bothriochloa barbinodis</i>	18-29	–
6				29-59	
	sand dropseed	SPCR	<i>Sporobolus cryptandrus</i>	29-59	–
7				18-29	
	hairy woollygrass	ERPI5	<i>Erioneuron pilosum</i>	18-29	–
8				6-18	
	ear muhly	MUAR	<i>Muhlenbergia arenacea</i>	6-18	–
9				6-18	
	New Mexico feathergrass	HENE5	<i>Hesperostipa neomexicana</i>	6-18	–
10				6-18	
	low woollygrass	DAPU7	<i>Dasyochloa pulchella</i>	6-18	–
11				18-29	
	Grass, perennial	2GP	<i>Grass, perennial</i>	18-29	–
Forb					
12				12-29	
	stemless four-nerve daisy	TEACE	<i>Tetranneuris acaulis</i> var. <i>epunctata</i>	12-29	–
13				6-18	
	woolly groundsel	PACA15	<i>Packera cana</i>	6-18	–
14				6-18	
	globemallow	SPHAE	<i>Sphaeralcea</i>	6-18	–
15				6-18	
	bladderpod	LESQU	<i>Lesquerella</i>	6-18	–
16				6-18	
	cassia	CASSI	<i>Cassia</i>	6-18	–
17				12-29	
	Forb (herbaceous, not grass nor grass-like)	2FORB	<i>Forb (herbaceous, not grass nor grass-like)</i>	12-29	–
Shrub/Vine					
18				6-18	
	littleleaf sumac	RHMI3	<i>Rhus microphylla</i>	6-18	–
19				6-18	

	creosote bush	LATR2	<i>Larrea tridentata</i>	6-18	–
20				6-18	
	littleleaf ratany	KRER	<i>Krameria erecta</i>	6-18	–
21				6-18	
	javelina bush	COER5	<i>Condalia ericoides</i>	6-18	–
22				6-18	
	American tarwort	FLCE	<i>Flourensia cernua</i>	6-18	–
23				6-18	
	crown of thorns	KOSP	<i>Koeberlinia spinosa</i>	6-18	–
24				12-29	
	honey mesquite	PRGL2	<i>Prosopis glandulosa</i>	12-29	–
	honey mesquite	PRGL2	<i>Prosopis glandulosa</i>	12-29	–
25				6-18	
	catclaw mimosa	MIACB	<i>Mimosa aculeaticarpa var. biuncifera</i>	6-18	–
26				6-18	
	pricklypear	OPUNT	<i>Opuntia</i>	6-18	–
27				12-29	
	mariola	PAIN2	<i>Parthenium incanum</i>	12-29	–
	mariola	PAIN2	<i>Parthenium incanum</i>	12-29	–
28				6-18	
	broom snakeweed	GUSA2	<i>Gutierrezia sarothrae</i>	6-18	–
29				18-29	
	Shrub (>.5m)	2SHRUB	<i>Shrub (>.5m)</i>	18-29	–

Table 6. Community 2.1 plant community composition

Group	Common Name	Symbol	Scientific Name	Annual Production ()	Foliar Cover (%)
-------	-------------	--------	-----------------	----------------------	------------------

Animal community

This site provides habitats which support a resident animal community that is characterized by desert cottontail, spotted ground squirrel, Merriam's kangaroo rat, cactus mouse, white-throated woodrat, gray fox, spotted skunk, roadrunner, Swainson's hawk, white-necked raven, cactus wren, pyrrhuloxia, lark sparrow, mourning dove, scaled quail, leopard lizard, round-tailed horned lizard, prairie rattlesnake, marbled whiptail, and greater earless lizard. Where associated with limestone hills, mule deer utilize this site. Where large woody shrubs occur, most resident birds and scissor-tailed flycatcher, morning dove, lark sparrow and Swainson's hawk nest.

Hydrological functions

The runoff curve numbers are determined by field investigations using hydraulic cover conditions and hydrologic soil groups. Hydrologic Interpretations Soil Series----- Hydrologic Group Lozier----- D Potter----- C Tencee----- D Upton----- C Kimbrough----- D Upton----- D Ector----- D

Recreational uses

This site offers recreation potential for hiking, horseback riding, rock hunting, nature photography and bird hunting and birding. During years of abundant spring moisture, a colorful array of wild flowers is displayed during May and June. A few summer and fall flowers also occur.

Wood products

This site has no potential for wood production.

Other products

This site is suited for grazing by all kinds and classes of livestock during all seasons of the year. Missmanagement will cause a decrease in black grama, sideoats grama, and blue grama, bush muhly and New Mexico feathergrass. A corresponding increase in bare ground will occur. There will also be an increase in muhlys, fluffgrass, creosotebush, javalinabush, catclaw, and mesquite. This site will respond best to a system of management that rotates the season of use.

Other information

Guide to Suggested Initial Stocking Rate Acres per Animal Unit Month Similarity Index----- Ac/AUM 100 - 76----- 3.7 – 4.5 75 – 51----- 4.3 – 5.5 50 – 26----- 5.3 – 10.0 25 – 0----- 10.1 +

Inventory data references

Data collection for this site was done in conjunction with the progressive soil surveys within the Southern Desertic Basins, Plains and Mountains, Major Land Resource Areas of New Mexico (SD-3). This site has been mapped and correlated with soils in the following soil surveys. Eddy County, Lea County, and Chaves County.

Other references

Literature Cited:

1. Humphrey, R.R. 1974. Fire in the deserts and desert grassland of North America. In: Kozlowski, T. T.; Ahlgren, C. E., eds. Fire and ecosystems. New York: Academic Press: 365-400.
2. Hennessy, J.T., R.P. Gibbens, J.M. Tromble, and M. Cardenas. 1983. Water properties of caliche. J. Range Manage. 36: 723-726.
3. U.S. Department of Agriculture, Natural Resources Conservation Service. 2001. Soil Quality Information Sheets. Rangeland Soil Quality—Infiltration, Organic Matter, Rangeland Sheets 5,6. [Online]. Available: <http://www.statlab.iastate.edu/survey/SQI/range.html>

Contributors

David Trujillo
Don Sylvester

Rangeland health reference sheet

Interpreting Indicators of Rangeland Health is a qualitative assessment protocol used to determine ecosystem condition based on benchmark characteristics described in the Reference Sheet. A suite of 17 (or more) indicators are typically considered in an assessment. The ecological site(s) representative of an assessment location must be known prior to applying the protocol and must be verified based on soils and climate. Current plant community cannot be used to identify the ecological site.

Author(s)/participant(s)	
Contact for lead author	
Date	
Approved by	
Approval date	
Composition (Indicators 10 and 12) based on	Annual Production

Indicators

1. Number and extent of rills:

2. Presence of water flow patterns:

3. Number and height of erosional pedestals or terracettes:

4. Bare ground from Ecological Site Description or other studies (rock, litter, lichen, moss, plant canopy are not bare ground):

5. Number of gullies and erosion associated with gullies:

6. Extent of wind scoured, blowouts and/or depositional areas:

7. Amount of litter movement (describe size and distance expected to travel):

8. Soil surface (top few mm) resistance to erosion (stability values are averages - most sites will show a range of values):

9. Soil surface structure and SOM content (include type of structure and A-horizon color and thickness):

10. Effect of community phase composition (relative proportion of different functional groups) and spatial distribution on infiltration and runoff:

11. Presence and thickness of compaction layer (usually none; describe soil profile features which may be mistaken for compaction on this site):

12. Functional/Structural Groups (list in order of descending dominance by above-ground annual-production or live foliar cover using symbols: >>, >, = to indicate much greater than, greater than, and equal to):

Dominant:

Sub-dominant:

Other:

Additional:

13. Amount of plant mortality and decadence (include which functional groups are expected to show mortality or decadence):

14. Average percent litter cover (%) and depth (in):

15. Expected annual annual-production (this is TOTAL above-ground annual-production, not just forage annual-production):

16. Potential invasive (including noxious) species (native and non-native). List species which BOTH characterize degraded states and have the potential to become a dominant or co-dominant species on the ecological site if their future establishment and growth is not actively controlled by management interventions. Species that become dominant for only one to several years (e.g., short-term response to drought or wildfire) are not invasive plants. Note that unlike other indicators, we are describing what is NOT expected in the reference state for the ecological site:

17. Perennial plant reproductive capability:
