

Ecological site R072XY100KS

Loamy Tableland

Last updated: 6/06/2025

Accessed: 09/01/2026

General information

Approved. An approved ecological site description has undergone quality control and quality assurance review. It contains a working state and transition model, enough information to identify the ecological site, and full documentation for all ecosystem states contained in the state and transition model.

MLRA notes

Major Land Resource Area (MLRA): 072X--Central High Tableland

Major Land Resource Area (MLRA) 72--Central High Tableland. This area is in Kansas (54 percent), Nebraska (25 percent), and Colorado (21 percent). A very small part of the area is in Wyoming. The area makes up about 34,550 square miles (89,535 square kilometers). It includes the towns of Garden City, Goodland, and Colby, Kansas; Imperial, North Platte, Ogallala, and Sidney, Nebraska; and Holyoke and Wray, Colorado. Interstate 70 bisects the area, and Interstates 76 and 80 follow the south side of the South and North Platte Rivers, respectively. The Cimarron National Grasslands occur in the southeast corner of the MLRA.

Classification relationships

MLRA 72 is in the Central Great Plains Winter Wheat and Range Land Resource Region. It is in the High Plains Section, of the Great Plains Province, of the Interior Plains (USDA, 2006). MLRAs can be defined by climate, landscapes, geology, and annual precipitation zones (PZ). Other features such as landforms, soil properties, and key vegetation further refine these concepts, and are described at the Ecological Site Description (ESD) level. Relationship to Other Classifications: Natural Resource Conservation Service (NRCS) Classification Hierarchy: Physiographic Division Physiographic Province Physiographic Section Land Resource Region Major Land Resource Area Land Resource Unit (Fenneman, 1946) United States Forest Service (USFS) Classification Hierarchy: Domain Division Province Section Subsection Landtype Association Landtype Landtype Phase (Cleland et al, 1997) Revision Notes: The Loamy Tableland site was developed by an earlier version of the Loamy Upland (2001). The earlier version of the Loamy Upland ESD was based upon input from NRCS (formerly Soil Conservation Service) and historical information obtained from the Loamy Upland Range Site Description (1967, updated1983).

Ecological site concept

The Loamy Tableland ecological site is located on plains, rises and hillslopes on tablelands. Soils correlated with this site are moderately deep to very deep and have a surface that is >7 (18cm)inches mollic color. The majority of the site silt loam, loam, and very fine sandy loam. Soils that are correlated to Loamy Tableland have free carbonates below 4 inches (10cm). This site is dominated by loess parent material.

Associated sites

R072XY101KS	Limy Slopes The Limy Slopes ecological site is located on shoulders and backslopes on hillslopes on tableland landscapes. Soils correlated with this site are moderately deep to very deep and have a surface that is <8 inches (20cm). The soil surface texture ranges from silt loam to loam, with most of the site surface textures being silt loam. Soils that are correlated to Limy Slopes have free carbonates occurring within 4 inches (10cm) of the surface. This site is dominated by loess parent material.
-------------	---

R072XY115KS	Closed Upland Depression The Closed Upland Depression ecological site occurs on broad upland divides (Tablelands). This site occurs in small to large depressions called playas that have very slow to slow permeability classes and a range of low to high available water capacity. This site receives water runoff from areas higher on the landscape.
R072XY108KS	Loamy Lowland Alluvial soils make up the parent material of this ecological site. This site occurs on flood plains or low stream terraces and is characterized by a seasonal or perennial high water table greater than 6 feet from the surface. The soils characteristic of this site are deep and well drained that formed in calcareous alluvial sediments. The Loamy Tableland ecological site occurs on broad upland divides and can be found adjacent to this site.

Similar sites

R072XY115KS	Closed Upland Depression The Closed Upland Depression ecological site occurs on broad upland divides (Tablelands). This site occurs in small to large depressions called playas that have very slow to slow permeability classes and a range of low to high available water capacity. This site receives water runoff from areas higher on the landscape.
R072XY117KS	Tableland Swale The Tableland Swale ecological site is located in swales and draws on tablelands. These sites are most often found adjacent to the Loamy Tableland sites and occur in water- gathering positions on the landscape.
R072XY108KS	Loamy Lowland Alluvial soils make up the parent material of this ecological site. This site occurs on flood plains or low stream terraces and is characterized by a seasonal or perennial high water table greater than 6 feet from the surface. The soils characteristic of this site are deep and well drained that formed in calcareous alluvial sediments.
R072XY116KS	Salt Flat The Salt Flat ecological site occurs on base and side slopes of closed depressions and alluvial fans. This site has saline and sodium affected soils.

Table 1. Dominant plant species

Tree	Not specified
Shrub	Not specified
Herbaceous	(1) <i>Bouteloua gracilis</i> (2) <i>Pascopyrum smithii</i>

Physiographic features

This site occurs on nearly level to sloping hillslopes, rises and plains on tablelands. This site is dominated by Quaternary Period loess deposits and eolian deposits of Tertiary Period material from the local rivers and streams. This site consists of very deep upland soils with silty or loamy surface layers and subsoil. This site produces runoff to areas lower on the landscape. This site is subject to erosion by wind and water if the vegetative cover is reduced or absent by such things as overgrazing and fire events. Elevation for this site ranges from

2,200 to 5,400 feet.

The extent of the major Hydrologic Unit Areas (identified by four-digit numbers) that make up MLRA 72 is as follows: Republican (1025), 38 percent; Middle Arkansas (1103), 20 percent; Smoky Hill (1026), 15 percent; South Platte (1019), 13 percent; Upper Cimarron (1104), 11 percent; North Platte (1018), 2 percent; and Upper Arkansas (1102), 1 percent. The North Platte River forms the northern boundary of this MLRA. The South Platte River joins the North Platte River at the town of North Platte, Nebraska. The Arkansas River bisects the southern part of the MLRA. Other large rivers between the North Platte and Arkansas Rivers in the area include the Republican, Sappa, Prairie Dog, Solomon, Saline, and Smoky Hill Rivers. The Cimarron River is the southern boundary of MLRA 72 in the western part of this area.

Figure 1. MLRA72 ESD block diagram

Table 2. Representative physiographic features

Landforms	(1) Plain (2) Hill (3) Rise
Flooding frequency	None
Ponding frequency	None
Elevation	670 – 1,650 m
Slope	0 – 20 %
Water table depth	200 cm

Climatic features

The average annual precipitation in this area is 14 to 25 inches (355 to 635 millimeters). It fluctuates widely from year to year. Most of the rainfall occurs as high-intensity, convective thunderstorms during the growing season. The maximum precipitation occurs from late spring through early autumn. Precipitation in winter occurs as snow. The annual snowfall ranges from about 16 inches (40 centimeters) in the southern part of the area to 35 inches (90 centimeters) in the northern part. The average annual temperature is 46 to 57 degrees F (8 to 14 degrees C). The freeze-free period averages 159 days and ranges from 135 to 210 days, increasing in length from northwest to southeast. Climate data comes from the Natural Resources Conservation Service (NRCS) National Water and Climate Center. The data set is from 1981-2010.

Table 3 Representative climatic features

Frost-free period (characteristic range)	120-130 days
Freeze-free period (characteristic range)	140-160 days
Precipitation total (characteristic range)	460-530 mm
Frost-free period (actual range)	110-140 days

Freeze-free period (actual range)	140-170 days
Precipitation total (actual range)	430-530 mm
Frost-free period (average)	130 days
Freeze-free period (average)	150 days
Precipitation total (average)	480 mm

- (1) TRENTON DAM [USC00258628], Trenton, NE
- (2) JULESBURG [USC00054413], Julesburg, CO
- (3) YUMA [USC00059295], Yuma, CO
- (4) ULYSSES 3NE [USC00148287], Ulysses, KS
- (5) KIMBALL 2NE [USC00254440], Kimball, NE
- (6) BREWSTER 4W [USC00141029], Brewster, KS
- (7) TRIBUNE 1W [USC00148235], Tribune, KS
- (8) SCOTT CITY [USC00147271], Scott City, KS
- (9) WALLACE 2W [USC00258920], Wallace, NE

Influencing water features

There are no water features of the ecological site or adjacent wetland/riparian regimes that influence the vegetation and/or management of the site that make it distinctive from other ecological sites. Many of the mapunits in this ESD have a 1-3% hydric component for closed upland depressions (playas).




Figure 8. Fig. 7-1 is from the National Range and Pasture Ha

Soil features

The soils on this site are well drained and are moderately deep to very deep on slopes ranging from nearly level to gently sloping. Most of the soils on this site have loess as parent material and are dominated by silt loam and silty clay loam textures. The geology is made up by river valley deposits of material from the Tertiary Period that have been wind deposited on the adjacent tableland or by Peoria Loess, which is glacial debris from the glacial retreat to the north during the Quaternary period.

The surface layer of the soils in this site is primarily moderately-fine textured to medium textured (but the range includes fine-textured). The surface layer ranges from a depth of 4 to 11 inches thick and permeability is moderately slow to moderate. The subsoil and underlying material have a similar range in texture as the surface layer. Contrasting sandy (gravelly) or very clayey layers may occur at depths below 40 inches in several of the listed soil series. Soils in this site are generally high in fertility and can have free carbonates below 6 inches. These soils are susceptible to erosion by water and wind. The potential for water erosion accelerates with increasing slope.

Surface soil structure is granular to subangular blocky, and structure below the surface is prismatic or subangular blocky. Soil structure describes the manner in which soil particles are aggregated and defines the nature of the system of pores and channels in a soil. Together, soil texture and structure help determine the ability of the soil to hold and conduct the water and air necessary for sustaining life.

Available water holding capacity (AWC) ranges from 1.9 to 2.7 inches of water per foot of soil depth. Available water is the portion of water in a soil that can be readily absorbed by plant roots. This is the amount of water released between the field capacity and the permanent wilting point. As fineness of texture increases, there is a general increase in available moisture storage from sands to loams and silt loams. Soil organic matter (SOM) averages 1 to 2 percent.

Major soil series correlated to this ecological site include: Ulysses, Richfield, Keith, Kuma, Rosebud, Blackwood, Mace, McConaughy, Norka, and Satanta.

Figure 9. Ulysses, 2 to 7 percent slope

Table 4. Representative soil features

Surface texture	(1) Silt loam (2) Loam (3) Very fine sandy loam
Family particle size	(1) Loamy
Drainage class	Well drained
Permeability class	Moderately slow to moderate
Depth to restrictive layer	Not specified
Soil depth	50 – 200 cm
Surface fragment cover <=3"	0 – 20 %
Surface fragment cover >3"	Not specified

Available water capacity (0-101.6cm)	4.83 – 6.86 cm
Calcium carbonate equivalent (0-101.6cm)	0 – 10 %
Electrical conductivity (0-101.6cm)	Not specified
Sodium adsorption ratio (0-101.6cm)	Not specified
Soil reaction (1:1 water) (0-101.6cm)	6.6 – 8.4
Subsurface fragment volume <=3" (Depth not specified)	10 – 40 %
Subsurface fragment volume >3" (Depth not specified)	Not specified

Ecological dynamics

The plant communities for this site are dynamic due to the complex interaction of many ecological processes. The interpretive plant community for this site is the Reference Plant Community. The reference community has been determined by the study of rangeland relic areas, areas protected from excessive disturbance, areas under long term rotational grazing strategies, literature of plant communities from the early 1900's and local expertise. Trends in plant community dynamics ranging from heavily grazed to lightly grazed areas, seasonal use pastures, and historical accounts have also been used.

This ecological site is made up of a grassland state, a sod-bound state and a tillage state. The grassland state is characterized by non-broken land (no tillage), both warm and cool- season bunchgrasses, sod-forming grasses, forbs and shrubs. The sod-bound state is characterized by a warm-season, shortgrass plant community made up of primarily buffalograss and blue grama with few remnant western wheatgrass motts and very few forbs. The tillage state has been mechanically disturbed (broken) by equipment and includes either a variety of reseeded warm-season bunch and sod forming grasses or early successional plants to include the latter as well as annual grasses and forbs.

Vegetation changes are expected within this ecological site and will be dependent on the site's geographical location inside Major Land Resource Area 72 (MLRA). The northern part of MLRA 72 is characterized by cooler temperatures and shorter growing season in respect to the southern end. As a result, cool-season bunchgrasses and sod formers proliferate. Growth of native cool-season plants begins about April 15, and continues to about June 15. Native warm-season plants begin growth about May 15, and continue to about August 15. Green up of cool-season plants may occur in September and October if adequate moisture is available (weather data from National Climate Data Center 1980-2010).

This site developed with occasional fires being part of the ecological processes. Historically, it is believed that the fires were infrequent, randomly distributed, and started by lightning at various times throughout the season when thunderstorms were likely to occur. It is also believed that pre-European inhabitants may have used fire as a management tool for attracting herds of large migratory herbivores (bison, elk, deer and pronghorn). The impact of fire over the past 100 years has been relatively insignificant due to the human control of

wildfires and the lack of acceptance of prescribed fire as a management tool in the semiarid, High Plains area.

The degree of herbivory (feeding on herbaceous plants) has a significant impact on the dynamics of the site. Historically, periodic grazing by herds of large migratory herbivores was a primary influence. Secondary influences of herbivory by species such as prairie dogs, grasshoppers, gophers and root feeding organisms impacted the vegetation historically, and continue to this day.

The management of herbivory by humans through grazing of domestic livestock and/or manipulation of wildlife populations has been a major influence on the ecological dynamics of the site. This management coupled with the High Plains climate largely dictates the plant communities for the site.

Drought cycles were part of the natural range of variability within the site and have historically had a major impact upon the vegetation. The species composition changes according to the duration and severity of the drought cycle (Albertson, F. W., Weaver, J. E.).

This site occurs on exposed interfluvies. Due to the landform position, these level areas are preferred by livestock, which can lead to grazing distribution problems.

Historically, mechanical treatment of this site was practiced. The theory of mechanical treatment was that it improves production and plant composition on rangeland. These mechanical treatments include such things as contour furrowing, contour pitting, terracing, chiseling, disking and inter-seeding. Many of these treatments were implemented during the 1930's through the 1960's and have shown to have no significant long-term benefits for improving production. Many of these practices result in a permanently rough ground surface. Inter-seeding may be beneficial depending upon stand achieved and management used after seeding.

State and transition model

Additional community tables

Table 5. Community 1.1 plant community composition

Group	Common Name	Symbol	Scientific Name	Annual Production ()	Foliar Cover (%)
Grass/Grasslike					
1	Shortgrass warm-season; Dominant 40%			336-762	
	blue grama	BOGR2	<i>Bouteloua gracilis</i>	252-493	–
	buffalograss	BODA2	<i>Bouteloua dactyloides</i>	84-269	–
2	Tall/Midgrass warm-season; Subdominant 21%			67-404	
	sideoats grama	BOCU	<i>Bouteloua curtipendula</i>	22-168	–
	little bluestem	SCSC	<i>Schizachyrium scoparium</i>	22-112	–
	composite dropseed	SPCO16	<i>Sporobolus compositus</i>	0-50	–
	sand dropseed	SPCR	<i>Sporobolus cryptandrus</i>	0-50	–
	big bluestem	ANGE	<i>Andropogon gerardii</i>	0-50	–
	Fendler threeawn	ARPUL	<i>Aristida purpurea</i> var. <i>longiseta</i>	0-17	–
3	Midgrass cool-season; Subdominant 18%			224-336	
	western wheatgrass	PASM	<i>Pascopyrum smithii</i>	224-303	–
	green needlegrass	NAVI4	<i>Nassella viridula</i>	0-135	–
	needle and thread	HECO26	<i>Hesperostipa comata</i>	0-90	–
	squirreltail	ELEL5	<i>Elymus elymoides</i>	0-17	–
4	Sedges; Minor 6%			0-118	
	threadleaf sedge	CAFI	<i>Carex filifolia</i>	0-56	–
	sun sedge	CAINH2	<i>Carex inops</i> ssp. <i>heliophila</i>	0-34	–
	sedge	CAREX	<i>Carex</i>	0-28	–
Forb					
5	Forbs; Minor 10%			17-191	
	scarlet globemallow	SPCO	<i>Sphaeralcea coccinea</i>	6-28	–
	white heath aster	SYERE	<i>Symphotrichum ericoides</i> var. <i>ericoides</i>	0-22	–
	upright prairie coneflower	RACO3	<i>Ratibida columnifera</i>	6-22	–
	dotted blazing star	LIPU	<i>Liatris punctata</i>	0-22	–

	Cuman ragweed	AMPS	<i>Ambrosia psilostachya</i>	6-22	–
	white sagebrush	ARLU	<i>Artemisia ludoviciana</i>	6-22	–
	purple prairie clover	DAPU5	<i>Dalea purpurea</i>	0-17	–
	scarlet beeblossom	GACO5	<i>Gaura coccinea</i>	0-17	–
	lacy tansyaster	MAPI4	<i>Machaeranthera pinnatifida ssp. pinnatifida var. pinnatifida</i>	0-17	–
	broadbeard beardtongue	PEAN4	<i>Penstemon angustifolius</i>	0-17	–
	slimflower scurfpea	PSTE5	<i>Psoralidium tenuiflorum</i>	6-17	–
	American vetch	VIAM	<i>Vicia americana</i>	0-17	–
	common sunflower	HEAN3	<i>Helianthus annuus</i>	0-11	–
	rush skeletonplant	LYJU	<i>Lygodesmia juncea</i>	0-11	–
	woolly locoweed	ASMO7	<i>Astragalus mollissimus</i>	0-11	–
	yellowspine thistle	CIOC2	<i>Cirsium ochrocentrum</i>	0-6	–
	Texas croton	CRTE4	<i>Croton texensis</i>	0-6	–
Shrub/Vine					
6	Shrubs; Minor 5%			17-95	
	fourwing saltbush	ATCA2	<i>Atriplex canescens</i>	0-45	–
	prairie sagewort	ARFR4	<i>Artemisia frigida</i>	0-34	–
	broom snakeweed	GUSA2	<i>Gutierrezia sarothrae</i>	0-22	–
	winterfat	KRLA2	<i>Krascheninnikovia lanata</i>	0-22	–
	plains pricklypear	OPPO	<i>Opuntia polyacantha</i>	6-22	–
	soapweed yucca	YUGL	<i>Yucca glauca</i>	6-22	–
	leadplant	AMCA6	<i>Amorpha canescens</i>	6-17	–

Table 6. Community 1.2 plant community composition

Group	Common Name	Symbol	Scientific Name	Annual Production ()	Foliar Cover (%)
-------	-------------	--------	-----------------	----------------------	------------------

Table 7. Community 2.1 plant community composition

Group	Common Name	Symbol	Scientific Name	Annual Production ()	Foliar Cover (%)
-------	-------------	--------	-----------------	----------------------	------------------

Table 8. Community 3.1 plant community composition

Group	Common Name	Symbol	Scientific Name	Annual Production ()	Foliar Cover (%)
-------	-------------	--------	-----------------	----------------------	------------------

Table 9. Community 3.2 plant community composition

Group	Common Name	Symbol	Scientific Name	Annual Production ()	Foliar Cover (%)
-------	-------------	--------	-----------------	----------------------	------------------

Animal community

Grazing by domestic livestock is one of the major income-producing industries in the area. Rangelands in this area provide year-long forage under prescribed grazing for cattle, sheep, horses and other herbivores. During the dormant period, livestock may need supplementation based on reliable forage analysis. Grazing Interpretations Calculating Safe Stocking Rates: Proper stocking rates should be incorporated into a grazing management strategy that protects the resource, maintains or improves rangeland health, and is consistent with management objectives. In addition to usable forage, safe stocking rates should consider ecological condition, trend of the site, past grazing use history, season of use, stock density, kind and class of livestock, forage digestibility, forage nutritional value, variation of harvest efficiency based upon desirability preference of plant species and/or grazing system and site grazability factors (such as steep slopes, site inaccessibility, or distance to drinking water). Often the current plant community does not entirely match any given community phase as described in this Ecological Site Description. Because of this, a resource inventory is necessary to document plant composition and production. Proper interpretation of inventory data will permit the establishment of a safe initial stocking rate. No two years have the same weather conditions. For this reason, year-to-year and season-to season fluctuations in forage production are to be expected on grazing lands. Livestock producers must make timely adjustments in the numbers of animals or in the length of grazing periods to avoid overuse of forage plants when production is unfavorable and to make advantageous adjustments when forage supplies are above average. Initial stocking rates should be improved though the use of vegetation monitoring and actual use records that include number and type of livestock, the timing and duration of grazing, and utilization levels. Actual use records over time will assist in making

stocking rate adjustments based on the variability factors. Average annual production must be measured or estimated to properly assess useable forage production and stocking rates.

Hydrological functions

Water is the principal factor limiting forage production on this site. This site is dominated by soils in hydrologic group A and B. Infiltration and runoff potential for this site varies from moderate to high depending on soil hydrologic group and ground cover. In many cases, areas with greater than 75 percent ground cover have the greatest potential for high infiltration and lower runoff. An example of an exception would be where shortgrasses form a strong sod and dominate the site. Areas where ground cover is less than 50 percent have the greatest potential to have reduced infiltration and higher runoff. (Refer to NRCS Section 4, National Engineering Handbook (NEH-4) for runoff quantities and hydrologic curves).

Recreational uses

Because of the deep fertile soils and gentle slopes of this site, it is continually in danger of development for cropland, home sites, roads, and urban uses. The site exhibits little visual contrast but does present a panoramic view of the wide-open spaces cherished by many in the Great Plains states. Hunting opportunities for upland game species abound.

Wood products

No appreciable wood products are present on the site.

Other products

None noted

Other information

Site development and testing plan. This ESD has gone through the approval process.

Inventory data references

Information presented here has been derived from NRCS clipping data and other inventory data. Field observations from range trained personnel was also used. Those involved in developing the initial ESD in 2001 include: Tim Watson, Amanda Shaw, Susan Francis, Jon Deege, Harvey Sprock, Robert Schiffner, Josh Saunders, Paul Billig, Ted Houser, Rick Cantu, JoAnn Geinger, Chuck Markley, Jeff Nichols, Mary Schrader, and Carol Eakins. Range Condition Guides and Technical Range Site Descriptions for Kansas, Loamy Upland, USDA, Soil Conservation Service, August, 1967 Range Site Description for Kansas, Loamy Upland, USDA-Soil Conservation Service, September, 1983 Range Site Description for Colorado, Loamy Plains, USDA-Soil Conservation Service, December 1975 Range Site Description for Colorado, Loamy Plains, USDA-Soil Conservation Service, January, 1982 Guide for determining range condition and suggestive initial stocking rates for Nebraska, Silty, Vegetative Zone 1, USDA-Soil Conservation Service, April 1983 Range Site Description for Nebraska, Silty, USDA-Soil Conservation Service, August, 1981 Schacht, Walter H., Larsen, Dana. Section III Range Sites, Silty Range Site, The Board of Regents of the University of Nebraska, publication Ecological Site Description for Kansas, Loamy Upland North (R072XA015KS) and South (R072XB015KS), located in Ecological Site Information System (ESIS), 2007 Ecological Site Description for Colorado, Loamy Plains (R072XY001CO), located in Ecological Site Information System (ESIS)

Other references

USDA Handbook 296, LRR and MLRA of the U.S., the Caribbean, and the Pacific Basin, 2006.

National Range and Pasture Handbook, USDA-NRCS, Chapter 7, Rangeland and Pastureland Hydrology and Erosion.

Soil Series—Official Series Descriptions, <https://soilseries.sc.egov.usda.gov/osdname.asp>

USDA-Natural Resources Conservation Service—Soil Surveys and Web Soil Survey

National Climatic Data Center, Weather data, web site <http://www.ncdc.noaa.gov/>

USDA-ARS-85 Changes in Vegetation and Land Use in Eastern Colorado, September 1991, A photographic study, 1904 to 1986

Range Management: principles and practices/ Jerry L., Holechek, Rex D. Pieper, Carlton H. Herbel.—5th ed.

Life History and Habits of Blue Grama, Andrew Riegel, Transactions of the Kansas Academy of Science (1903-) Vol. 44 (April 3-5, 1941). pp. 76-85

A New Vegetation Map of Kansas. A. W. Kuchler, Ecology (1974) 55: pp. 586-604

Kansas Grazing land Water Quality Education Program Final Comprehensive Report, Ohlenbusch et al, 1997

Understanding Grass Growth, Steven S. Waller, Lowell E. Moser, Patrick E. Reece and George A. Gates, 1985.

Rangeland Cover Types of the United States, Society for Range Management 1994.

Vegetation and Cattle Responses to Different Intensities of Grazing on Short-Grass Ranges of the Central Great Plains, Technical Bulletin No. 1216, USDA. July 1960

History of the Native Vegetation of Western Kansas During Seven Years of Continuous Drought, F.W. Albertson, J.E. Weaver, Ecological Monographs, Vol. 12, No. 1 Jan. 1942 pp. 23-51

Deterioration of Midwestern Ranges, J. E. Weaver, F. W. Albertson, Ecology, Vol. 21, No. 2, April 1940, pp. 216-236

The effect of stocking rate on cattle gains and on native shortgrass vegetation in west-central Kansas, J. L. Launchbaugh, Bulletin 394, November 1957, Ft. Hays Branch, Kansas Ag. Exp. Station, Hays KS

The nature and properties of soils, Nyle C. Brady, Ray R. Weil- 14th ed, 2008.

Cleland, D.T., P.E. Avers, W.H. McNab, M.E. Jensen, R.G. Bailey, T. King, and W.E. Russell. 1997. National hierarchical framework of ecological units. In: M.S. Boyce and A. Haney (eds.) Ecosystem management applications for sustainable forest and wildlife resources, Yale University Press, New Haven, CT.

Fenneman, N.M., and D.W. Johnson. 1946. Physical divisions of the United States. U.S. Geological Survey, Physiographic Committee. Scale 1:700,000.

Quality control review: MLRA 72 Loamy Tableland Workshop, Goodland KS, September 21-22, 2015. David Kraft, Hutchinson, KS Area office 9-28-2015

Quality assurance review: David Kraft (acting QA for region 5 and 9)

Site Development and Testing Plan: This ESD went through the approval process.

Contributors

Chris Tecklenburg

Approval

Suzanne Mayne-Kinney, 6/06/2025

Acknowledgments

The ecological site development process is a collaborative effort, conceptual in nature, dynamic and is never considered complete. I thank all those who set the foundational work in the early 2000's in regard to this ESD. I thank all those who contributed to the development of this site. In advance, I thank those who would provide insight, comments and questions about this ESD in the future.

Rangeland health reference sheet

Interpreting Indicators of Rangeland Health is a qualitative assessment protocol used to determine ecosystem condition based on benchmark characteristics described in the Reference Sheet. A suite of 17 (or more) indicators are typically considered in an assessment. The ecological site(s) representative of an assessment location must be known prior to applying the protocol and must be verified based on soils and climate. Current plant community cannot be used to identify the ecological site.

Author(s)/participant(s)	Chris Tecklenburg Revision2 5-25-2021; Revision1 9-15-2015 David Kraft, John Henry, Doug Spencer and Dwayne Rice Original Authors 2-2005
--------------------------	--

Contact for lead author	Chris Tecklenburg (chris.tecklenburg@usda.gov)
Date	09/16/2015
Approved by	
Approval date	
Composition (Indicators 10 and 12) based on	Annual Production

Indicators

1. Number and extent of rills: None

2. Presence of water flow patterns: Typically none. If present (steeper slopes following intense storms) short and not connected.

3. Number and height of erosional pedestals or terracettes: None, due to the slope percentage and amount of cover. Pedestals and terracettes are indicators of soil being moved by water and/or wind.

4. Bare ground from Ecological Site Description or other studies (rock, litter, lichen, moss, plant canopy are not bare ground): Less than 5% bare ground is found on this site, with bare patches generally less than 2-3 inches in diameter. Extended drought can cause bare ground to increase. Bare ground is the remaining ground cover after accounting for ground surface covered by vegetation (basal and foliar canopy), litter, standing dead vegetation, gravel/rock, and visible biological crust (e.g., lichen, mosses, algae).

5. Number of gullies and erosion associated with gullies: None. Gullies are not a natural feature of this landscape and site.

6. Extent of wind scoured, blowouts and/or depositional areas: None. The vegetative cover in the reference state is sufficient to limit wind-scoured or blowout areas. This site is not a depositional area for offsite wind erosion.

7. Amount of litter movement (describe size and distance expected to travel): None. No evidence of litter movement (i.e., dead plant material that is in contact with the soil surface). The inherent capacity for litter movement on a soil is a function of its slope and landscape position.

8. Soil surface (top few mm) resistance to erosion (stability values are averages - most sites will show a range of values): Stability class ratings from the soil stability test should be =5 overall, with ratings of 5 or greater in the interspaces and under perennial plant canopy. Plant canopy is large enough to intercept the majority of raindrops. No physical crusts apparent.

9. Soil surface structure and SOM content (include type of structure and A-horizon color and thickness): The surface horizon (A) should be a minimum 7" (roots growing throughout) with weak fine granular to moderate fine subangular blocky structure. The surface (A) horizon color is very dark grayish brown 10YR 3/2 (moist), and the subsurface (B) horizon color is dark grayish brown 10YR 4/2 (moist).

10. Effect of community phase composition (relative proportion of different functional groups) and spatial distribution on infiltration and runoff: Diverse grass, forb, and shrub canopy and root structure reduces raindrop impact and slows overland flow providing increased time for infiltration to occur. Warm season shortgrasses are dominant. Tall and mid grasses and mid grass cool seasons are co-subdominants. Sedges, forbs, and shrubs are all minor components. Infiltration will decrease if tall and midgrasses and mid grass cool season groups become minor or trace. Spatial distribution of functional/structural groups varies from north to south. In the northern part of this MLRA expect more cool seasons and shrubs than in the southern end.

11. Presence and thickness of compaction layer (usually none; describe soil profile features which may be mistaken for compaction on this site): There is no evidence of a compacted soil layer less than 6 inches from the soil surface. Soil structure is similar to that described in Indicator 9.

12. Functional/Structural Groups (list in order of descending dominance by above-ground annual-production or live foliar cover using symbols: >>, >, = to indicate much greater than, greater than, and equal to):

Dominant: Group 1 Shortgrass Dominant 40% 680 lbs. blue grama 225-440, buffalograss 75-240

Sub-dominant: Group 2 Tall and mid warm season grasses co-subdominant 21% 360 lbs. sideoats grama 20-150, little bluestem 20-100, big bluestem 0-45, composite dropseed 0-45, sand dropseed 0-45 Group 3 Midgrass cool season co-subdominant 18% 300 lbs. western wheatgrass 200-270, green needlegrass 0-120, needle and thread 0-80, Fendler threeawn 0-15, squirreltail 0-15

Other: Group 4 Sedges minor component 6% 0-105 lbs. threadleaf sedge 0-50, sun sedge 0-30, sedge 0-25 Group 5 Forbs minor component 10% 170 lbs. see ESD for species list minimum 5 species. Group 6 Shrubs minor component 5% 85 lbs. see ESD for species list minimum 3 species.

Additional:

13. Amount of plant mortality and decadence (include which functional groups are expected to show mortality or decadence): The majority of plants are alive and vigorous. Some (<10%) mortality and decadence is expected for the site. This in part is due to drought, unexpected wildfire or a combination of the two events. This would be expected for both dominant and sub-dominant groups.

14. Average percent litter cover (%) and depth (in): Total litter cover is expected to be 60-80% and at an average depth of 0.25 inches. Litter may be reduced to 20-40% in cover and near zero depth for 1-2 years following wildfire or multiyear drought.

15. **Expected annual annual-production (this is TOTAL above-ground annual-production, not just forage annual-production):**
Annual production is 1,700 pounds per acre in a year with normal precipitation and temperatures. Low and high production years should yield 900 and 2,200 pounds per acre, respectively. Annual production may be reduced by 40-60% the first year following a wildfire or following a multiyear drought.
-

16. **Potential invasive (including noxious) species (native and non-native).** List species which BOTH characterize degraded states and have the potential to become a dominant or co-dominant species on the ecological site if their future establishment and growth is not actively controlled by management interventions. Species that become dominant for only one to several years (e.g., short-term response to drought or wildfire) are not invasive plants. Note that unlike other indicators, we are describing what is NOT expected in the reference state for the ecological site: Cheatgrass, Russian thistle, kochia, and other non-native annuals may occur in trace amounts in community 1.1 but they do not have the potential to increase to a subdominant or dominant in the absence of wildfire and act as an invasive on this site. The listed invasives are not expected to occur in the reference state.
-

17. **Perennial plant reproductive capability:** Plants in all functional/structural groups should be capable of reproducing annually under normal weather conditions. The number and distribution of tillers or rhizomes is assessed on perennial plants occupying the evaluation area. Vigor and reproductive capability may be somewhat reduced during drought or for 1 year following a wildfire. At least 50% of plants should still have reproductive capability during droughts that last 1-2 years.
-