

Ecological site R072XY104KS

Saline Lowland

Accessed: 09/05/2026

General information

Approved. An approved ecological site description has undergone quality control and quality assurance review. It contains a working state and transition model, enough information to identify the ecological site, and full documentation for all ecosystem states contained in the state and transition model.

MLRA notes

Major Land Resource Area (MLRA): 072X--Central High Tableland

Major Land Resource Area (MLRA) 72--Central High Tableland. This area is in Kansas (54 percent), Nebraska (25 percent), and Colorado (21 percent). A very small part of the area is in Wyoming. The area makes up about 34,550 square miles (89,535 square kilometers). It includes the towns of Garden City, Goodland, and Colby, Kansas; Imperial, North Platte, Ogallala, and Sidney, Nebraska; and Holyoke and Wray, Colorado. Interstate 70 bisects the area, and Interstates 76 and 80 follow the south side of the South and North Platte Rivers, respectively. The Cimarron National Grasslands occur in the southwest corner of the MLRA.

Classification relationships

Major land resource area (MLRA): 072-Central High Tableland

Ecological site concept

This site occurs on somewhat poorly drained soils in alluvium, on flood plains, and low stream terraces. This site consists of alluvial soils that usually contain a high concentration of salts. A seasonal or perennial water table greater than 6 feet from the surface characterizes this site. This ESD was developed using the existing ESDs developed in 2001: saline lowland North R072XA019KS, saline lowland South R072XB019KS and salt flat R072XY034CO. Changes were made in accordance with policy using the national ecological site handbook. Text changes were made in regards to clarity and style. Content was changed reflecting the range of variability of the site.

Associated sites

R072XY102KS	Saline Subirrigated Saline Subirrigated shares the same landform and parent material as Saline Lowland.
R072XY107KS	Sandy Lowland Sandy Lowland can be found adjacent to Saline Lowland site.

Similar sites

R072XY102KS	Saline Subirrigated Saline Subirrigated as a water table less than 6 feet from the surface and Saline Lowland has a water table greater than 6 feet. Plant communities can resemble one another.
-------------	--

Table 1. Dominant plant species

Tree	Not specified
Shrub	Not specified
Herbaceous	(1) <i>Sporobolus airoides</i> (2) <i>Pascopyrum smithii</i>

Physiographic features

The Saline Lowland site occurs on nearly level to moderately sloping floodplains, and low terraces adjacent to streams and rivers. This site may also occur on a Dune Field interdune. Salts have accumulated within the root zone of this site. This site receives runoff from areas higher on the landscape. The soils of this site are typically subject to rare flooding. Some soils in this site are not in positions that flood (interdune, terrace). NOTE: Saline lowland ecological site is characterized by a water table >6 feet from the surface. Haigler soil series associated with saline lowland can have a water table at 36".

Figure 1. MLRA 72 block diagram

Table 2. Representative physiographic features

Landforms	(1) Terrace (2) Flood plain (3) Interdune
Flooding duration	Brief (2 to 7 days)
Flooding frequency	None to rare
Ponding frequency	None
Elevation	760 – 1,140 m
Slope	0 %
Ponding depth	0 cm
Water table depth	90 – 200 cm

Climatic features

The average annual precipitation in this area is 14 to 25 inches (355 to 635 millimeters). It fluctuates widely from year to year. Most of the rainfall occurs as high-intensity, convective thunderstorms during the growing season. The maximum precipitation occurs from late spring through early autumn. Precipitation in winter occurs as snow. The annual snowfall ranges from about 16 inches (40 centimeters) in the southern part of the area, to 35 inches (90 centimeters) in the northern part. The average annual temperature is 46 to 57 degrees F (8 to 14 degrees C). The freeze-free period averages 161 days and ranges from 135 to 210 days, increasing in length from northwest to southeast. Climate data comes from the Natural Resources Conservation Service (NRCS) National Water and Climate Center. The data

set is from 1981-2010.

Table 3 Representative climatic features

Frost-free period (average)	150 days
Freeze-free period (average)	170 days
Precipitation total (average)	510 mm

- (1) SCOTT CITY [USC00147271], Scott City, KS
- (2) SHARON SPRINGS [USC00147397], Sharon Springs, KS
- (3) SYRACUSE 1NE [USC00148038], Syracuse, KS
- (4) HAIGLER [USC00253515], Haigler, NE
- (5) TRENTON DAM [USC00258628], Trenton, NE
- (6) GARDEN CITY EXP STN [USC00142980], Garden City, KS
- (7) CULBERTSON [USC00252065], Culbertson, NE

Influencing water features

A seasonal water table, (approximately greater than 6 feet) and adjacent perennial water table have influence on the kinds and amounts of vegetation on this site.




Figure 6. Fig. 7-1 from National Rangeland and Pasture Handb

Soil features

This site consists of very deep saline and saline-alkali soils. The high concentration of salts affect both the kind and amount of vegetation present. These soils generally do not have free water within 60 inches of the surface, but subsoil and underlying layers may have a seasonal high water table below a depth of about 36 inches in the Haigler series. Laird soils once had a higher water table, but presently the water table has dropped between 15 to 50 feet below the soil surface. The surface soil ranges from 3 to 20 inches thick. It is dark colored in some of these series and light colored in others. Texture of the surface soil ranges widely from silty clay loam to loamy fine

sand. The underlying material is lighter colored than the surface soil, and occasionally has redoximorphic concentrations in the lower part. It ranges widely in texture. The soils in this site are commonly calcareous to the surface. The excessive saline and alkali salts present affect other soil properties such as available water capacity, soil structure, infiltration, and permeability. Soils in this site generally have low to moderate content of organic matter.

Slick spots are high sodium areas that contain no vegetation. They are inherent to the site and are intermixed with areas of vegetation.

Major soil series correlated to this ecological site include Las, Haigler, Laird, Caruso, and Arvada.

Other soil series that have been correlated to this site include Craft, Las Animas, Lincoln, Bridgeport, Sanborn, and Saltine.

These attributes represent 0-40 inches in depth or to the first restrictive layer.

Figure 7. Las soil (OSD) dendrogram

Table 4. Representative soil features

Surface texture	(1) Silty clay loam (2) Silt loam (3) Loamy fine sand
Family particle size	(1) Loamy
Drainage class	Moderately well drained to well drained
Permeability class	Very slow to moderately rapid
Soil depth	150 – 200 cm
Surface fragment cover <=3"	Not specified
Surface fragment cover >3"	Not specified
Available water capacity (0-101.6cm)	6.76 – 21.82 cm
Calcium carbonate equivalent (0-101.6cm)	0 – 40 %
Electrical conductivity (0-101.6cm)	0 – 20 mmhos/cm

Sodium adsorption ratio (0-101.6cm)	0 – 130
Soil reaction (1:1 water) (0-101.6cm)	5.6 – 10
Subsurface fragment volume <=3" (Depth not specified)	Not specified
Subsurface fragment volume >3" (Depth not specified)	Not specified

Ecological dynamics

The plant community for this site is dynamic due to the complex interaction of many ecological processes. The interpretive plant community for this site is the Reference Plant Community. This community has been determined by the study of rangeland relic areas, areas protected from excessive disturbance, and areas under long-term rotational grazing strategies. Trends in plant community dynamics ranging from heavily grazed to lightly grazed areas, seasonal use pastures, and historical accounts have also been used.

The Saline Lowland site developed with occasional fires as part of the ecological processes. Historically, it is believed that the fires were infrequent, randomly distributed, and started by lightning at various times throughout the season when thunderstorms were likely to occur. It is also believed that pre-European inhabitants may have used fire as a management tool for attracting herds of large migratory herbivores (bison, elk, deer, and pronghorn). The impact of fire over the past 100 years has been relatively insignificant due to the human control of wildfires, and the lack of acceptance of prescribed fire as a management tool in the semi-arid, High Plains area.

This site occurs on terraces adjacent to streams or rivers, floodplains, and the walls of closed depressions. Historically, the flooding potential on these sites range from none to rare.

Soils of this site are characterized by moderate to high salinity, or alkaline conditions. There will often be white or gray deposits on the soil surface early in the spring or following wet periods. Slick spots (bare exposed areas high in sodium) are an inherent characteristic occupying less than 3 percent of the community. The degree of salinity or alkali condition has a major influence on the plant community that will persist on this site. Plants that can tolerate these conditions include alkali sacaton, western wheatgrass, and inland saltgrass.

Continuous grazing without adequate recovery periods on this site will result in the loss of vigor and reproductive potential for such species as alkali sacaton and western wheatgrass. Inland saltgrass and blue grama will increase. Once the plant community has shifted to inland saltgrass/blue grama sod, it is very difficult to reverse the shift.

The degree of herbivory (feeding on herbaceous plants) has a significant impact on the dynamics of the site. Historically, periodic grazing by herds of large migratory herbivores was a primary influence. Secondary influences of herbivory by species such as prairie dogs, grasshoppers, gophers, and root feeding organisms impacted the vegetation historically, and continue to this day.

The management of herbivory by humans through grazing of domestic livestock and/or manipulation of wildlife populations has been a major influence on the ecological dynamics of the site. This management coupled with the High Plains climate largely dictates the plant communities for the site.

Typically, drought cycles have not had a major impact upon the vegetation due to the presence of a water table. Depletion of ground water reserves may change species composition and possibly cause a shift in ecological sites.

Growth of native cool season plants begins about April 15, and continues to about June 15. Native warm season plants begin growth about May 15, and continue to about August 15. Green up of cool season plants may occur in September and October if adequate moisture is available.

The following diagram illustrates the common plant communities that can occur on the site and the transition pathways (arrows) among communities. Bold lines surrounding each state represent ecological thresholds. The ecological processes are discussed in more detail in

the plant community descriptions following the diagram.

State and transition model

Additional community tables

Table 5. Community 1.1 plant community composition

Group	Common Name	Symbol	Scientific Name	Annual Production ()	Foliar Cover (%)
Grass/Grasslike					
1	Tall-mid warm season Dominant 43%			336-1059	
	alkali sacaton	SPAI	<i>Sporobolus airoides</i>	336-673	–
	saltgrass	DISP	<i>Distichlis spicata</i>	56-336	–
	switchgrass	PAVI2	<i>Panicum virgatum</i>	0-78	–
	sand dropseed	SPCR	<i>Sporobolus cryptandrus</i>	0-39	–
	alkali cordgrass	SPGR	<i>Spartina gracilis</i>	0-39	–
2	Mid-short cool season grasses Sub-dominant 32%			336-790	
	western wheatgrass	PASM	<i>Pascopyrum smithii</i>	336-673	–
	slender wheatgrass	ELTR7	<i>Elymus trachycaulus</i>	0-56	–
	Canada wildrye	ELCA4	<i>Elymus canadensis</i>	0-39	–
	squirreltail	ELELE	<i>Elymus elymoides ssp. elymoides</i>	0-22	–
	Nuttall's alkaligrass	PUNU2	<i>Puccinellia nuttalliana</i>	0-22	–
	little barley	HOPU	<i>Hordeum pusillum</i>	0-22	–
	green needlegrass	NAVI4	<i>Nassella viridula</i>	0-22	–
3	Short warm season Minor component 5%			0-135	
	buffalograss	BODA2	<i>Bouteloua dactyloides</i>	0-73	–
	blue grama	BOGR2	<i>Bouteloua gracilis</i>	0-73	–
	ring muhly	MUTO2	<i>Muhlenbergia torreyi</i>	0-22	–
4	Mid warm season Minor component 3%			0-78	
	sideoats grama	BOCU	<i>Bouteloua curtipendula</i>	0-45	–
	scratchgrass	MUAS	<i>Muhlenbergia asperifolia</i>	0-17	–
	Fendler threeawn	ARPUL	<i>Aristida purpurea var. longiseta</i>	0-17	–
5	Sedges-rushes Trace component 2%			0-56	
	sun sedge	CAINH2	<i>Carex inops ssp. heliophila</i>	0-28	–
	sedge	CAREX	<i>Carex</i>	0-28	–
	rush	JUNCU	<i>Juncus</i>	0-28	–
Forb					
6	Forbs Minor component 10%			22-252	
	white heath aster	SYERE	<i>Symphyotrichum ericoides var. ericoides</i>	0-22	–
	common dandelion	TAOF	<i>Taraxacum officinale</i>	0-22	–
	Cuman ragweed	AMPS	<i>Ambrosia psilostachya</i>	0-22	–
	white sagebrush	ARLUM2	<i>Artemisia ludoviciana ssp. mexicana</i>	0-22	–
	twogrooved milkvetch	ASBI2	<i>Astragalus bisulcatus</i>	0-22	–
	Maximilian sunflower	HEMA2	<i>Helianthus maximiliani</i>	0-22	–
	rush skeletonplant	LYJU	<i>Lygodesmia juncea</i>	0-22	–
	scarlet beeblossom	OESU3	<i>Oenothera suffrutescens</i>	0-22	–
	leafy false goldenweed	OOFOF	<i>Oenopsis foliosa var. foliosa</i>	6-17	–
	scarlet globemallow	SPCO	<i>Sphaeralcea coccinea</i>	6-17	–
	American vetch	VIAM	<i>Vicia americana</i>	6-17	–

	desert princesplume	STPIP	<i>Stanleya pinnata</i> var. <i>pinnata</i>	0-11	–
	upright prairie coneflower	RACO3	<i>Ratibida columnifera</i>	0-11	–
	lacy tansyaster	MAPIP4	<i>Machaeranthera pinnatifida</i> ssp. <i>pinnatifida</i> var. <i>pinnatifida</i>	0-11	–
	hairy false goldenaster	HEVI4	<i>Heterotheca villosa</i>	0-11	–
	broadleaf milkweed	ASLA4	<i>Asclepias latifolia</i>	0-11	–
	scrambled eggs	COAU2	<i>Corydalis aurea</i>	0-11	–
	purple prairie clover	DAPUP	<i>Dalea purpurea</i> var. <i>purpurea</i>	0-11	–
	fetid marigold	DYPA	<i>Dyssodia papposa</i>	0-11	–
	curlycup gumweed	GRSQ	<i>Grindelia squarrosa</i>	0-11	–
Shrub/Vine					
7	Shrubs Minor component 5%			0-112	
	broom snakeweed	GUSA2	<i>Gutierrezia sarothrae</i>	0-45	–
	devil's-tongue	OPHU	<i>Opuntia humifusa</i>	0-45	–
	plains pricklypear	OPPO	<i>Opuntia polyacantha</i>	0-45	–
	fragrant sumac	RHAR4	<i>Rhus aromatica</i>	0-22	–
	willow	SALIX	<i>Salix</i>	0-22	–
	winterfat	KRLA2	<i>Krascheninnikovia lanata</i>	0-22	–
	fourwing saltbush	ATCA2	<i>Atriplex canescens</i>	0-22	–
	rubber rabbitbrush	ERNAG	<i>Ericameria nauseosa</i> ssp. <i>nauseosa</i> var. <i>glabrata</i>	0-22	–

Table 6. Community 1.2 plant community composition

Group	Common Name	Symbol	Scientific Name	Annual Production ()	Foliar Cover (%)
-------	-------------	--------	-----------------	----------------------	------------------

Table 7. Community 2.1 plant community composition

Group	Common Name	Symbol	Scientific Name	Annual Production ()	Foliar Cover (%)
-------	-------------	--------	-----------------	----------------------	------------------

Table 8. Community 3.1 plant community composition

Group	Common Name	Symbol	Scientific Name	Annual Production ()	Foliar Cover (%)
-------	-------------	--------	-----------------	----------------------	------------------

Table 9. Community 3.2 plant community composition

Group	Common Name	Symbol	Scientific Name	Annual Production ()	Foliar Cover (%)
-------	-------------	--------	-----------------	----------------------	------------------

Animal community

Grazing by domestic livestock is one of the major income-producing industries in the area. Rangelands in this area provide yearlong forage under prescribed grazing for cattle, sheep, horses, and other herbivores. During the dormant period, livestock may need supplementation based on reliable forage analysis. Grazing Interpretations Calculating safe stocking rates: Proper stocking rates should be incorporated into a grazing management strategy that protects the resource, maintains, or improves rangeland health, and is consistent with management objectives. In addition to usable forage, safe stocking rates should consider ecological condition, trend of the site, past grazing use history, season of use, stock density, kind and class of livestock, forage digestibility, forage nutritional value, variation of harvest efficiency based on desirability preference of plant species, and/or grazing system and site grazability factors (such as steep slopes, site inaccessibility, or distance to drinking water). Often the current plant community does not entirely match any particular community phase as described in this ecological site description. Because of this, a resource inventory is necessary to document plant composition and production. Proper interpretation of inventory data will permit the establishment of a safe initial stocking rate. No two years have exactly the same weather conditions. For this reason, year-to-year and season-to season fluctuations in forage production are to be expected on grazing lands. Livestock producers must make timely adjustments in the numbers of animals or in the length of grazing periods to avoid overuse of forage plants when production is unfavorable, and to make advantageous adjustments when forage supplies are above average. Initial stocking rates should be improved through the use of vegetation monitoring and actual use records that include number and type of livestock, the timing and duration of grazing, and utilization levels. Actual use records over time will assist in making stocking rate adjustments based on the variability factors. Average annual production must be measured or estimated to properly assess usable forage production and stocking rates.

Hydrological functions

Water is the principal factor limiting forage production on this site. Infiltration is moderate to slow and runoff potential for this site varies from moderate to high depending on soil hydrologic group and ground cover. In many cases, areas with greater than 75 percent ground cover have the greatest potential for high infiltration and lower runoff. An example of an exception would be where shortgrasses form a strong sod and dominate the site. Areas where ground cover is less than 50 percent have the greatest potential to have reduced infiltration and higher runoff. (Refer to NRCS Section 4, National Engineering Handbook (NEH-4) for runoff quantities and hydrologic curves).

Recreational uses

This site provides hunting, hiking, photography, bird watching and other opportunities. The wide varieties of plants that bloom from spring until fall have an esthetic value that appeals to visitors.

Wood products

No appreciable wood products are present on the site.

Other products

None noted.

Other information

Site Development and Testing Plan. Future work (for approved ESD) includes field visits to verify ecological site concepts with field staff. Field staff include but not limited to project office leader, area soil scientist, state soil scientist, ecological site specialist, state rangeland conservationist, area rangeland management specialist, and local field personnel. This site includes collaboration between Kansas, Colorado and Nebraska. Field visits are to be determined by spatial extent of the site as well as personal knowledge of the site. Activity during field visits will include but not limited to: identifying the soil, landform, plant community, and verifying existing site concepts.

Inventory data references

Information presented here has been derived from NRCS clipping data, numerous ocular estimates, and other inventory data. Field observations from experienced range personnel was used extensively to develop this ecological site description. Those involved in developing the Saline Lowland ecological site North include Harvey Sprock from Colorado; Carol Eakins, Chuck Markley, Jeff Nichols, and Mary Schrader from Nebraska; Joan Gienger and Ted Houser from Kansas. Those involved in developing Saline Lowland South include: Tim Watson, Amanda Shaw, Susan Francis, Jon Deege, Harvey Sprock, Robert Schiffner, and Josh Saunders. Range Condition Guides and Technical Range Site Descriptions for Kansas, Saline Lowland, USDA, Soil Conservation Service, August, 1967. Range Site Description for Kansas, Saline Lowland, USDA-Soil Conservation Service, September, 1983. Range Site Description for Colorado, Salt Flat, USDA-Soil Conservation Service, December 1975. Guide for determining range condition and suggestive initial stocking rates for Nebraska, Saline Lowland, Vegetative Zone 1 and II, USDA-Soil Conservation Service, April 1983. Range Site Description for Nebraska, Saline Lowland, USDA-Soil Conservation Service, August, 1981 Schacht, Walter H., Larsen, Dana. Section III. Range Sites, Saline Lowland Range Site, The Board of Regents of the University of Nebraska, publication. Ecological Site Description for Kansas, Saline Lowland North (R072XA019KS) and South (R072XB019KS), located in Ecological Site Information System (ESIS), 2007. Ecological Site Description for Colorado, Salt Flat (R072XY034CO), located in Ecological Site Information System (ESIS).

Other references

High Plains Regional Climate Center, University of Nebraska, 830728 Chase Hall, Lincoln, NE 68583-0728. (<http://hpcc.unl.edu>)

USDA, NRCS. National Water and Climate Center, 101 SW Main, Suite 1600, Portland, OR 97204-3224. (<http://wcc.nrcs.usda.gov>)

USDA, NRCS. National Range and Pasture Handbook, September 1997

USDA, NRCS. National Soil Information System, Information Technology Center, 2150 Centre Avenue, Building A, Fort Collins, CO 80526. (<http://nasis.nrcs.usda.gov>)

USDA, NRCS. 2002. The PLANTS Database, Version 3.5 (<http://plants.usda.gov>). National Plant Data Center, Baton Rouge, LA 70874-4490 USA.

Contributors

Chris Tecklenburg

Acknowledgments

The ecological site development process is a collaborative effort, conceptual in nature, dynamic, and is never considered complete. I thank all those who set the foundational work in the early 2000s in regards to this ESD. I thank all those who contributed to the development of this site. In advance, I thank those who would provide insight, comments and questions about this ESD in the future.

Rangeland health reference sheet

Interpreting Indicators of Rangeland Health is a qualitative assessment protocol used to determine ecosystem condition based on benchmark characteristics described in the Reference Sheet. A suite of 17 (or more) indicators are typically considered in an assessment. The ecological site(s) representative of an assessment location must be known prior to applying the protocol and must be verified based on soils and climate. Current plant community cannot be used to identify the ecological site.

Author(s)/participant(s)	Update 3-08-2016 Chris Tecklenburg. Original authors Feb 2005 David Kraft, John Henry, Doug Spencer, and Dwayne Rice and from Harvey Sprock, Dan Nosal, Blake Hendon on ES formerly known as Salt Flat 1-19-2005.
Contact for lead author	Chris Tecklenburg chris.tecklenburg@ks.usda.gov and David Kraft david.kraft@ks.usda.gov
Date	03/08/2016
Approved by	
Approval date	
Composition (Indicators 10 and 12) based on	Annual Production

Indicators

1. Number and extent of rills: None

2. Presence of water flow patterns: None where vegetation is continuous. Slick spots (high sodium areas) can pond water and concentrate overland flow. Flow paths should be short in length and disconnected.

3. Number and height of erosional pedestals or terracettes: There is no evidence of pedestaled plants or terracettes on the site.

4. Bare ground from Ecological Site Description or other studies (rock, litter, lichen, moss, plant canopy are not bare ground):
Less than 5% bare ground is found on this site. Bare areas can range from 3-4 inches in diameter. Extended drought may cause bare ground to increase up to 10%. Slick spots occur on the site and support some vegetation.
-

5. Number of gullies and erosion associated with gullies: None
-

6. Extent of wind scoured, blowouts and/or depositional areas: None
-

7. Amount of litter movement (describe size and distance expected to travel): None
-

8. Soil surface (top few mm) resistance to erosion (stability values are averages - most sites will show a range of values):
Plant canopy is large enough to intercept the majority of raindrops. A soil fragment will not “melt” or lose its structure when immersed in water for 30 seconds. There is no evidence of pedestaled plants or terracettes. Soil stability scores will range from 3-4.
-

9. Soil surface structure and SOM content (include type of structure and A-horizon color and thickness): SOM ranges from 1-3 % . Soils are deep well drained, sodic, saline, and strongly alkaline. Surface texture ranges from clay loam to sandy loam. A-horizon color is grayish brown (10YR 4/2) moist at 0-12 inches in depth. Weak granular structure.
-

10. Effect of community phase composition (relative proportion of different functional groups) and spatial distribution on infiltration and runoff: There is no negative effect on water infiltration and/or runoff due to plant composition or distribution. Diverse grass, forb, shrub functional/structural groups and diverse root structure/patterns reduces raindrop impact that slows overland flow providing increased time for infiltration to occur.
-

11. Presence and thickness of compaction layer (usually none; describe soil profile features which may be mistaken for compaction on this site): None
-

12. Functional/Structural Groups (list in order of descending dominance by above-ground annual-production or live foliar cover using symbols: >>, >, = to indicate much greater than, greater than, and equal to):

Dominant: Two groups of sub-dominant when combined make up 64% of the plant community

Sub-dominant: tallgrass warm season (alkali sacaton 300-600) = mid-short cool season (western wheatgrass 300-600) > midgrass warm season (inland saltgrass 50-300)

Other: Minor—Forbs > shortgrass warm season (blue grama, buffalograss) = Shrubs > sedges and rushes

Additional:

13. Amount of plant mortality and decadence (include which functional groups are expected to show mortality or decadence):
The majority of plants are alive and vigorous. Some mortality and decadence is expected for the site. This in part is due to drought, unexpected wildfire or a combination of the two events. This would be expected for both dominant and sub-dominant groups.
-
14. Average percent litter cover (%) and depth (in): Plant litter is distributed evenly throughout the site. When prescribed burning is practiced there will be little litter the first half of the growing season. 35-50% litter cover at 0.25-0.50 inch depth. Litter cover during and following extended drought ranges from 25-35%.
-
15. Expected annual annual-production (this is TOTAL above-ground annual-production, not just forage annual-production):
1000 lbs./ac. low precipitation years, 2200 lbs./ac. average precipitation years, 3000 lbs./ac. high precipitation years. After extended drought or the first growing season following wildfire, production may be significantly reduced by 500 – 700 lbs./ac.
-
16. Potential invasive (including noxious) species (native and non-native). List species which BOTH characterize degraded states and have the potential to become a dominant or co-dominant species on the ecological site if their future establishment and growth is not actively controlled by management interventions. Species that become dominant for only one to several years (e.g., short-term response to drought or wildfire) are not invasive plants. Note that unlike other indicators, we are describing what is NOT expected in the reference state for the ecological site: Invasive plants should not occur in reference plant community. However Russian thistle, kochia or other non-native alkali tolerant species may invade following extended drought assuming a seed source is available. Inland saltgrass is the major native (non-invasive) increaser on this site, but rabbitbrush and the muhlys may also increase.
-
17. Perennial plant reproductive capability: The only limitations are weather-related, wildfire, natural disease, and insects that may temporarily reduce reproductive capability.
-