

Ecological site R080BY153TX

Loamy Sand

26-33" PZ

Last updated: 9/19/2023
Accessed: 06/22/2026

General information

Provisional. A provisional ecological site description has undergone quality control and quality assurance review. It contains a working state and transition model and enough information to identify the ecological site.

MLRA notes

Major Land Resource Area (MLRA): 080B–Texas North-Central Prairies

MLRA 80B consists of gently rolling, dissected plains with very steep hillsides and sideslopes and narrow flood plains associated with small streams. Loamy and clayey soils range from very shallow to deep and developed in sandstones, shales, and limestones of Pennsylvanian age.

Classification relationships

This ecological site is correlated to soil components at the Major Land Resource Area (MLRA) level which is further described in USDA Ag Handbook 296.

Ecological site concept

These sites occur over deep non calcareous loamy sand soils on uplands. The reference vegetation includes native perennial tallgrasses with numerous forbs and scattered oaks. Woody canopy cover is typically under thirty percent under reference conditions. Without periodic fire or brush management, woody species may increase and dominate the site.

Associated sites

R080BY158TX	SANDY 26-33" PZ These sites frequently occur adjacent to each other or may be intermingled in the same landscape position. Deep sandy soils.
R080BY159TX	Sandy Loam 26-33" PZ These sites frequently occur adjacent to each other or may be intermingled in the same landscape position. Sandy loam soils.

Similar sites

R080BY159TX	Sandy Loam 26-33" PZ Similar species, greater proportion of tallgrasses, fewer trees. Slightly more production potential.
--------------------	---

R080BY158TX	<p>SANDY 26-33" PZ</p> <p>Similar species, lower proportions of tallgrasses and forbs. Greater proportion of midgrasses. Somewhat lower production potential.</p>
--------------------	--

Table 1. Dominant plant species

Tree	(1) <i>Quercus stellata</i> (2) <i>Quercus marilandica</i>
Shrub	Not specified
Herbaceous	(1) <i>Sorghastrum nutans</i> (2) <i>Andropogon gerardii</i>

Physiographic features

This site occurs on linear to concave base slopes and interfluves of dip slopes and on treads of stream terraces in the Texas North-Central Prairies. Slopes are typically less than percent.

Table 2. Representative physiographic features

Landforms	(1) Hills > Ridge (2) Hills > Dip slope (3) Alluvial plain > Stream terrace
Runoff class	Very low to medium
Elevation	230 – 730 m
Slope	0 – 10 %
Aspect	Aspect is not a significant factor

Climatic features

The climate is subtropical subhumid and is characterized by hot humid summers and relatively mild winters. Tropical maritime air controls the climate during spring, summer and fall. In winter and early spring, frequent surges of polar Canadian air cause sudden drops in temperatures and add considerable variety to the daily weather. The average first frost generally occurs about November 5 and the last freeze of the season usually occurs about March 19. The average frost free period ranges from 215 days in the northern counties, to 240 days in the south.

The average relative humidity in mid-afternoon is about 60 percent in the summer months. Humidity is higher at night, and the average at dawn is about 80 percent. The sun shines 75 percent of the time possible during the summer and 50 percent in winter. The prevailing wind direction is from the southwest and highest windspeeds occur during the spring months.

Approximately 75% of annual rainfall occurs between April 1 and October 31. Rainfall during the months of April through September typically occurs during thunderstorms which tend to be intense and brief, resulting in large amounts of rain in a short time. The wettest months of the year are May, June, September, and October. The driest months during the growing season are July and August. The

winter months of November, December, January, and February are the driest months overall.

Average annual precipitation for the entire MLRA is approximately 28 inches. There is a noticeable difference in the average annual precipitation in the northern counties in comparison to the southern and western counties of this Major Land Resource Area. Jack, Clay, Young, and Palo Pinto Counties all have an average annual precipitation of more than 31 inches. Stephens, Eastland, McCulloch, and San Saba Counties all have an average annual precipitation of less than 28 inches.

Winters tend to be mild, with occasional periods of very cold temperatures which can be accompanied by strong northerly winds and freezing precipitation. Snow is infrequent and significant accumulations are rare. These periods of very cold weather are generally short-lived. Summers tend to be hot and dry. Drought conditions are common during most summers. Air temperatures of more than 95oF are common from mid-June through September. In the northern counties nearest to the Red River, temperatures are generally slightly cooler during winter months and slightly warmer during summer months than in the other counties in the North Central Prairie.

Table 3 Representative climatic features

Frost-free period (characteristic range)	180-200 days
Freeze-free period (characteristic range)	210-230 days
Precipitation total (characteristic range)	760-810 mm
Frost-free period (actual range)	180-200 days
Freeze-free period (actual range)	210-230 days
Precipitation total (actual range)	740-840 mm
Frost-free period (average)	190 days
Freeze-free period (average)	220 days
Precipitation total (average)	790 mm

- (1) SAN SABA 7NW [USC00417994], Richland Springs, TX
- (2) BROWNWOOD 2ENE [USC00411138], Early, TX
- (3) EASTLAND [USC00412715], Eastland, TX
- (4) MINERAL WELLS AP [USW00093985], Millsap, TX
- (5) BRECKENRIDGE [USC00411042], Breckenridge, TX
- (6) GRAHAM [USC00413668], Graham, TX
- (7) JACKSBORO [USC00414517], Jacksboro, TX

Influencing water features

These sites may receive some runoff from other sites but also shed some water via overland flow. However, the presence of good ground cover and deep rooted grasses can help facilitate water infiltration into the soil. Theses sites are not associated with wetlands.

Wetland description

NA




Figure 7.

Soil features

Representative soil components for this ecological site include: Decordova, Newcastle, Vashti

The site is characterized by moderately deep to very deep loamy sand soils with a high infiltration rate.

Table 4. Representative soil features

Parent material	(1) Residuum – sandstone (2) Alluvium – sandstone (3) Residuum – claystone (4) Alluvium – claystone
Surface texture	(1) Loamy fine sand (2) Stony loamy fine sand
Drainage class	Moderately well drained to well drained
Permeability class	Slow to moderately rapid
Soil depth	50 cm
Surface fragment cover <=3"	Not specified

Surface fragment cover >3"	Not specified
Available water capacity (0-101.6cm)	10.16 – 17.78 cm
Electrical conductivity (0-101.6cm)	Not specified
Sodium adsorption ratio (0-101.6cm)	Not specified
Soil reaction (1:1 water) (0-101.6cm)	5.6 – 7.3
Subsurface fragment volume <=3" (Depth not specified)	0 – 10 %
Subsurface fragment volume >3" (Depth not specified)	Not specified

Ecological dynamics

The reference plant community for the Loamy Sand ecological site is a tallgrass/midgrass oak savanna. Evidence of the historic vegetation can be found in the journals and records of explorers, military expeditions, and boundary survey teams.

Climate is a major factor influencing vegetation on the site. Long-term droughts lasting multiple years or growing seasons are infrequent, but when they do occur, they can have a negative impact on the vegetation. If abusive grazing occurs during or immediately following the drought period, the results can be devastating. The effects of erratic seasonal moisture and short-term dry spells lasting a few months are not as severe as those caused by long-term droughts. However, the lower the ecological status of the site, the greater the negative impact will be during drought periods regardless of duration.

Fire was an important part of the ecosystem. Most ecosystems in the North Central Prairie developed in a 4 to 6 year regime of recurring fires. Many of these fires resulted from lightning strikes during thunderstorms. Native Americans frequently set fires to manipulate the movement of bison and other animals as well as a defensive or offensive technique when dealing with their enemies. These historic fires were usually severe because of the amount of grass fuel available to carry the fire. The intensity of fires kept shrubs and sapling trees suppressed and allowed grasses and forbs to flourish. Tallgrass species are fire tolerant and are enhanced by periodic burning. Forbs usually increase for a year or two following these fires before the grasses become dominant again.

Prior to settlement, this site was subject to periodic grazing and browsing by vast herds of bison, wild cattle, wild horses, and deer. At times these grazing and browsing episodes were intense and severe, but periods of heavy use were followed by long periods of non-use as the herds migrated to fresh grazing areas before returning to previously grazed areas. The grazed areas had an opportunity to rest, regrow, regain vigor, and reproduce prior to the next grazing event. Intervals between grazing periods were frequently influenced by the amount of time that had elapsed since the last fire on the area.

As the region was settled, fire was reduced or eliminated and grasslands were fenced off to control movement and facilitate grazing by domestic livestock. As a result of abusive grazing or lack of grazing and/or the elimination of fire, in association with extreme climatic events, the tallgrass plant community has been eliminated or significantly altered on most Loamy Sand sites.

Lack of fire allows herbaceous vegetation to become senescent and may eventually lead to the loss of the most desirable species. The density and canopy of oaks, elms, shrubs, and greenbriar increase significantly in the absence of fire. Seedlings of non-native brush species and invasive weeds may encroach on the site from adjacent sites

Further deterioration leads to the loss of the perennial warm-season midgrass and forb plant community and an increase in short grasses, annuals, and bare ground. This provides the opportunity for woody species such as mesquite and juniper to encroach from adjacent sites.

Selective individual removal of unwanted trees and shrubs is relatively easy and more practical when brush plants initially appear on the site. The increase of brush can be fairly rapid and the plants per acre will soon become too numerous for individual control to be feasible. Once woody plants become mature or develop into dense stands, control is expensive, uneconomical, impractical, and difficult to achieve. Brush management is most successful using a systems approach. Initial treatment by mechanical methods can be followed by using approved herbicides, and using prescribed fire as a maintenance technique. Prescribed grazing with a reasonable stocking rate can sustain the grass species composition and production at a near reference community level.

Changes in plant communities and vegetation states on the Loamy Sand site are result of the combined influences of natural events (rainfall, temperature, droughts, etc.) and the accompanying management systems implemented on the area (prescribed fire, grazing management, and brush management).

Rangeland Health Reference Worksheets have been posted for this site on the Texas NRCS website (www.tx.nrcs.usda.gov) in Section II of the eFOTG under (F) Ecological Site Descriptions.

State and Transitional Pathways:

The State and Transition Diagram which follows provides information on some of the most typical pathways that the vegetation on this site can follow as the result of natural events, management inputs, and application of conservation treatments. There may be other plant communities that can exist on this site under certain conditions. Consultation with local experts and professionals is recommended prior to application of practices or management strategies in order to ensure that specific objectives will be met.

State and transition model

Additional community tables

Table 5. Community 1.1 plant community composition

Group	Common Name	Symbol	Scientific Name	Annual Production ()	Foliar Cover (%)
Grass/Grasslike					
1	Tallgrass			560-1681	
	little bluestem	SCSC	<i>Schizachyrium scoparium</i>	560-1681	–
2	Tallgrasses			112-1681	
	big bluestem	ANGE	<i>Andropogon gerardii</i>	112-1681	–
	switchgrass	PAVI2	<i>Panicum virgatum</i>	0-1681	–
	Indiangrass	SONU2	<i>Sorghastrum nutans</i>	112-1681	–
	purpletop tridens	TRFL2	<i>Tridens flavus</i>	0-560	–
3	Midgrasses			560-1121	
	sideoats grama	BOCU	<i>Bouteloua curtipendula</i>	280-1121	–
	sand lovegrass	ERTR3	<i>Eragrostis trichodes</i>	280-1121	–
	Texas wintergrass	NALE3	<i>Nassella leucotricha</i>	0-280	–
	Texas bluegrass	POAR	<i>Poa arachnifera</i>	0-280	–
	yellow foxtail	SEPUP2	<i>Setaria pumila ssp. pumila</i>	0-280	–
	composite dropseed	SPCOC2	<i>Sporobolus compositus var. compositus</i>	112-280	–
	Drummond's dropseed	SPCOD3	<i>Sporobolus compositus var. drummondii</i>	0-280	–
	sand dropseed	SPCR	<i>Sporobolus cryptandrus</i>	112-280	–
	silver beardgrass	BOLAT	<i>Bothriochloa laguroides ssp. torreyana</i>	112-280	–
	Arizona cottontop	DICA8	<i>Digitaria californica</i>	0-280	–
	Canada wildrye	ELCA4	<i>Elymus canadensis</i>	0-280	–

	plains lovegrass	ERIN	<i>Eragrostis intermedia</i>	0-280	-
	fringed signalgrass	URCI	<i>Urochloa ciliatissima</i>	0-112	-
	thin paspalum	PASE5	<i>Paspalum setaceum</i>	0-112	-
4	Mid/shortgrasses			112-560	
	blue grama	BOGR2	<i>Bouteloua gracilis</i>	0-280	-
	Scribner's rosette grass	DIOLS	<i>Dichanthelium oligosanthes var. scribnerianum</i>	0-280	-
	hooded windmill grass	CHCU2	<i>Chloris cucullata</i>	0-280	-
	tumble windmill grass	CHVE2	<i>Chloris verticillata</i>	0-168	-
	fall witchgrass	DICO6	<i>Digitaria cognata</i>	0-168	-
	red lovegrass	ERSEO	<i>Eragrostis secundiflora ssp. oxylepis</i>	0-168	-
	curly-mesquite	HIBE	<i>Hilaria belangeri</i>	0-168	-
	hairy grama	BOHI2	<i>Bouteloua hirsuta</i>	0-168	-
	purple threeawn	ARPUP6	<i>Aristida purpurea var. purpurea</i>	0-168	-
	Wright's threeawn	ARPUW	<i>Aristida purpurea var. wrightii</i>	0-168	-
	buffalograss	BODA2	<i>Bouteloua dactyloides</i>	0-168	-
Forb					
5	Forbs			280-560	
	Cuman ragweed	AMPS	<i>Ambrosia psilostachya</i>	0-280	-
	Engelmann's daisy	ENPE4	<i>Engelmannia peristenia</i>	0-280	-
	Maximilian sunflower	HEMA2	<i>Helianthus maximiliani</i>	0-280	-
	partridge pea	CHFA2	<i>Chamaecrista fasciculata</i>	0-112	-
	white sagebrush	ARLUM2	<i>Artemisia ludoviciana ssp. mexicana</i>	0-112	-
	Illinois bundleflower	DEIL	<i>Desmanthus illinoensis</i>	0-112	-
	bractless blazingstar	MENU	<i>Mentzelia nuda</i>	0-56	-
	Nuttall's sensitive-briar	MINU6	<i>Mimosa nuttallii</i>	0-56	-
	yellow puff	NELU2	<i>Neptunia lutea</i>	0-56	-
	evening primrose	OENOT	<i>Oenothera</i>	0-56	-
	groundcherry	PHYSA	<i>Physalis</i>	0-56	-
	scurfpea	PSORA2	<i>Psoralegium</i>	0-56	-
	pitcher sage	SAAZG	<i>Salvia azurea var. grandiflora</i>	0-56	-
	globemallow	SPHAE	<i>Sphaeralcea</i>	0-56	-
	amberique-bean	STHE9	<i>Strophostyles helvola</i>	0-56	-
	white heath aster	SYERE	<i>Symphotrichum ericoides var. ericoides</i>	0-56	-
	prairie spiderwort	TROC	<i>Tradescantia occidentalis</i>	0-56	-
	Texas vervain	VEHA	<i>Verbena halei</i>	0-56	-
	purple poppymallow	CAIN2	<i>Callirhoe involucrata</i>	0-56	-
	yellow sundrops	CASE12	<i>Calylophus serrulatus</i>	0-56	-
	bluestem pricklypoppy	ARAL3	<i>Argemone albiflora</i>	0-56	-
	whitemouth dayflower	COER	<i>Commelina erecta</i>	0-56	-
	prairie clover	DALEA	<i>Dalea</i>	0-56	-
	purple prairie clover	DAPU5	<i>Dalea purpurea</i>	0-56	-
	hymenopappus	HYMEN4	<i>Hymenopappus</i>	0-56	-
	lespedeza	LESPE	<i>Lespedeza</i>	0-56	-
	buckwheat	ERIOG	<i>Eriogonum</i>	0-56	-
	beeblossom	GAURA	<i>Gaura</i>	0-56	-
Shrub/Vine					

6	Shrubs/Vines			112-336	
	plum	PRUNU	<i>Prunus</i>	0-112	-
	skunkbush sumac	RHTR	<i>Rhus trilobata</i>	0-112	-
	gum bully	SILA20	<i>Sideroxylon lanuginosum</i>	0-112	-
	pricklyash	ZANTH	<i>Zanthoxylum</i>	0-112	-
	greenbrier	SMILA2	<i>Smilax</i>	0-56	-
	yucca	YUCCA	<i>Yucca</i>	0-56	-
	catclaw acacia	ACGRG3	<i>Acacia greggii</i> var. <i>greggii</i>	0-56	-
	Christmas cactus	CYLE8	<i>Cylindropuntia leptocaulis</i>	0-56	-
	pricklypear	OPUNT	<i>Opuntia</i>	0-56	-
Tree					
7	Trees			112-336	
	post oak	QUST	<i>Quercus stellata</i>	112-560	-
	American elm	ULAM	<i>Ulmus americana</i>	0-112	-
	sugarberry	CELAL	<i>Celtis laevigata</i> var. <i>laevigata</i>	0-112	-
	netleaf hackberry	CELAR	<i>Celtis laevigata</i> var. <i>reticulata</i>	0-112	-
	honey mesquite	PRGL2	<i>Prosopis glandulosa</i>	0-112	-
	blackjack oak	QUMA3	<i>Quercus marilandica</i>	0-112	-
	ash	FRAXI	<i>Fraxinus</i>	0-56	-
	slippery elm	ULRU	<i>Ulmus rubra</i>	0-56	-
	wingleaf soapberry	SASA4	<i>Sapindus saponaria</i>	0-56	-
	winged elm	ULAL	<i>Ulmus alata</i>	0-56	-

Table 6. Community 1.2 plant community composition

Group	Common Name	Symbol	Scientific Name	Annual Production ()	Foliar Cover (%)
-------	-------------	--------	-----------------	----------------------	------------------

Table 7. Community 1.3 plant community composition

Group	Common Name	Symbol	Scientific Name	Annual Production ()	Foliar Cover (%)
-------	-------------	--------	-----------------	----------------------	------------------

Table 8. Community 2.1 plant community composition

Group	Common Name	Symbol	Scientific Name	Annual Production ()	Foliar Cover (%)
-------	-------------	--------	-----------------	----------------------	------------------

Table 9. Community 3.1 plant community composition

Group	Common Name	Symbol	Scientific Name	Annual Production ()	Foliar Cover (%)
-------	-------------	--------	-----------------	----------------------	------------------

Table 10. Community 4.1 plant community composition

Group	Common Name	Symbol	Scientific Name	Annual Production ()	Foliar Cover (%)
-------	-------------	--------	-----------------	----------------------	------------------

Table 11. Community 4.2 plant community composition

Group	Common Name	Symbol	Scientific Name	Annual Production ()	Foliar Cover (%)
-------	-------------	--------	-----------------	----------------------	------------------

Animal community

Historically, the Loamy Sand site was inhabited permanently and intermittently by a wide variety of mammals, reptiles, and birds. Several historical references and journals written in the 18th and 19th century by explorers, survey parties, and military expeditions refer to herds of bison, wild cattle, wild horses, and antelope roaming freely across the North Central Prairie and adjacent regions. Currently, the site is utilized by deer, quail, dove, numerous species of birds, and a variety of small fur-bearing mammals. Animal species and populations fluctuate as the vegetation cycles through temporary phases and different ecological stages. Livestock grazing should be controlled by implementing grazing management systems that incorporate frequent and timely deferment periods to prevent abusive grazing.

Hydrological functions

When herbaceous vegetation and ground cover are maintained in a healthy and vigorous status, water infiltration into the soil profile is increased significantly, resulting in less runoff. A thick, healthy grass cover also results in improved water quality because it serves as a filter or trap to reduce sediments and pollutants before the water flows offsite.

Recreational uses

These scenic areas offer outdoor activities including photography, shaded picnic areas, bird watching, hiking, camping, horseback riding, and off-road vehicle use. The site offers opportunities for hunting hunting quail, dove, deer, and turkey because of its diverse plant communities, abundance of forbs, and variety of trees and shrubs.

Wood products

Post oak, elms, and hackberry have limited value as fence posts, firewood, specialty products, and crafts.

Other products

None.

Other information

None.

Inventory data references

Vegetation data for this site was obtained from existing Range Site Descriptions, SCS-RANGE -417 Production and Composition Records for Native Grazing Lands, and on-site inventories by the author and local experts including ranchers, natural resource specialists from federal and state agencies, and personnel from cooperating agencies and organizations. A total of 7 SCS-RANGE-417's containing data collected from 3 counties during the period 12/30/1981 to 12/12/1986 were reviewed for this site.

References

. 2021 (Date accessed). **USDA PLANTS Database**. <http://plants.usda.gov>.

Bailey, V. 1905. **Biological Survey of Texas**. North American Fauna 25:1–222.

Other references

Ajilvsgi, Geyata. *Wildflowers of Texas*. Sharer Publishing, Bryan, TX. 1984.

Bachand, Richard. *The American Prairie: Going, Going, Gone?* National Wildlife Federation. Rocky Mountain Natural Resource Center. Boulder, CO. 2001.

Coffey, Chuck R., and Russell Stevens. *Grasses of Southern Oklahoma and North Texas: A Pictorial Guide*. The Samuel Roberts Noble Foundation, Ardmore, OK. 2004

Diggs, George M., Jr., Barney L. Lipscomb, and Robert J. O'Kennon. *Illustrated Flora of North Central Texas*. Botanical Research Institute of Texas. Fort Worth, TX 1999.

Egan, Dave and Evelyn A. Howell. *The Historical Ecology Handbook...A Restorationist's Guide to Reference Ecosystems*. Island Press, Washington, DC. 2001.

Enquist, Marshall. *Wildflowers of the Texas Hill Country*. Lone Star Botanical, Austin, TX. 1987.

Flores, Dan. "Indian Use of Range Resources" presented at 20th Annual Ranch Management Conference. Lubbock, TX, September 30, 1983.

Gould, Frank W., *The Grasses of Texas*. Texas A&M University Press, College Station, TX. 1975.

Hatch, Stephan L., Kancheepuram N. Gandhi, and Larry E. Brown. Checklist of the Vascular Plants of Texas. Texas Agricultural Experiment Station MP-1655. College Station, TX. 1990

Hatch, Stephan L., Jennifer Pluhar. Texas Range Plants. Texas A&M University Press, College Station, TX. 1993.

Kelton, Elmer. "History of Rancher Use of Range Resources" presented at 20th Annual Ranch Management Conference. Lubbock, TX, September 30, 1983.

Ladd, Doug. Tallgrass Prairie Wildflowers. Falcon Press, Helena and Billings, MT. 1995.

Larrabee, Aimee and John Altman. Last Stand of the Tallgrass Prairie. Sterling Publishing Co., New York, NY. 2001.

Merz, Dalton. Personal communication. 9/29/2007.

Nelson, Paul W. The Terrestrial Natural Communities of Missouri. Missouri Department of Natural Resources. 1985.

Packard, Stephen and Cornelia F. Mutel. The Tallgrass Restoration Handbook for Prairies, Savannas, and Woodlands. Island Press, Washington, DC. 1997.

Parker, W.B. Through Unexplored Texas In The Summer and Fall of 1854. The Texas State Historical Commission. Austin, TX 1984

Smith, Jared G. Grazing Problems in the Southwest and How to Meet Them. United States Department of Agriculture Division of Agrostology. Washington, DC. 1899.

Texas Almanac Sesquicentennial Edition 1857-2007. Dallas Morning News. Dallas, TX. 2006.

Tyrl, Ronald J., Terrence G. Bidwell, and Ronald E. Masters. Field Guide to Oklahoma Plants. Oklahoma State University, Stillwater, OK. 2002.

United States Department of Agriculture Natural Resources Conservation Service, National Plant Data Center, Baton Rouge, LA. The PLANTS Database. <http://plants.usda.gov> 2007.

United States Department of Agriculture Natural Resources Conservation Service, Ag Handbook 296. Land Resource Regions and Major Land Resource Areas of the United States, the Caribbean, and the Pacific Basin. 2006.

United States Department of Agriculture Natural Resources Conservation Service, Temple, TX. Loamy Sand Ecological Site Descriptions R078BY082TX and R084BY171TX. 2006.

United States Department of Agriculture Soil Conservation Service, Temple, TX. Production and Composition Record for Native Grazing Lands. SCS-RANGE 417 data from Brown, Eastland, Jack, Stephens, and Young Counties. 1981-1986.

United States Department of Agriculture Soil Conservation Service, Washington, DC. Web Soil Survey <http://websoilsurvey.nrcs.usda.gov/app/>. 2007

United States Department of Agriculture Soil Conservation Service, Temple, TX. Published Soil Surveys: Brown and Mills, Jack, Palo Pinto, Stephens, and Young Counties. Various publication dates.

United States Department of Agriculture Soil Conservation Service, Temple, TX. Range Site Descriptions for the North Central Prairie counties. Various publication dates.

Vines, Robert A. Trees of North Texas. University of Texas Press, Austin, TX. 1982

Weniger, Del. The Explorers' Texas. Eakin Publications. Austin, TX. 1984.

Williams, Gerald W. References On The American Indian Use Of Fire in Ecosystems. United States Department of Agriculture – Forest Service, Washington, DC. 2005.

ACKNOWLEDGEMENTS: I would like to express my thanks and appreciation to the following for their cooperation, assistance, and support in the development of this Ecological Site Description:

Tony Dean, NRCS – Henrietta, TX

Matt Gregory, NRCS – Jacksboro, TX

John Paclik, NRCS – Graham, TX

Richards Ranch – Jacksboro, TX

Reviewers:

Lem Creswell, RMS, NRCS, Weatherford, Texas
Justin Clary, RMS, NRCS, Temple, Texas

Contributors

Dan Caudle, DMC Natural Resources Management, Weatherford Texas

Approval

Bryan Christensen, 9/19/2023

Acknowledgments

Site Development and Testing Plan: Future work, as described in a Project Plan, to validate the information in this Provisional Ecological Site Description is needed. This will include field activities to collect low, medium and high intensity sampling, soil correlations, and analysis of that data. Annual field reviews should be done by soil scientists and vegetation specialists. A final field review, peer review, quality control, and quality assurance reviews of the ESD will be needed to produce the final document. Annual reviews of the Project Plan are to be conducted by the Ecological Site Technical Team.

Rangeland health reference sheet

Interpreting Indicators of Rangeland Health is a qualitative assessment protocol used to determine ecosystem condition based on benchmark characteristics described in the Reference Sheet. A suite of 17 (or more) indicators are typically considered in an assessment. The ecological site(s) representative of an assessment location must be known prior to applying the protocol and must be verified based on soils and climate. Current plant community cannot be used to identify the ecological site.

Author(s)/participant(s)	Lem Creswell, Zone RMS, NRCS, Weatherford, Texas
Contact for lead author	817-596-2865
Date	10/22/2007
Approved by	
Approval date	
Composition (Indicators 10 and 12) based on	Annual Production

Indicators

1. **Number and extent of rills:** None.
-

2. **Presence of water flow patterns:** Deposition or erosion is uncommon during normal rainfall events, but may occur in limited areas during intense rainfall events.

3. **Number and height of erosional pedestals or terracettes:** None.

4. **Bare ground from Ecological Site Description or other studies (rock, litter, lichen, moss, plant canopy are not bare ground):** Expect no more than 10% bare ground scattered randomly throughout the site.

5. **Number of gullies and erosion associated with gullies:** Few rills and very few gullies may occur on sidedrains into perennial and intermittent streams. Gullies and rills should be vegetated and stable.

6. **Extent of wind scoured, blowouts and/or depositional areas:** Uncommon.

7. **Amount of litter movement (describe size and distance expected to travel):** Little or no litter movement or deposition during normal rainfall events.

8. **Soil surface (top few mm) resistance to erosion (stability values are averages - most sites will show a range of values):** Soil surface is moderately susceptible to wind erosion in this MLRA.

9. **Soil surface structure and SOM content (include type of structure and A-horizon color and thickness):** 0-8" thick - weak, fine granular structure. SOM is approximately 1-6%. See Official Series Description for more information.

10. **Effect of community phase composition (relative proportion of different functional groups) and spatial distribution on infiltration and runoff:** The tallgrass/midgrass savanna with abundant forbs, adequate litter, and little bare ground provides for maximum infiltration and negligible runoff.

11. **Presence and thickness of compaction layer (usually none; describe soil profile features which may be mistaken for compaction on this site):** None.

12. **Functional/Structural Groups (list in order of descending dominance by above-ground annual-production or live foliar cover using symbols: >>, >, = to indicate much greater than, greater than, and equal to):**

Dominant: Warm-season tallgrasses >> warm-season midgrasses >

Sub-dominant: Warm-season midgrasses > Trees >

Other: Forbs > Shrubs/Vines > Cool-season grasses

Additional:

13. **Amount of plant mortality and decadence (include which functional groups are expected to show mortality or decadence):**
Perennial grasses will naturally exhibit a minor amount (less than 5%) of senescence and some mortality every year.
-

14. **Average percent litter cover (%) and depth (in):** Litter is dominantly herbaceous.
-

15. **Expected annual annual-production (this is TOTAL above-ground annual-production, not just forage annual-production):**
2000 to 5500 pounds per acre.
-

16. **Potential invasive (including noxious) species (native and non-native). List species which BOTH characterize degraded states and have the potential to become a dominant or co-dominant species on the ecological site if their future establishment and growth is not actively controlled by management interventions. Species that become dominant for only one to several years (e.g., short-term response to drought or wildfire) are not invasive plants. Note that unlike other indicators, we are describing what is NOT expected in the reference state for the ecological site:** Mesquite, pricklypear, yucca, tasajillo, pricklyash, lotebush, bermudagrass, johnsongrass, King Ranch bluestem, weeping lovegrass, annual broomweed
-

17. **Perennial plant reproductive capability:** All perennial species should be capable of reproducing every year unless disrupted by extended drought, overgrazing, wildfire, insect damage, or other events occurring immediately prior to, or during the reproductive phase.
-