

Ecological site R083AY021TX Sandy

Last updated: 9/19/2023
Accessed: 04/17/2026

General information

Provisional. A provisional ecological site description has undergone quality control and quality assurance review. It contains a working state and transition model and enough information to identify the ecological site.

MLRA notes

Major Land Resource Area (MLRA): 083A–Northern Rio Grande Plain

This area is entirely in Texas and south of San Antonio. It makes up about 11,115 square miles (28,805 square kilometers). The towns of Uvalde, Cotulla, and Hondo are in the western part of the area, and Beeville, Goliad, and Kenedy are in the eastern part. The town of Alice is just outside the southern edge of the area. Interstate Highways 35 and 37 cross this area. This area is comprised of inland, dissected coastal plains.

Classification relationships

USDA-Natural Resources Conservation Service, 2006. -Major Land Resource Area (MLRA) 83A

Ecological site concept

The Sandy ecological sites are very deep and are moderately well to excessively drained. Soils typically have a thick sandy surface from 40 to 80 inches with a loamy or clayey subsoil.

Associated sites

R083AY023TX	Sandy Loam
R083AY024TX	Tight Sandy Loam
R083AY022TX	Loamy Sand

Similar sites

R083CY021TX	Sandy
R083EY021TX	Sandy

Table 1. Dominant plant species

Tree	(1) <i>Quercus virginiana</i>
Shrub	Not specified
Herbaceous	(1) <i>Schizachyrium scoparium</i> (2) <i>Sorghastrum nutans</i>

Physiographic features

The Sandy ecological sites are found on nearly level to gently sloping stream terraces of the Coastal Plains. The soils developed in fine sands and loamy fine sands, presumably of recent eolian origin over loamy sediments. The sediments were formed from the Carrizo Sand geologic formation. Slopes range from 0 to 5 percent. Elevation ranges from 200 to 1,000 feet. This area is comprised of inland, dissected coastal plains.

Table 2. Representative physiographic features

Landforms	(1) Coastal plain > Stream terrace
Runoff class	Negligible to medium
Elevation	20 – 310 m
Slope	0 – 10 %
Aspect	Aspect is not a significant factor

Climatic features

MLRA 83A is subtropical, subhumid on the western boundary and subtropical humid on the eastern boundary. Winters are dry and mild and the summers are hot and humid. Tropical maritime air masses predominate throughout spring, summer, and fall. Modified polar air masses exert considerable influence during winter, creating a continental climate characterized by large variations in temperature. Average precipitation for MLRA 83A is 20 inches on the western boundary and 35 inches on the eastern boundary. Peak rainfall, because of rain showers, occurs late in spring and a secondary peak occurs early in fall. Heavy thunderstorm activities increase in April, May, and June. July is hot and dry with little weather variations. Rainfall increases again in late August and September as tropical disturbances increase and become more frequent. Tropical air masses from the Gulf of Mexico dominate during the spring, summer, and fall. Prevailing winds are southerly to southeasterly throughout the year except in December when winds are predominately northerly.

Table 3 Representative climatic features

Frost-free period (characteristic range)	230-250 days
Freeze-free period (characteristic range)	260-370 days
Precipitation total (characteristic range)	640-810 mm
Frost-free period (actual range)	220-260 days

Freeze-free period (actual range)	250-370 days
Precipitation total (actual range)	610-940 mm
Frost-free period (average)	240 days
Freeze-free period (average)	310 days
Precipitation total (average)	740 mm

- (1) BEEVILLE 5 NE [USC00410639], Beeville, TX
- (2) CHEAPSIDE [USC00411671], Gonzales, TX
- (3) CUERO [USC00412173], Cuero, TX
- (4) GOLIAD [USC00413618], Goliad, TX
- (5) NIXON [USC00416368], Stockdale, TX
- (6) CARRIZO SPRINGS 3W [USC00411486], Carrizo Springs, TX
- (7) FOWLERTON [USC00413299], Fowlerton, TX
- (8) HONDO [USC00414254], Hondo, TX
- (9) KARNES CITY 2N [USC00414696], Karnes City, TX
- (10) PEARSALL [USC00416879], Pearsall, TX
- (11) CHARLOTTE 5 NNW [USC00411663], Charlotte, TX
- (12) MATHIS 4 SSW [USC00415661], Mathis, TX
- (13) TILDEN 4 SSE [USC00419031], Tilden, TX
- (14) UVALDE 3 SW [USC00419268], Uvalde, TX
- (15) CROSS [USC00412125], Tilden, TX
- (16) DILLEY [USC00412458], Dilley, TX
- (17) FLORESVILLE [USC00413201], Floresville, TX
- (18) LYTTLE 3W [USC00415454], Natalia, TX
- (19) PLEASANTON [USC00417111], Pleasanton, TX
- (20) HONDO MUNI AP [USW00012962], Hondo, TX
- (21) CALLIHAM [USC00411337], Calliham, TX

Influencing water features

Many sites are somewhat excessively drained in the surface until the water contacts the argillic, where it becomes a very slow to moderately permeable layer. Some soils may exhibit a perched water table after very heavy rains for a short duration.

Wetland description

N/A.

Soil features

The soils in this site are very deep, moderate to excessively drained, with moderate or moderately slow permeability in the subsoil. Ochric epipedons range from 40 to 80 inches over a loamy subsoil. Other features include sandy surface textures, little to no salinity or sodicity, and moderately acid to slightly alkaline soil reaction. Soil series correlated to this site include: Antosa, Bobillo, Nusil, Rhymes, and Ruiz.

Table 4. Representative soil features

Parent material	(1) Alluvium – sandstone
-----------------	--------------------------

Surface texture	(1) Fine sand (2) Loamy fine sand
Family particle size	(1) Loamy
Drainage class	Moderately well drained to somewhat excessively drained
Permeability class	Very slow to moderately rapid
Surface fragment cover <=3"	Not specified
Surface fragment cover >3"	Not specified
Available water capacity (0-101.6cm)	5.08 – 10.16 cm
Calcium carbonate equivalent (0-101.6cm)	0 – 10 %
Electrical conductivity (0-101.6cm)	Not specified
Sodium adsorption ratio (0-101.6cm)	0 – 10
Soil reaction (1:1 water) (0-101.6cm)	5.6 – 8.4
Subsurface fragment volume <=3" (Depth not specified)	Not specified
Subsurface fragment volume >3" (Depth not specified)	0 – 10 %

Ecological dynamics

The plant communities of this site are dynamic and community composition may vary dramatically in annual rainfall, grazing, and fire. The site is subject to extreme variation in rainfall. During the years 1900 to 1983, 36 percent were drought years and 34 percent were wet years. During dry periods the amount of bare ground increases. Bare ground may predominate during droughts. Shortgrasses such as hairy grama (*Bouteloua hirsuta*), thin paspalum (*Paspalum setaceum*), fringed signalgrass (*Brachiaria ciliatissima*), red lovegrass (*Eragrostis secundiflora*), sandbur (*Cenchrus* spp.), and forbs increase in abundance at the expense of the taller grasses. During wet years, tallgrasses such as big bluestem (*Andropogon gerardii*) increase in importance. The shortgrasses and forbs occur as an understory component forming a multi-layered community.

Early explorers provide some insight into the general landscape but most lack site-specific information. Their observations do provide useful information into the general aspect of the landscape. In some cases the exact position of the explorers can be determined. In 1821, Stephen F. Austin crossed the San Antonio River to Cabeza Creek in Goliad County. He observed the area to be sandy in places and there was not too much mesquite and underbrush. In 1846, Sitgreaves crossed Karnes County and stated that the whole distance was over dry, sandy, rolling prairie covered with mesquite. He reported timber was more abundant. Ponce de Leon in 1689 observed in the common county corners of La Salle, Frio, Atascosa and McMullen counties, describing the country as level, with fine pasturage, very pleasant glades, and occasionally, little mottes of oak. Overall the upland country was described with small amounts of brush or mottes interspersed in the prairie. The increase of brush generally coincides with settlement.

Historically, fire was an important factor in the ecology of this site. Native Americans set periodic fires for hunting and reducing insects. Fires reduced woody plant cover, kept oak mottes scattered and isolated, and maintained the open stretches of grassland witnessed by Berlandier. Wildfires are common on this site at present. White-tailed deer (*Odocoileus virginianus*) and pronghorns (*Antilocapra americana*) were significant herbivores on this site at the time of colonization by Europeans. The extent to which bison (*Bos bison*) utilized the site is uncertain. The reports of bison were not nearly as abundant as farther north in the southern plains region.

The reference plant community is a grassland with scattered live oak mottes and occasional mesquite trees. Little bluestem (*Schizachyrium scoparium*) was the prevailing dominant species. Other important associated grasses include big bluestem, brownseed paspalum (*Paspalum plicatulum*), Indiangrass (*Sorghastrum* spp.), switchgrass (*Panicum virgatum*), tanglehead (*Heteropogon contortus*), and thin paspalum. The reference plant community supports a diverse understory community of perennial legumes and other forbs.

Continued overuse by livestock results in a decline of little bluestem and other perennial grasses and an increase in forbs, particularly camphor daisy (*Rayjacksonia phyllocephala*), partridgepea (*Chamaecrista fasciculata*) and Crotons (*Croton* spp.). Pan-American balsamscale, three-awns (*Aristida* spp.), and thin paspalum increase in abundance with heavy grazing but decline on severely grazed rangeland. On severely grazed rangeland, little bluestem is virtually absent. Sandbur, fringed signalgrass, red lovegrass, camphor daisy, and other forbs dominate severely grazed sites. Severe overuse results in a large amount of bare ground. The oak colonies can become thicketed, and take on a low stature with high stem density rather than forming large, single-trunked trees. Mesquite increases once established. After the mesquites reach sufficient size, understory shrubs including granjeno (*Celtis pallida*), brasil (*Condalia hookeri*), and lime prickly-ash (*Zanthoxylum fagara*) establish beneath them, forming brush mottes.

State and transition model

Figure 7. STM

Additional community tables

Table 5. Community 1.1 plant community composition

Group	Common Name	Symbol	Scientific Name	Annual Production ()	Foliar Cover (%)
Grass/Grasslike					
1	Tallgrass			1121-2522	
	little bluestem	SCSC	<i>Schizachyrium scoparium</i>	1121-2522	–
2	Midgrass			112-336	
	brownseed paspalum	PAPL3	<i>Paspalum plicatulum</i>	112-336	–
3	Tallgrasses			224-560	
	Indiangrass	SONU2	<i>Sorghastrum nutans</i>	112-392	–
	crinkleawn grass	TRACH2	<i>Trachypogon</i>	112-392	–
	switchgrass	PAVI2	<i>Panicum virgatum</i>	112-336	–
4	Midgrass			112-448	
	tanglehead	HECO10	<i>Heteropogon contortus</i>	112-448	–
5	Midgrass			112-252	
	fringed signalgrass	URCI	<i>Urochloa ciliatissima</i>	112-252	–

6	Midgrasses			112-252	
	balsamscale grass	ELION	<i>Elionurus</i>	56-140	-
	purple dropseed	SPPU3	<i>Sporobolus purpurascens</i>	56-140	-
	Texasgrass	VAMU	<i>Vaseyochloa multinervosa</i>	56-140	-
7	Shortgrasses			112-224	
	Wright's threeawn	ARPUW	<i>Aristida purpurea var. wrightii</i>	56-112	-
	hooded windmill grass	CHCU2	<i>Chloris cucullata</i>	56-112	-
8	Shortgrasses			224-448	
	sand crabgrass	DIAR7	<i>Digitaria arenicola</i>	56-112	-
	fall witchgrass	DICO6	<i>Digitaria cognata</i>	56-112	-
	gulfdune paspalum	PAMO4	<i>Paspalum monostachyum</i>	56-112	-
Forb					
9	Forbs			67-151	
	Texas bullnettle	CNTE	<i>Cnidocolus texanus</i>	28-84	-
	coastal indigo	INMI	<i>Indigofera miniata</i>	28-84	-
	dotted blazing star	LIPU	<i>Liatris punctata</i>	28-84	-
	sensitive plant	MIMOS	<i>Mimosa</i>	28-84	-
	snoutbean	RHYNC2	<i>Rhynchosia</i>	28-84	-
10	Forbs			45-129	
	Forb, annual	2FA	<i>Forb, annual</i>	0-112	-
	partridge pea	CHFA2	<i>Chamaecrista fasciculata</i>	28-84	-
	croton	CROTO	<i>Croton</i>	28-84	-
	snakecotton	FROEL	<i>Froelichia</i>	28-84	-
	lantana	LANTA	<i>Lantana</i>	28-84	-
	beebalm	MONAR	<i>Monarda</i>	28-84	-
Shrub/Vine					
11	Shrubs			0-280	
	mesquite	PROSO	<i>Prosopis</i>	0-280	-
Tree					
12	Trees			112-280	
	live oak	QUVI	<i>Quercus virginiana</i>	112-280	-

Table 6. Community 1.2 plant community composition

Group	Common Name	Symbol	Scientific Name	Annual Production ()	Foliar Cover (%)
-------	-------------	--------	-----------------	----------------------	------------------

Table 7. Community 2.1 plant community composition

Group	Common Name	Symbol	Scientific Name	Annual Production ()	Foliar Cover (%)
-------	-------------	--------	-----------------	----------------------	------------------

Table 8. Community 2.2 plant community composition

Group	Common Name	Symbol	Scientific Name	Annual Production ()	Foliar Cover (%)
-------	-------------	--------	-----------------	----------------------	------------------

Animal community

As a historic tall/midgrass prairie, this site was occupied by bison, antelope, deer, quail, turkey, and dove. This site was also used by many species of grassland songbirds, migratory waterfowl, and coyotes. This site now provides forage for livestock and is still used by quail, dove, migratory waterfowl, grassland birds, coyotes, and deer. Feral hogs (*Sus scrofa*) can be found on most ecological sites in Texas. Damage caused by feral hogs each year includes, crop damage by rutting up crops, destroyed fences, livestock watering areas, and predation on native wildlife, and ground-nesting birds. Feral hogs have few natural predators, thus allowing their population to grow to high numbers. Wildlife habitat is a complex of many different plant communities and ecological sites across the landscape. Most

animals use the landscape differently to find food, shelter, protection, and mates. Working on a conservation plan for the whole property, with a local professional, will help managers make the decisions that allow them to realize their goals for wildlife and livestock. Grassland State (1): This state provides the maximum amount of forage for livestock such as cattle. It is also utilized by deer, quail and other birds as a source of food. When a site is in the reference plant community phase (1.1) it will also be used by some birds for nesting, if other habitat requirements like thermal and escape cover are near. Shrubland State (2): This state can be maintained to meet the habitat requirements of cattle and wildlife. Land managers can find a balance that meets their goals and allows them flexibility to manage for livestock and wildlife. Forbs for deer and birds like quail will be more plentiful in this state. There will also be more trees and shrubs to provide thermal and escape cover for birds as well as cover for deer. This rating system provides general guidance as to animal preference for plant species. It also indicates possible competition between kinds of herbivores for various plants. Grazing preference changes from time to time, especially between seasons, and between animal kinds and classes. Grazing preference does not necessarily reflect the ecological status of the plant within the plant community. For wildlife, plant preferences for food and plant suitability for cover are rated. Refer to habitat guides for a more complete description of a species habitat needs.

Hydrological functions

Water infiltration is rapid in the fine sands of the site. Therefore, runoff and soil erosion from water is seldom a problem.

Recreational uses

Hunting, birdwatching, and eco-tourism are common uses.

Inventory data references

Information presented was derived from the revised Range Site, literature, limited NRCS clipping data (417s), field observations, and personal contacts with range-trained personnel.

Other references

- AgriLife. 2009. Managing Feral Hogs Not a One-shot Endeavor. AgNews, April 23, 2009. <http://agnews.tamu.edu/showstory.php?id=903>.
- Archer, S. 1995. Herbivore mediation of grass-woody plant interactions. *Tropical Grasslands*, 29:218-235.
- Archer, S. 1995. Tree-grass dynamics in a Prosopis-thornscrub savanna parkland: reconstructing the past and predicting the future. *Ecoscience*, 2:83-99.
- Archer, S. 1994. Woody plant encroachment into southwestern grasslands and savannas: rates, patterns and proximate causes. *Ecological implications of livestock herbivory in the West*, 13-68.
- Archer, S. and F. E. Smeins. 1991. Ecosystem-level Processes. In *Grazing Management: An Ecological Perspective*. Edited by R.K. Heischmidt and J.W. Stuth. Timber Press, Portland, OR.
- Baen, J. S. 1997. The growing importance and value implications of recreational hunting leases to agricultural land investors. *Journal of Real Estate Research*, 14:399-414.
- Bailey, V. 1905. North American Fauna No. 25: Biological Survey of Texas. United States Department of Agriculture Biological Survey. Government Printing Office, Washington D. C.
- Bestelmeyer, B. T., J.R. Brown, K. M. Havstad, R. Alexander, G. Chavez, and J. E. Herrick. 2003. Development and use of state-and-transition models for rangelands. *Journal of Range Management*, 56(2):114-126.
- Box, T. W. 1960. Herbage production on four range plant communities in South Texas. *Journal of Range Management*, 13:72-76.
- Briske, B B, B. T. Bestelmeyer, T. K. Stringham, and P. L. Shaver. 2008. Recommendations for development of resilience-based State-and-Transition Models. *Rangeland Ecology and Management*, 61:359-367.
- Brown, J. R. and S. Archer. 1999. Shrub invasion of grassland: recruitment is continuous and not regulated by herbaceous biomass or density. *Ecology*, 80(7):2385-2396.
- Diamond, D. D. and T. E. Fulbright. 1990. Contemporary plant communities of upland grasslands of the Coastal Sand Plain, Texas. *Southwestern Naturalist*, 35:385-392.
- Dillehay T. 1974. Late quaternary bison population changes on the Southern Plains. *Plains Anthropologist*, 19:180-96.
- Edward, D. B. 1836. The history of Texas; or, the immigrants, farmers, and politicians guide to the character, climate, soil and production

- of that country. Geographically arranged from personal observation and experience. J. A. James and Co., Cincinnati, OH.
- Everitt, J. H., D. L. Drawe, and R. I. Leonard. 2002. *Trees, Shrubs, and Cacti of South Texas*. Texas Tech University Press, Lubbock, TX.
- Everitt, J. H., D. L. Drawe, and R. I. Leonard. 1999. *Field Guide to the Broad-Leaved Herbaceous Plants of South Texas*. Texas Tech University Press, Lubbock, TX.
- Foster, J. H. 1917. Pre-settlement fire frequency regions of the United States: a first approximation. *Tall Timbers Fire Ecology Conference Proceedings* No. 20.
- Foster, W. C., ed. 1998. *The La Salle Expedition to Texas: The Journal of Henry Joutel, 1684-1687*. Texas State Historical Association, Austin, TX.
- Frost, C. C. 1995. Presettlement fire regimes in southeastern marshes, peatlands, and swamps. In: *Proceedings, 19th Tall Timbers fire ecology conference*, 39-60. Tall Timbers Research Station, Tallahassee, FL.
- Fulbright, T. E. and S. L. Beasom. 1987. Long-term effects of mechanical treatment on white-tailed deer browse. *Wildlife Society Bulletin*, 15:560-564.
- Fulbright, T. E., J. A. Ortega-Santos, A. Lozano-Cavazos, and L. E. Ramirez-Yanez. 2006. Establishing vegetation on migrating inland sand dunes in Texas. *Rangeland Ecology and Management*, 59:549-556.
- Fulbright, T. E., D. D. Diamond, J. Rappole, and J. Norwine. *The Coastal Sand Plain of Southern Texas*. *Rangelands*, 12:337-340.
- Gould, F. W. 1975. *The Grasses of Texas*. Texas A&M University Press, College Station, TX.
- Grace, J. B., L. K. Allain, H. Q. Baldwin, A. G. Billock, W. R. Eddleman, A. M. Given, C. W. Jeske, and R. Moss. 2005. Effects of prescribed fire in the coastal prairies of Texas. *USGS Open File Report 2005-1287*.
- Hamilton, W. and D. Ueckert. 2005. Rangeland Woody Plant Control: Past, Present, and Future. In: *Brush Management: Past, Present, and Future*, 3-16. Texas A&M University Press, College Station, TX.
- Hansmire, J. A., D. L. Drawe, B. B. Wester and C.M. Britton. 1988. Effect of winter burns on forbs and grasses of the Texas Coastal Prairie. *The Southwestern Naturalist*, 33(3):333-338.
- Heitschmidt R. K., Stuth J. W., eds. 1991. *Grazing management: an ecological perspective*. Timberline Press, Portland, OR.
- Inglis, J. M. 1964. *A history of vegetation of the Rio Grande Plains*. Texas Parks and Wildlife Department Bulletin No. 45, Austin, TX.
- Kneuper, C. L., C. B. Scott, and W. E. Pinchak. 2003. Consumption and dispersion of mesquite seeds by ruminants. *Journal of Range Management*, 56:255-259.
- Kramp, B., R. Ansley, and D. Jones. 1998. Effect of prescribed fire on mesquite seedlings. *Texas Tech University Research Highlights - Range, Wildlife and Fisheries Management*, 29:13.
- Le Houerou, H. N. and J. Norwine. 1988. The ecoclimatology of South Texas. In *Arid lands: today and tomorrow*. Edited by E. E. Whitehead, C. F. Hutchinson, B. N. Timmesman, and R. G. Varady, 417-444. Westview Press, Boulder, CO.
- Lehman, V. W. 1965. Fire in the range of Attwater's prairie chicken. *Tall Timbers Fire Ecology Conference*, 4:127-143.
- Lehman, V. W. 1969. *Forgotten Legions: Sheep in the Rio Grande Plain of Texas*. Texas Western Press, El Paso, TX.
- Mann, C. 2004. 1491. *New Revelations of the Americas before Columbus*. Vintage Books, New York City, NY.
- Mapston, M. E. 2009. *Feral Hogs in Texas*. Rep. Texas Cooperative Extension. 23 Apr. 2009 <http://icwdm.org/Publications/pdf/Feral%20Pig/Txferalhogs.pdf>
- McClendon, T. 1991. Preliminary description of the vegetation of South Texas exclusive of the Coastal Saline Zones. *Texas Journal of Science*, 43:13-32.
- McGinty A., D. N. Ueckert. 2001. The Brush Busters success story. *Rangelands*, 23:3-8.
- McLendon, T. 1991. Preliminary description of the vegetation of south Texas exclusive of coastal saline zones. *Texas Journal of Science*, 43:13-32.

- Norwine, J. 1978. Twentieth-century semiarid climates and climatic fluctuations in Texas and northeastern Mexico. *Journal of Arid Environments*, 1:313-325.
- Norwine, J. and R. Bingham. 1986. Frequency and severity of droughts in South Texas: 1900-1983, 1-17. In *Livestock and wildlife management during drought*. Edited by R. D. Brown. Caesar Kleberg Wildlife Research Institute, Kingsville, TX.
- Olmsted, F. L. 1857. *A journey through Texas, or a saddle trip on the Southwest frontier: with a statistical appendix*. Dix, Edwards, and co., New York, London.
- Prichard, D. 1998. *A User Guide to Assessing Proper Functioning Condition and the Supporting Science for Lentic Areas*. Bureau of Land Management. National Applied Resource Sciences Center, CO.
- Rappole, J. H. and G. W. Blacklock. 1994. *A field guide: Birds of Texas*. Texas A&M University Press, College Station, TX.
- Rhyne, M. Z. 1998. Optimization of wildlife and recreation earnings for private landowners. M. S. Thesis, Texas A&M University-Kingsville, Kingsville, TX.
- Schindler, J. R. and T. E. Fulbright. 2003. Roller chopping effects on Tamaulipan scrub community composition. *Journal of Range Management*, 56:585-590.
- Schmidley, D. J. 1983. *Texas mammals east of the Balcones Fault zone*. Texas A&M University Press, College Station, TX.
- Scifres C. J., W. T. Hamilton, J. R. Conner, J. M. Inglis, and G. A. Rasmussen. 1985. *Integrated Brush Management Systems for South Texas: Development and Implementation*. Texas Agricultural Experiment Station, College Station, TX.
- Scifres, C. J. and W. T. Hamilton. 1993. *Prescribed burning for brushland management: the South Texas example*. Texas A&M Press, College Station, TX.
- Scifres, C. J. 1975. Systems for improving McCartney rose infested coastal prairie rangeland. *Texas Agricultural Experiment Station Bulletin MP 1225*.
- Smeins, F. E., S. Fuhlendorf, and C. Taylor, Jr. 1997. Environmental and Land Use Changes: A Long Term Perspective. In *Juniper Symposium*, 1-21. Texas Agricultural Experiment Station.
- Smeins, F. E., D. D. Diamond, and W. Hanselka. 1991. Coastal prairie, 269-290. In *Ecosystems of the World: Natural Grasslands*. Edited by R. T. Coupland. Elsevier Press, Amsterdam, Netherlands.
- Soil Survey Staff, Natural Resources Conservation Service, United States Department of Agriculture. *Soil Survey Geographic (SSURGO) Database*.
- Snyder, R. A. and C. L. Boss. 2002. Recovery and stability in barrier island plant communities. *Journal of Coastal Research*, 18:530-536.
- Stiles, H. R., ed. 1906. *Joutel's journal of La Salle's last voyage, 1686-1687*. Joseph McDonough, Albany, NY.
- Stringham, T. K., W. C. Krueger, and P. L. Shaver. 2001. State and transition modeling: and ecological process approach. *Journal of Range Management*, 56(2):106-113.
- Texas A&M Research and Extension Center. 2000. *Native Plants of South Texas* <http://uvalde.tamu.edu/herbarium/index.html>.
- Texas Agriculture Experiment Station. 2007. Benny Simpson's Texas Native Trees <http://aggie-horticulture.tamu.edu/ornamentals/natives/>.
- Texas Parks and Wildlife Department. 2007. *List of White-tailed Deer Browse and Ratings. District 8*.
- Tharp, B. C. 1926. Structure of Texas Vegetation east of the 98th meridian. *Bulletin 2606*. University of Texas, Austin. TX.
- Thurow, T. L. 1991. Hydrology and Erosion. In: *Grazing Management: An Ecological Perspective*. Edited by R.K. Heitschmidt and J.W. Stuth. Timber Press, Portland, OR.
- Urbatsch, L. 2000. Chinese tallow tree (*Triadica sebifera* (L.) Small). *USDA-NRCS Plant Guide*.
- USDA-NRCS Plant Database. 2018. <https://plants.usda.gov/>.
- Van't Hul, J. T., R. S. Lutz and N. E. Mathews. 1997. Impact of prescribed burning on vegetation and bird abundance on Matagorda Island, Texas. *Journal of Range Management*, 50:346-360.

Vines, R. A. 1984. Trees of Central Texas. University of Texas Press, Austin, TX.

Wade, D. D., B. L. Brock, P. H. Brose, J. B. Grace, G. A. Hoch, and W. A. Patterson III. 2000. Fire in Eastern ecosystems. In Wildland fire in ecosystems: effects of fire on flora. Edited by J. K. Brown and J. Kaplers. United States Forest Service, Rocky Mountain Research Station, Ogden, UT.

Weltz, M. A. and W. H. Blackburn. 1995. Water budget for south Texas rangelands. Journal of Range Management, 48:45-52.

Whittaker, R. H., L. E. Gilbert, and J. H. Connell. 1979. Analysis of a two-phase pattern in a mesquite grassland, Texas. Journal of Ecology, 67:935-52.

Wright, B. D., R. K. Lyons, J. C. Cathey, and S. Cooper. 2002. White-tailed deer browse preferences for South Texas and the Edwards Plateau. Texas Cooperative Extension Bulletin B-6130.

Wright, H.A. and A.W. Bailey. 1982. Fire Ecology: United States and Southern Canada. John Wiley & Sons, Inc., Hoboken, NJ.

Approval

Bryan Christensen, 9/19/2023

Acknowledgments

Reviewers and Technical Contributors: Jason Hohlt, RMS, NRCS, Kingsville, Texas Vivian Garcia, RMS, NRCS, Corpus Christi, Texas Shanna Dunn, RSS, NRCS, Corpus Christi, Texas Mark Moseley, RMS, NRCS, Boerne, Texas Justin Clary, RMS, NRCS, Temple, Texas

Rangeland health reference sheet

Interpreting Indicators of Rangeland Health is a qualitative assessment protocol used to determine ecosystem condition based on benchmark characteristics described in the Reference Sheet. A suite of 17 (or more) indicators are typically considered in an assessment. The ecological site(s) representative of an assessment location must be known prior to applying the protocol and must be verified based on soils and climate. Current plant community cannot be used to identify the ecological site.

Author(s)/participant(s)	
Contact for lead author	
Date	04/17/2026
Approved by	
Approval date	
Composition (Indicators 10 and 12) based on	Annual Production

Indicators

1. Number and extent of rills:

2. Presence of water flow patterns:

3. Number and height of erosional pedestals or terracettes:

4. Bare ground from Ecological Site Description or other studies (rock, litter, lichen, moss, plant canopy are not bare ground):

5. Number of gullies and erosion associated with gullies:

6. Extent of wind scoured, blowouts and/or depositional areas:

7. Amount of litter movement (describe size and distance expected to travel):

8. Soil surface (top few mm) resistance to erosion (stability values are averages - most sites will show a range of values):

9. Soil surface structure and SOM content (include type of structure and A-horizon color and thickness):

10. Effect of community phase composition (relative proportion of different functional groups) and spatial distribution on infiltration and runoff:

11. Presence and thickness of compaction layer (usually none; describe soil profile features which may be mistaken for compaction on this site):

12. Functional/Structural Groups (list in order of descending dominance by above-ground annual-production or live foliar cover using symbols: >>, >, = to indicate much greater than, greater than, and equal to):

Dominant:

Sub-dominant:

Other:

Additional:

13. Amount of plant mortality and decadence (include which functional groups are expected to show mortality or decadence):

14. Average percent litter cover (%) and depth (in):

15. Expected annual annual-production (this is TOTAL above-ground annual-production, not just forage annual-production):

16. Potential invasive (including noxious) species (native and non-native). List species which BOTH characterize degraded states and have the potential to become a dominant or co-dominant species on the ecological site if their future establishment and growth is not actively controlled by management interventions. Species that become dominant for only one to several years (e.g., short-term response to drought or wildfire) are not invasive plants. Note that unlike other indicators, we are describing what is NOT expected in the reference state for the ecological site:

17. Perennial plant reproductive capability:
